

INTEGRATING NITROGEN FERTILIZER STRATEGIES, MODELS, AND COVER CROPS
TO OPTIMIZE CORN PRODUCTION AND SOIL HEALTH

By

Jeffrey Alan Rutan

A DISSERTATION

Submitted to
Michigan State University
in partial fulfillment of the requirements
for the degree of

Crop and Soil Sciences – Doctor of Philosophy

2017

ProQuest Number: 10687019

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



ProQuest 10687019

Published by ProQuest LLC (2017). Copyright of the Dissertation is held by the Author.

All rights reserved.

This work is protected against unauthorized copying under Title 17, United States Code
Microform Edition © ProQuest LLC.

ProQuest LLC.
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106 – 1346

ABSTRACT

INTEGRATING NITROGEN FERTILIZER STRATEGIES, MODELS, AND COVER CROPS TO OPTIMIZE CORN PRODUCTION AND SOIL HEALTH

By

Jeffrey Alan Rutan

Degradation of Great Lakes water quality and weather variability emphasize improved nitrogen (N) management strategies that synchronize N availability with corn (*Zea mays* L.) N uptake. Field studies were conducted at either Lansing or Richville, Michigan from 2014 to 2016 and included investigations of cover crops (radish [*Raphanus sativus* (L.) var. The Buster], forage oats [*Avena sativa* (L.) var. Magnum], or no cover crop), and N management strategies (starter N as in-furrow [IF] or sub-surface banded 5-cm beside and 5-cm below the seed furrow [5x5] followed by sidedress [SD] at V4, V10, or split 50/50, pre-plant incorporated N with or without enhanced efficiency N, poultry litter [PL] followed by V11 SD, and zero N). Assessments included cover crop biomass quantity and quality, soil NO₃-N levels, corn grain production and profitability, soil basal CO₂ respiration, and soil bacterial community composition (16S rRNA sequencing). Radish and oat cover crops (CC) seeded in August reduced autumn soil NO₃-N levels 78 – 84% from the no CC control. The following year CC residues reduced N availability to a corn cash crop and reduced effectiveness of 5x5 starter N to maintain yield potential when full SD was applied at V11. Sidedress N delayed from V4 to V11 resulted in similar or reduced yield and profitability. Soil bacterial diversity at R1 was inversely proportional to corn grain yield and suggested bacterial relative abundance may increase indication of yield rather than diversity.

Copyright by
JEFFREY ALAN RUTAN
2017

Dedicated to my family and friends who have provided encouragement along the way.

ACKNOWLEDGEMENTS

My experience in the soil fertility and nutrient management program at Michigan State University has been eclectic and rewarding. First I would like to thank my major professor, Dr. Kurt Steinke. I cannot quantify how much I have learned from you. Thank you for sharing your knowledge and guidance as well as your experience, which will serve me well. Thank you for providing great opportunities to learn, network with peers, and apply myself abroad. Education goes far beyond the classroom, and experience by doing and being involved will provide an invaluable knowledge base. Thank you as well to committee members Christina DiFonzo, Karen Renner, and Noah Rosenzweig for dissertation advice and suggestions.

Thank you to the MSU soil fertility field technician Andy Chomas for your assistance and guidance in my field studies. As a 'hybrid' grower and research investigator, your technical skills and knowledge are an asset to the soil fertility research program. Thank you to MSU Agronomy Farm personnel Tom Galecka for assistance with control applications to field studies, and sharing your knowledge as a grower. Thank you to Bill Widdicombe for your assistance and guidance with corn harvest each year. Field studies for this dissertation were also conducted offsite. I would like to thank Paul Horny and Dennis Fleischmann of the MSU Saginaw Valley Research and Extension Center for facilitating field research off campus.

Thank you to the office staff of the Plant, Soil and Microbial Sciences dept. Thank you as well to Cal Bricker for software technical assistance and support. Cal, your efforts were invaluable to keep mine and other student, staff, and faculty computers operating normally. Thank you to Zach Noel of the MSU Chilvers lab for assistance with microbial analyses. Thank you to fellow graduate students in the soil fertility program, current and past, Mike Swoish, Chris Bauer, Dan Quinn, and Taylor Purucker for field assistance and support. Your friendship is what helps to make a graduate program a memorable experience. Thank you also to all the undergraduates who have assisted with various aspects of my MSU field studies.

Thank you to my family and friends Keith, Susan, Amanda, and Markah for your love and continued support during my graduate studies at MSU. My degree and so many things would not have been possible without all of you. Finally, I thank God for providing diligence, perseverance, guidance, and spiritual growth. For of Him and through Him and to Him are all things, to whom be glory forever. Amen. (Romans 11:36).

TABLE OF CONTENTS

LIST OF TABLES.....	ix
CHAPTER 1	
LITERATURE REVIEW	1
Michigan Corn Production.....	1
Tissue Nitrogen.....	1
Climate Variability.....	2
Nitrogen Use Efficiency	2
Corn Response to N Fertilizer.....	3
4R Nutrient Management.....	4
Corn N application rates	5
Maximum return to nitrogen.....	6
Illinois soil N test and Michigan presidedress soil nitrate test.....	7
Corn N application timing	8
Corn N fertilizer source	10
Corn starter N fertilizer placement	11
In-furrow	12
Subsurface band.....	13
Corn Response to Starter N.....	13
Corn Nutrition Status	16
Chlorophyll meter	16
NDVI.....	17
Basal cornstalk nitrate test	17
Cover Crops and Soil N Dynamics.....	18
Radish cover crops.....	19
Oat cover crops	20
Plant, Soil, and Microbial Community Interactions	21
Effect of Plant Diversity on Soil Microbial Communities.....	22
Influence of Soil Properties on Microbial Community Composition.....	23
Rhizosphere Soil	24
Plant Growth Promoting Rhizobacteria	25
LITERATURE CITED	27
CHAPTER 2	
CORN GROWTH AND YIELD RESPONSES TO PRE-PLANT AND IN-SEASON	
NITROGEN COMBINATIONS	40
Abstract.....	40
Introduction.....	41
Materials and Methods.....	46
Results and Discussion	49
Growing conditions.....	49
Corn plant density	49

Grain yield	50
Profitability	53
Plant characteristics	56
NDVI.....	56
Chlorophyll content	58
Conclusions.....	60
Acknowledgements.....	61
APPENDICES	62
APPENDIX A- CHAPTER 2 TABLES	63
APPENDIX B- CHAPTER 2 DATA COLLECTED AND NOT USED FOR PUBLICATION.....	74
LITERATURE CITED	77

CHAPTER 3

NITROGEN CYCLING, CORN GROWTH AND GRAIN YIELD RESPONSE TO NON- LEGUMINOUS COVER CROPS AND FERTILIZER.....	83
Abstract.....	83
Introduction.....	84
Materials and Methods.....	87
Site characteristics and cultural practices	87
Experimental design and treatment application	88
Data collection and statistical analysis	89
Soil inorganic nitrogen.....	89
CC measurements	90
Corn measurements.....	90
Results and Discussion	91
Weather.....	91
Autumn CC biomass and nitrogen uptake, and soil NO ₃ -N.....	92
Spring CC biomass and nitrogen uptake, and monthly soil NO ₃ -N.....	93
Corn response to CCs and N strategies.....	95
Grain yield	95
Profitability	97
Normalized difference vegetation index.....	98
Chlorophyll content (SPAD).....	100
Conclusions.....	101
Acknowledgements.....	102
APPENDIX.....	103
LITERATURE CITED	114

CHAPTER 4

INFLUENCE OF COVER CROPS AND NITROGEN MANAGEMENT ON THE SOIL MICROBIOME TO OPTIMIZE CORN PRODUCTION.....	120
Abstract.....	120
Introduction.....	121
Materials and Methods.....	125
Site characteristics and cultural practices	125

Experimental design and treatment application	126
Data collection and processing	127
Cover crops	127
Soil chemical properties.....	128
Soil basal CO ₂ respiration and labile amino-N (SLAN)	128
Soil bacterial phylogenetic diversity.....	129
Data analysis	130
Results and Discussion	131
Environmental conditions	131
Cover crop biomass production and quality	132
Cover crop effects on base cation levels.....	133
Cover crop effects on corn grain yield.....	134
Cover crop effects on SLAN content and soil basal CO ₂ respiration	135
SLAN	135
Respiration	136
DNA sequence analysis	138
Alpha diversity indicators (Inverse Simpson [ISD], Shannon Evenness [Shannon] and Chao1 Richness Estimator [Chao]) before and after corn establishment	138
Before corn establishment.....	139
After corn establishment	140
Conclusions.....	144
Acknowledgements.....	146
APPENDIX.....	147
LITERATURE CITED	158

LIST OF TABLES

<i>Table 2.01.</i> Corn planting date, V4, and V11 sidedress N application dates for SCRf (Lansing, MI) and SVRF (Richville, MI), 2014-2016	63
<i>Table 2.02.</i> Prices received and variable input costs utilized used for profitability analysis, 2014-2016.....	64
<i>Table 2.03.</i> Growing season (April – September) and 30-yr mean temperature (°C) and precipitation (mm) data for SCRf (Lansing, MI) and SVRF (Richville, MI), 2014 – 2016.....	65
<i>Table 2.04.</i> Impact of N strategy on V3 corn plant density prior to in-season sidedress applications using multiple degree of freedom contrasts†, across sites, 2014 - 2016	66
<i>Table 2.05.</i> Analysis of variance for V3 stand count, V6 and V11 NDVI, R1 and R4 rel. chlorophyll meter (SPAD), and grain yield in Lansing, and Richville, MI, 2014 - 2016	67
<i>Table 2.06.</i> Corn grain yield† as affected by in-furrow (IF), pre-plant incorporated (PPI), and 5x5 N strategies in combination with sidedress timings, at SCRf (Lansing, MI) and SVRF (Richville, MI)‡ 2014 - 2016	68
<i>Table 2.07.</i> Corn profitability as affected N strategy at SCRf (Lansing, MI) and SVRF (Richville, MI) 2014 - 2016.....	69
<i>Table 2.08.</i> Mean V6 NDVI readings as affected by N strategy across locations at SCRf (Lansing, MI) and SVRF (Richville, MI) 2014 – 2016	70
<i>Table 2.09.</i> Mean V11 NDVI readings as affected by N strategy across locations and years in Lansing and Richville, MI, 2015 - 2016.....	71
<i>Table 2.10.</i> Mean R1 relative chlorophyll (CM) meter measurements as affected by N strategy at SCRf (Lansing, MI) and SVRF (Richville, MI), MI, 2014 – 2016	72
<i>Table 2.11.</i> Mean R4 relative chlorophyll (CM) meter measurements as affected by N strategy across locations and years, 2015-2016.....	73
<i>Table B.01.</i> Total vegetative N uptake at physiological maturity as affected by in-furrow (IF), pre-plant incorporated (PPI), and 5x5 N strategies in combination with sidedress timings, at SCRf (Lansing, MI) and SVRF (Richville, MI) 2014 - 2016.....	74
<i>Table B.02.</i> Soil residual nitrate at corn harvest as affected by in-furrow (IF), pre-plant incorporated (PPI), and 5x5 N strategies in combination with sidedress timings, at SCRf (Lansing, MI) and SVRF (Richville, MI) 2014 – 2016.....	75

<i>Table B.03.</i> Corn stalk nitrate at corn physiological maturity as affected by in-furrow (IF), pre-plant incorporated (PPI), and 5x5 N strategies in combination with sidedress timings, at SCRF (Lansing, MI) and SVRF (Richville, MI) 2014 – 2016	76
<i>Table 3.01.</i> Dates of field operations and observations in 2014 – 2015 and 2015 – 2016, Lansing, MI.....	104
<i>Table 3.02.</i> Mean monthly and 30-yr precipitation and temperature data for Lansing, MI, 2014 – 2016.....	105
<i>Table 3.03.</i> Main effects of cover crop on dry matter biomass production, total N uptake, and soil residual NO ₃ -N concentrations following autumn growth, Lansing, MI, 2014 – 2015.....	106
<i>Table 3.04.</i> Impact of preceding cover crops not receiving N fertilizer on ensuing monthly soil test nitrate (NO ₃ -N) (0 – 30 cm) concentrations and linear correlation to grain yield (r-value), Lansing, MI, 2015 – 2016.....	107
<i>Table 3.05.</i> ANOVA of cover crop, N strategy, and year fixed effects on corn V6 NDVI, R1 SPAD, R2 height, grain yield, and net return in Lansing, MI, 2015 – 2016	108
<i>Table 3.06.</i> Cover crop and N strategy interaction on corn grain yield (155 g kg ⁻¹ moisture) and impact of cover crop on response to N, Lansing, MI, 2015 – 2016.....	109
<i>Table 3.07.</i> Interaction of cover crop and N strategy on corn profitability, Lansing, MI, 2015 – 2016.....	110
<i>Table 3.08.</i> Main effect of N strategy on corn V6 NDVI in Lansing, MI, combined across 2015 – 2016.....	111
<i>Table 3.09.</i> Corn V6 NDVI measurements as affected by previous cover crop, Lansing, MI, 2015 – 2016.....	112
<i>Table 3.10.</i> Interaction of cover crop and N strategy on SPAD R1 chlorophyll meter readings, Lansing, MI [†] , 2015 – 2016	113
<i>Table 4.01.</i> Monthly precipitation and temperature departures from normal and 30-yr means for Lansing, MI, 2014 – 2016.....	148
<i>Table 4.02.</i> Two-year soil moisture (30-cm), precipitation (cm), and soil temperature (5-cm) data for the 14 d period prior to each soil sampling event (cover crop planting [CC plant], autumn cover crop harvest [CC harvest], corn planting [corn plant], corn tasseling [corn R1], and corn harvest [corn R6]) in Lansing, MI, 2014 – 2016	149
<i>Table 4.03.</i> Cover crop biomass and quality at autumn cover crop harvest, Lansing, MI, 2014 - 2015.....	150

<i>Table 4.04.</i> Main effects of cover crop and N strategy [†] on corn grain yield, Lansing, MI, 2015 – 2016.....	151
<i>Table 4.05.</i> Correlation coefficients (<i>r</i>) (<i>n</i> =10) relating indicators of soil microbial alpha diversity [†] indicators (inverse Simpson diversity [ISD], Chao1 richness [Chao] and Shannon evenness [Shannon]) or soil parameters [‡] (soil labile amino acid [SLAN], CO ₂ respiration, soil NO ₃ -N and NH ₄ -N) to weather factors observed during the 14 d period prior to cover crop planting, autumn cover crop harvest, corn planting, corn tasseling, and corn harvest in Lansing, MI, in 2015 - 2016	152
<i>Table 4.06.</i> Effect of cover crop and no N fertilizer on soil labile amino N (SLAN) and soil respiration (CO ₂) means observed in the autumn and the ensuing May at corn planting, Lansing, MI, 2014 - 2016	153
<i>Table 4.07.</i> Main effects of cover crop and N strategy on basal soil respiration combined across observations at flowering (R1) and physiological maturity (R6), Lansing, MI, 2015.....	154
<i>Table 4.08.</i> Cover crop and N management interaction (<i>P</i> =0.0374) on soil basal CO ₂ respiration combined across R1 and R6 sampling times, Lansing, MI 2016.....	155
<i>Table 4.09.</i> Effect of cover crop on alpha diversity indicators [†] (inverse Simpson [ISD], Chao1 Richness [Chao] and Shannon evenness [Shannon]) sampled in the autumn and spring prior to corn planting, Lansing, MI, 2014 – 2016.....	156
<i>Table 4.10.</i> Main effects of cover crop and N strategy [†] on alpha diversity indicators (inverse Simpson diversity [ISD], Chao1 richness [Chao] and Shannon evenness [Shannon]) across soil sampling zones (rhizosphere and bulk soil) and sample timings (corn tasseling and harvest), Lansing, MI, 2015- 2016	157

CHAPTER 1

LITERATURE REVIEW

Michigan Corn Production

In 2016 corn (*Zea mays* L.) production accounted for 27% of total United States crop acres and 32% of total Michigan (MI) crop acres (USDA-NASS, 2017). Corn grain production is Michigan's second largest cash receipt commodity (>1,000 million US\$) and was ranked 13th (2016) in average United States production (Mg^{-1}) (https://www.nass.usda.gov/Quick_Stats/Ag_Overview/stateOverview.php?state=MICHIGAN; verified 15 Oct. 2017). Crop nitrogen (N) fertilization represents 59% of major U.S. applied plant nutrients with corn accounting for >45% of total N fertilizer applied at a mean rate of 157 and 137 kg N ha⁻¹ in the U.S. and MI, respectively (USDA-ERS, 2016).

Tissue Nitrogen

Nitrogen is a constituent of chlorophyll molecules, proteins, enzymes, nucleic acids, a component of energy transfer compounds such as adenosine diphosphate (ADP) and adenosine triphosphate (ATP), and influences carbohydrate utilization during photosynthesis (Havlin et al., 2014). Nitrogen is often the most limiting nutrient in grain cropping and natural systems due to its large plant elemental composition (1 – 6%) (Fernández et al., 2009; Robertson and Vitousek, 2009). Like many crops fertilizer N is applied to corn to supplement native soil N supplies. Corn N requirements are a function of physiological requirements for biomass yield and tissue N content (Cassman et al., 2002). Corn grain has been found to remove 11.6, 4.8, and 3.5 kg m⁻³ of N, P₂O₅, and K₂O, respectively, and a corn yield of 10,000 kg ha⁻¹ may remove 190 kg N ha⁻¹

(Cassman et al., 2002; Warncke et al., 2009). Corn yield potential is affected by interacting factors including agronomic practices [e.g., fertility management], cultivars, and the environment (Evans and Fischer, 1999). Corn production will require accurate estimates of soil N supply and corn requirements across environments when determining fertilizer N application strategies.

Climate Variability

Recent studies project more heat waves (5°C above climatic normal) and increasing air temperatures (1.5 to 2.0°C) over the next 30 years may reduce early spring frost dates (1 wk) in the northern hemisphere in addition to increased precipitation frequency in winter and spring months with more intense events [> 50.8 mm in 48 hours] (Schwartz et al., 2006; Tebaldi et al., 2006; Hayhoe et al., 2007; Karl et al., 2009). Warming spring temperatures are likely to move corn planting dates earlier than currently optimal [1 and 7 May] (Lauer et al., 1999). In lieu of mounting concerns over Great Lakes Basin surface- and ground-water quality, weather variability may increase N loss opportunities and require further refining of N management strategies in response to changing climates (Scharf et al., 2002; Scavia et al., 2014; Dove and Chapra, 2015).

Nitrogen Use Efficiency

Nitrogen use efficiency (NUE) has been defined as the ratio of output (biological or economic yield) to N supply, where biological yield consists of total plant dry matter or total plant N, economic yield as grain yield or total grain N (Cassman et al., 2002; Ladha et al., 2005).

Nationally the price of urea-N and N-solutions, two commonly used corn N sources, have risen from \$159 and \$104 per ton in 2000 to \$472 and \$327 in 2013, respectively

(<http://usda.mannlib.cornell.edu/MannUsda/viewDocumentInfo.do?documentID=1002>; verified 14 Nov. 2017). Corn prices received by growers have sharply declined from \$6.89 in 2012 to \$3.65 in 2014 (https://www.nass.usda.gov/Quick_Stats/Lite/index.php; verified 14 Nov. 2017). Raun and Johnson (1999) reported worldwide NUE for cereal production was only 33%, while the remaining 67% represented a \$15.9 billion annual loss of N fertilizer. A regional survey of combined on-farm corn research experiments conducted across Illinois, Michigan, Minnesota, Missouri, Nebraska, and Wisconsin found that 37% of applied N fertilizer was recovered in aboveground plant biomass (Cassman et al., 2002). Previous studies indicate N management strategies that account for N loss opportunities may improve NUE (Raun and Johnson, 1999). Despite a linear trend for increasing NUE from 1980 to 2004, Bundy (2006) suggested that year to year variation of NUE illustrated a need for improved assessment of NUE factors such as N rate selection, accounting for non-fertilizer N sources, and managing applied N. Increased costs of fertilizer N, decreased grain prices, and continued degradation of Great Lakes water quality, suggests further improvement of N management strategies that simultaneously increase NUE and reduce opportunities for N loss (Dove and Chapra, 2015; USEPA, 2017).

Corn Response to N Fertilizer

Soil texture and weather can interact to change N availability in the corn rooting zone and affect corn response to applied N (Tremblay et al., 2012). A 25 mm rainfall has moved nitrate downward 25 mm in clay loam soils and 63.5 mm in sandy soils (Nelson and Huber, 2001). Heavy rainfall events (i.e., > 25.4 mm) in June may reduce corn N availability through denitrification or leaching beyond the active rooting zone (Kyveryga et al., 2010). Rainfall frequency may also impact corn response to N. Infrequent rainfall events may result in dry soil

conditions and reduce plant fertilizer N uptake and increase residual inorganic N levels (Gagnon et al., 2012; Jokela and Randall, 1997).

4R Nutrient Management

To provide a framework of site-specific nutrient management decisions, the 4R nutrient stewardship concept has been adopted by the world's fertilizer industry. The 4R concept emphasizes the right 'source' of nutrient application, applied at the right 'rate', at the right 'time', and in the right 'place' to guide best management practices (BMP) that ensure fertilizer uptake is optimized, and environmental loss is minimized (Roberts, 2007). While the 'right' source, placement, and time can help achieve synchrony between N supply and N uptake, correct N rate emphasizes N excess and deficiency. Optimizing the '4R's may help synchronize N availability with crop N demands (Randall and Mulla, 2001). The concept of BMPs is fundamental to efficient and judicious use of nutrients. General fertilizer recommendations can fail to account for external factors such as crop rotation, rainfall, pest pressure, and temperature (Mikkelsen, 2011). In conventional tillage systems spring pre-plant N applications to medium- and fine-textured soils is considered a BMP although spring pre-plant applications of anhydrous ammonia to no-till and strip-till systems may be undesirable to some growers due to delayed planting and compaction of wet soils (Vetsch and Randall, 2004). Growers should consider the ability of the soil to supply N, crop N requirements, climate, and management capabilities when determining 4R requirements (Roberts, 2007).

Corn N application rates

Corn N application rates are based on N demand minus indigenous N soil contributions (Cassman et al., 2002). Biological transformation can result in several mobile forms of N susceptible to leaching, volatilization, surface runoff, gaseous plant emission, denitrification, or erosion making N a dynamic and leaky system (Randall et al., 2008; Millar et al., 2010; Congreves and Eerd, 2015). The optimum N application rate has been regarded as the point where the last increment of N fertilizer results in a profit just large enough to pay for the N (EONR). Unlike phosphorus and potassium plant available N lacks the mechanism for long-term mineral soil storage of weatherable N pools (Robertson and Vitousek, 2009). Applications of N at the EONR can minimize N losses (Hong et al., 2007; Jokela and Randall, 1997; Stevens et al., 2005a). Predicting the optimum rate of N for corn production has both economic and environment implications.

Growers have previously relied on ‘insurance N’ where greater applications were made to ensure sufficient amounts are available for plant N uptake (Schröder et al., 2000). However, rates in excess of crop N uptake may influence the potential for NO_3^- leaching or denitrification, especially during wet and warm conditions (Dinnes et al., 2002). Karlen et al. (1998) found the potential for N loss was greatest when residual soil NO_3^- was present without active crop growth. Andraski et al. (2000) observed increased NO_3^- in groundwater where N rates were in excess of a corn crop EONR (150 kg ha^{-1}). Studies by Steven et al. (2005a; 2005b) determined where excessive corn N ($> 201 \text{ kg N ha}^{-1}$) applications were made: 1) mineralization of residual soil N is increased and 2) response of subsequent crops to fertilizer N may decrease due to increased mineralization of newly formed organic N pools and crop residues. Increased mineralization due

to excess N application ($>201 \text{ kg N ha}^{-1}$) increased the risk of N loss to the environment and emphasized the importance of fertilizing at optimal N levels (Stevens et al., 2005a).

Maximum return to nitrogen: Michigan N recommendations are based on crop N utilization and response to applied N. In 2006 Michigan adopted the Maximum Return to Nitrogen (MRTN) recommendation system. The MRTN is a pre-season general recommendation model (as compared to a field-specific or site-based model) that does not account for individual site variability nor variable in-season weather trends which can affect corn N response. The MRTN system was developed by Iowa State University as a regional guideline approach to N recommendations and has been adopted by Illinois, Michigan, Minnesota, Ohio, and Wisconsin. The MRTN approach assumes a linear increase in N cost and a grain yield plateau. The MRTN system is updated annually and N rate incorporate variability among years. The MRTN approach was preceded by yield-based N recommendations based on Stanford (1973), which utilized a mass-balance approach accounting for dry matter yield goal and non-fertilizer N contributions. Yield goal was multiplied by a 15.4 kg m^{-3} (1.2 lb N per bushel) factor, and adjusted for non-fertilizer N contributions (Sawyer et al., 2006). The rate constant was supported by research demonstrating that optimum yield (yield at the economically optimum N rate) divided by the optimum N rate was about 1.5 kg N hL^{-1} (Fernández et al., 2009). Problems associated with this system included unrealistic yield goals, lack of a general use consensus, and poor relationships of actual optimum N rates and N recommendations (Vanotti and Bundy, 1994). In Michigan, the yield goal approach was used to develop a linear N recommendation equation based on yield potential (Warncke et al., 2009). Using the yield goal approach, over-application of N could result from the assumption of limitless yield with increasing N application.

Illinois soil N test versus the Michigan sidedress soil nitrate test: Assessing the soil's ability to supply N for plant uptake may help synchronize available N with uptake and reduce the potential for NO_3^- leaching (Andraski et al., 2000; Dinnes et al., 2002). Soils where manure has been previously applied may not respond to additional N applications (Scharf et al., 2002). Soils unresponsive to N illustrate the importance of determining first year N availability from animal manure and credit this portion of available N towards N application rates (Oberle and Keeney, 1990). Mulvaney et al. (2006) utilized the Illinois soil N test (ISNT) to quantify corn response to N fertilization in nonresponsive soils in various rotations at a critical test level of 230 mg kg^{-1} amino sugar N. The ISNT measured amino sugar N compounds which may mineralize during the growing season. Fernández et al. (2009) stated when cool soils limited N mineralization into early June, high values of the ISNT did not correspond to reduced fertilizer N needs. In Michigan the sidedress soil nitrate (PSNT) test is used to measure residual NO_3^- from previous season inputs and early-season mineralized organic-N (Warncke et al., 2009). The PSNT involves sampling approximately 6 weeks after planting and indicates the additional amount of fertilizer N to supply and appropriate N credits when test levels are ≥ 10 and $\leq 25 \text{ mg NO}_3^- \text{ kg}^{-1}$ (Dinnes et al., 2002; Fernández et al., 2009). Sidedress N rates with PSNT adjustment have resulted in $< 8.0 \text{ mg L}^{-1}$ flow-weighted NO_3^- concentrations in percolate and 8-10% less total N applied when compared to unadjusted N rates on a Woodbridge fine sandy loam soil (Guillard et al., 1999).

Corn N application timing

Michigan State University does not recommend fall applications of N to corn in Michigan. May rainfall and soil temperature $>10^{\circ}\text{C}$ promoting nitrification can reduce corn grain yield and N uptake when N demand is low (Randall and Vetsch, 2005). A survey in 2005 indicated $<5\%$ of total annual corn N was fall applied in MI over 2.2 million acres (Randall and Sawyer, 2008). Sanchez and Blackmer (1988) found 49-64% of fall applied N to be lost from the upper 1.5 m of surface soils, and wet and warm soil conditions during the spring may exacerbate N losses (Vetsch and Randall, 2004). Nitrogen recovery and grain yield were reduced 42 and 20% (year 1), respectively, with fall versus spring applied N in addition to decreased corn grain N concentration from a poorly drained clay-loam soil and did not include a nitrification inhibitor (Vetsch and Randall, 2004). In northern states (i.e., Minnesota) soils can be poorly drained, and spring pre-plant N applications are often utilized in conventionally tilled soils (Vetsch and Randall, 2004).

Failure to supply N to corn at critical stages of growth may lead to yield loss (Binder et al., 2000; Subedi and Ma, 2005; Walsh et al., 2012). Weather, planting date, and N application timing may affect N uptake (Scharf et al., 2002). Traditional sidedress N timing in Michigan is V4 to V6 corn which leaves time for soil incorporation by rainfall and does not require the use of high-clearance equipment. Early studies by Karlen et al. (1988) suggested two patterns of N accumulation: 1) V12-V18, and 2) during grain fill. Recently Bender et al. (2013) observed N uptake by corn is greatest during V10-V14 and at reproductive (R) stages R2-R6 with limited uptake at tasseling (VT)/R1. Utilizing modern corn hybrids Bender et al. (2013) observed that corn had taken up 65% of total N by R1 and total grain N comprised 58% of total N uptake when

N was not limiting. Improved fertility practices should focus on matching available N during these periods of high vegetative uptake.

Applying corn fertilizer N after planting may be an effective strategy to increase NUE and reduce overall N rates (Gehl et al., 2005; Walsh et al., 2012). Utilizing split-N applications, Gehl et al., (2005) concluded yield-goal based N rates could be reduced 40% as compared to a single pre-plant application in Kansas. Split N applications (45 kg pre-plant/45kg V6 or V10) maximized NUE and doubling these N rates maximized grain yield (Walsh et al., 2012). Delayed N applications may avoid exposure to early-season weather events and reduce opportunities for N loss, but growers can risk irreversible yield loss due to N stress (Binder et al., 2000; Scharf et al., 2002). Jokela and Randall (1997) further noted N uptake was reduced from 54 to 48 kg ha⁻¹ when N application was delayed until V8 due to extremely dry soil conditions limiting N availability. Maximum grain production will require earlier N applications when corn is N stressed (Binder et al., 2000). A single N application at V6 was found to reduce yield up to 12% when the chlorophyll meter N sufficiency index was below 0.90 (Binder et al., 2000), but a grower missing the opportunity to apply N early could apply N up to silking with a 15% yield reduction (Scharf et al., 2002). Walsh et al. (2012) determined sidedress N applied at V6 allowed for crop recovery when no pre-plant N was supplied but sidedress N beyond V6 did not. When pre-plant N was provided (45 or 90 kg ha⁻¹) no yield differences were observed among sidedress timings of V6, V10, or tasseling (VT) (Walsh et al., 2012). In-season weather variables should guide sidedress decisions where increased June rainfall can reduce corn N availability (Gagnon and Ziadi, 2010; Kyveryga et al., 2010).

Corn N fertilizer source

Corn response to N fertilizer source can be affected by corn management and weather and may affect urea-N fertilizer sources. Up to 35% of applied N has been lost due to non-incorporated applications of urea and UAN if no rain fell within 6 days of application in no-till corn (Fox and Hoffman, 1981). Volatilization of surface applied urea to wheat were found to exceed 50% when urea was surface applied to Saskatchewan soil types (Aridic, Typic, Vertic, and Udic Haploborolls) (Fowler and Brydon, 1989). On a Kamouraska clay soil (Orthic Humic Gleysol) a split application of calcium ammonium nitrate (CAN) + UAN sidedressed at V6 produced comparable corn yields relative to polymer coated urea (PCU) applied in a single pre-plant incorporated (PPI) application (Gagnon et al., 2012). In the same study, PCU, UAN at V6 sidedress (injected), and urea with a nitrification inhibitor N sources resulted in yields $\geq 0.6 \text{ Mg ha}^{-1}$ as compared to urea PPI in wet soils when average June-July soil temperatures were cool (17.6°C). In cool and humid climates UAN has increased mean apparent N recovery 44.4% as compared to CAN and aqua ammonia (AA) (Gagnon and Ziadi, 2010). Increased apparent N recovery of UAN corresponded to a lower optimum N rate of 100 kg ha^{-1} required to maximize grain yield relative to 124 and 128 kg N ha^{-1} observed for CAN and AA, respectively (Gagnon and Zaidi, 2010).

Application of enhanced efficiency fertilizer (EEF) N sources can be used to extend N release for periods of corn N uptake and achieve comparable recovery efficiency (48-56%) to UAN sidedress V6 (Gagnon et al., 2012). Observing leaf chlorophyll values during grain fill, Hatfield and Parkin (2014) demonstrated the effectiveness of EEF sources (i.e., PCU, stabilized urea with urease and nitrification inhibitors, and UAN with or without a combination urease and

nitrification inhibitor). In their study delayed leaf senescence resulted in a greener canopy and corresponded to improved grain yield despite variable weather among study years. Economic analysis revealed PCU and UAN increased net return \$95 to \$263 ha⁻¹ in wet years relative to urea but in a warm and dry season UAN returned \$154 ha⁻¹ as compared to PCU \$-46 ha⁻¹ (Gagnon et al., 2012). Michigan studies on Capac loam soils attained mean corn yields of 9.7 Mg ha⁻¹ (NUE = 0.61 Mg grain kg N⁻¹) when PCU was blended with urea (75:25) and broadcast + incorporated 2 to 4 weeks before planting as compared to yields of 8.3 Mg ha⁻¹ (NUE = 0.46 Mg grain kg N⁻¹) achieved with a similar application of 100% urea (Steinke and Chomas, 2014). When 102 mm less rainfall was avoided with at-plant N applications, a 25:75 blend PCU + urea was sufficient to increase corn yield 2.7 Mg ha⁻¹ relative to pre-plant urea-N applications demonstrating the potential of PCU fertilizer to reduce N loss for surface N applications (Steinke and Chomas, 2014).

Corn starter N fertilizer placement

Collectively, fertilizers placed with or near the seed at planting are generally referred to as starter fertilizer. Starter fertilizer may utilize either single element or complete NPK mixtures even in soils testing high in P and K (Niehues et al., 2004; Rehm et al., 2009). Starter fertilizer increased corn yield in Wisconsin 0.27 Mg ha⁻¹ and decreased grain moisture at harvest 0.5% regardless of soil test phosphorus (STP) (Bundy and Andraski, 1999). Starter fertilizer has increased yield in Iowa 1.1-2.4% and early growth (V5-6 whole plant dry weight and N uptake) 20-32% but studies noted early season corn growth and nutrient uptake associated with starter N do not always correspond to yield improvements ($R=0.44$) or STP (Bermudez and Mallarino 2002; 2004; Kaiser et al., 2005). Plant biomass (V6-V8) has been increased more in irrigated corn (30%) at 5

sites as compared to rainfed corn (10%) at 3 of 5 sites with starter fertilizer application although grain yield was only increased 0.86 Mg ha⁻¹ at irrigated sites (Wortman et al., 2006).

In-furrow: Michigan growers sometimes place fertilizer with corn seed at planting to improve early corn growth. In-furrow nutrient placements (sometimes referred to as "pop-up" or "seed-placed" starter) can provide immediate nutrient access to young corn roots (Niehues et al., 2004). In-furrow placements have been more sensitive to fertilizer rate than sub-surface band placements (Niehues et al., 2004). Salt rates (N+K₂O) in excess of 11.2 kg ha⁻¹ could delay seedling emergence until precipitation occurs and result in uneven corn stands (Raun et al., 1986). Reported salt indices for N sources utilized by corn growers are 20.0, 63.0, and 74.4 for ammonium polyphosphate (10-34-0), UAN 28%, and urea, respectively (Kaiser and Rubin, 2013). Niehues et al. (2004) found 56 kg N ha⁻¹ placed in-furrow reduced corn populations 12,000 and 21,000 plants ha⁻¹ in two notably dry years on a silt loam textured soil. When April + May rainfall increased 100%, little effect on corn stand was observed at the 45 and 56 kg N + K₂O rates illustrating the potential dilution of reduced N rates at planting (Niehues et al., 2004). Niehues et al. (2004) concluded the N component was responsible for reduced seedling emergence in contrast to Rehm and Lamb (2009) who recommended the sum of N+K₂O should guide considerations due to salt damage. Common in-furrow corn fertilizers include 10-34-0, 7-21-7, and 9-18-9. At equal P₂O₅ rates (0, 14, 28, and 56 kg ha⁻¹), 10-34-0 and 7-21-7 have resulted in less corn stand reduction (>93%) as compared to 9-18-9 (41 to 66%) (Gerwing et al., 1996). To prevent seedling injury Michigan recommendations for in-furrow applications to corn are ≤5.6 kg ha⁻¹ N + K₂O with seed where CEC is < 7 cmol(+) kg⁻¹, and ≤ 9 kg ha⁻¹ N+K₂O where CEC is ≥ 8.

Subsurface band: Subsurface placement can vary relative to seed placement with bands 5cm below and 5cm laterally (“2x2” or “sub-surface [SS]band”) as compared to within the seed furrow "pop-up" or "in-furrow" (Niehues et al., 2004). In-furrow and SSband placements have been utilized at corn planting in Michigan to encourage early season plant growth and nutrient uptake where cool and moist soil conditions associated with early planting dates can affect corn ontogeny (Bollero et al., 1996). Rates of SSband N up to 134 kg N ha⁻¹ did not affect plant population while 34 kg N ha⁻¹ optimized yield (5.4 to 8.3 Mg ha⁻¹) on silt loam textured soils in Kansas. Increased yield consistency has been observed with the use of SSband placement as compared to in-furrow placements (Ritchie et al., 1996).

Corn Response to Starter N

Predicting corn response to starter fertilizer application has been inconsistent (Randall and Hoef, 1988). A review of placement methods by Randall and Hoef (1988) concluded that efficacy of starter fertilizer placement was dependent on crop, soil texture, tillage system, and climatic conditions. Magnitude of corn yield response to starter fertilizer has been more likely when STP is <16 mg kg⁻¹ suggesting response to a P component (Bermudez and Mallarino, 2002; 2004; Kaiser et al., 2005; Wortman et al., 2006). Grid-point soil sampling confirmed spatially-variable yield response to starter NPK to portions of fields with low STP (15 mg kg⁻¹) deemed otherwise unresponsive ($P=0.16$) (Bermudez and Mallarino, 2004). Starter fertilizer has also increased corn yield where soil STP ≥ 24 and attributed to an N or P component of the product (Bermudez and Mallarino, 2002; 2004; Vetsch and Randall, 2002). For example corn yield response to starter fertilizer was increased (0.20 to 0.67 Mg ha⁻¹) in fields where STP was low (<16 mg P kg⁻¹,

Bray-P₁) (Bermudez and Mallarino, 2002). In soils with high STP (21 to 30 mg kg⁻¹) corn yield response to in-furrow placement (4.5 to 9.1 kg N ha⁻¹) was less (0.08 to 0.19 Mg ha⁻¹) compared to a SSband placement (16.3 to 27.2 kg N ha⁻¹) (0.17 to 0.47 Mg ha⁻¹) (Bermudez and Mallarino, 2002). Bundy and Andraski (1999) determined an arithmetic combination of relative hybrid corn maturity date (increasing maturity date) + plant date (increasing plant date) (PDRM) increased the profitable yield response 34% where soil K levels were below excessively high (EH; <140ppm) and 17% where soils were above EH. Bundy and Andraski (1999) attributed increased yield from SSband placement relative to a no SSband control to increased yield potential realization due to early season growth stimulation. Positive economical response to the SSband placement was most likely when PDRM > 235 and soil K levels are < 140 mg kg⁻¹ (Bundy and Andraski, 1999).

Tillage has affected corn response to starter fertilizer. Spring (beginning Julian day 100) soil temperatures and gravimetric water content have been increased 3 to 5° and reduced 20 to 40 g kg⁻¹, respectively, in conventional till (CT) (fall chisel 20 cm + spring disk [2x] 10 cm) as compared to reduced (fall) or no-till (NT) corn on a silt loam soil in Wisconsin (Wolkowski, 2000). The authors reported yield increased 0.5 Mg ha⁻¹ with SSband starter as compared to a no starter control across tillage systems despite excessively high STP (48 mg kg⁻¹) and STK (167 mg kg⁻¹) but observed no yield increase when SSband was used in a CT system (Wolkowski, 2000). On a silt loam soil in Minnesota corn emergence and yield were reduced 2.7 d and 7%, respectively, in NT continuous corn (CC) as compared to CT (Vetsch and Randall, 2000). An SSband starter N-P-K increased corn height >5 cm observed 35 d after emergence and yield 0.5 Mg ha⁻¹ as compared to a no starter control in a corn-soybean (CS) and CC rotation despite high

STP and high STK (≥ 24) and ≥ 130 mg kg⁻¹, respectively) (Vetsch and Randall, 2000). Deep placement of starter fertilizer (15 cm) was not an effective placement as compared to SSband for increasing yield in the north (Minnesota and Wisconsin) (Vetsch and Randall, 2000; Wolkowski, 2000). In Iowa corn response to liquid starter placement (SSband or in-furrow N-P-K) was similar across CS tillage systems (NT or spring tillage) in reference to a no starter control when P and K were broadcast 1 to 4 mo. before tillage (Bermudez and Mallarino, 2004). The authors further noted soil test levels (P, K, pH, and OM) and soil series were not useful predictors of a yield increase to starter fertilizer (Bermudez and Mallarino, 2004).

Soil texture (clay loam and silty clay loam as compared to loamy fine sand) has affected corn yield response to N rate (13.1 as compared to 6.6 kg ha⁻¹) using 10-15-0 (placed either in-furrow, 1.25, or 1.90 cm away from seed) (Rehm and Lamb, 2009). Grain yield reductions of 6.9 and 14.3% corresponded to reduced emergence of 8.6 and 31.9%, respectively, when 10-15-0 was applied at 13.1 kg N ha⁻¹ on a loamy fine sand textured soil (Rehm and Lamb, 2009). Grain yield was not increased relative to a control (without starter) regardless of starter grade, rate, or placement in contrast to increased V4-5 whole plant weight and P uptake on clay loam and silty clay loam textured soils (Rehm and Lamb, 2009). Stand loss was noted on sandy soils with in-furrow placement of 93.5 L ha⁻¹ 10-15-0 but equivalent rates of 4-4-8 N-P-K and 63.6 L ha⁻¹ 3-8-15 N-P-K did not (Rehm and Lamb, 2009). In fine-textured soils placement of all grades did not impact emergence demonstrating the importance of soil texture when considering in-furrow sources and rates (Rehm and Lamb, 2009).

Corn Nutrition Status

Chlorophyll meter

Determining corn N status may be a useful tool to correct in-season N deficiencies and improve N management (Hawkins et al., 2007; Varvel et al., 1997). The Konica Minolta SPAD 502 chlorophyll meter (CM) has been used to determine chlorophyll content of plants relative to corn receiving non-limiting rates of N (rel. chlorophyll). Light passing through a leaf is captured by another sensor below the leaf and the filtered wavelengths is a relative indication of chlorophyll content per leaf unit area (Richardson et al., 2002). Studies have determined chlorophyll contents obtained in corn leaves 1 week before silking ($a + b = \text{total}$) were correlated to tissue N content ($r^2 = 0.97$) and a predictor of N sufficiency (Asghari and Hanson, 1984; Lohry, 1989). Leaf tissue N has been correlated to CM readings while differences in greenness associated with hybrids emphasize the importance of reference strips (Rorie et al., 2011; Schepers et al., 1992; Varvel et al., 1997). Readings have been used to detect N deficiencies in irrigated corn to initiate additional N applications as well as N sufficiency where normalized readings were determined in regards to a high N reference strip to estimate excessive N fertilization (Blackmer and Schepers, 1995; Piekielek et al., 1995). Nitrogen sufficiency indices (i.e., sufficiency index) can be used to monitor crop N status and are determined using readings taken for a crop divided by the reference or reading taken from the crop where N was not limiting. Varvel et al. (1997) used the CM to separate N deficient and sufficient corn beginning at V8 using a sufficiency index of 95% and used this to make 30 kg N ha^{-1} applications beginning at V8 corn to obtain maximum yields. Corn N supply was deemed insufficient when V8 corn fell below a sufficiency index of 90% and full yield was not achieved with further N additions (Varvel et al., 1997). At the late milk stage Piekielek et al. (1995) observed a corn N sufficiency index of 0.93. Samborski et al. (2009)

provided a comprehensive review of methods and strategies which included normalization of CM data used to establish relationships between N rates and CM data and indicated the need for accurate reference strips.

NDVI

Normalized difference vegetation index (NDVI) is an indicator of green biomass, indicates plant vigor, and has been used as a tool to assess in-season corn N status and adjust N rates (Tucker, 1979; Teal et al., 2006; Samborski et al., 2009). Deriving N rates utilizing NDVI requires correlation of NDVI and yield to determine yield potential at a particular corn growth stage (Teal et al., 2006). Using the GreenSeeker sensor, Teal et al. (2006) observed significant relationships between yield and NDVI and noted V8 corn resulted in the strongest correlation ($R^2=0.77$). In-season measurements at V9 could not determine treatment differences due to canopy closure (Teal et al., 2006). Normalizing NDVI with growing degree days (GDD_{50}) increased the timeframe for assessment plus N application where NDVI sensed at 800 to 1000 GDD's was correlated with yield ($R^2=0.76$), which encompassed V7-9 corn (Teal et al., 2006).

Basal cornstalk nitrate test

The Basal Cornstalk Nitrate Test (CSNT) is a late-season assessment of deficient/sufficient/excess corn N supply for the growing season taken 1-3 weeks after black layer formation on 80% of kernels (Fox et al., 2001). The CSNT indicates N supply to the corn plant enabling growers to assess optimization of N management practices. In Michigan, 'low' ($<700 \text{ mg NO}_3^- \text{ kg}^{-1}$), 'optimal' (700 to 2000 $\text{mg NO}_3^- \text{ kg}^{-1}$) and 'excessive' ($>2000 \text{ mg NO}_3^- \text{ kg}^{-1}$) CSNT levels are used as criteria (Silva, 2011). Brouder et al. (2000) determined a critical

cornstalk $\text{NO}_3\text{-N}$ level of 1.67 g kg^{-1} separated sufficient from excessive N categories while at 2900 mg kg^{-1} fertilizer efficiency approached zero. Values of CSNT have been positively correlated to residual soil nitrate illustrating the test's ability to detect luxury consumption (Forrestal et al., 2012; Fox et al., 2001). Precipitation can impact results of the CSNT. Less than $2000 \text{ mg NO}_3^- \text{ kg}^{-1}$ (excessive) did not always correspond to low residual soil $\text{NO}_3\text{-N}$ values where drought reduced corn production (Forrestal et al., 2012). Likewise, excessive rainfall in June has been found to reduce N availability which decreased the likelihood of obtaining 'excessive' CSNT values (Kyveryga et al., 2010).

Cover Crops and Soil N Dynamics

High N demands of crop systems coupled with soil N mobility present a niche where the use of living ground covers can be used to alter N cycling. Cover crops (CC) have been found useful to reduce soil erosion, increase OM, reduce soil compaction, and provide suppression of weeds (Kaspar and Singer, 2011). Winter annual CCs have been used to reduce the risk of N loss in the non-cash crop growing season of Michigan corn rotations, but spring kill date can impact synchrony of N mineralization and corn N needs (Crandall et al., 2005; McSwiney et al., 2010). In Michigan crop rotations winter wheat (soft red or soft white) (*Triticum aestivum* L.) sometimes precedes corn, and is often harvested in July leaving approximately a 90 d period of fallow soils prior to the first fall freeze (-2.2°C) that CCs may be established (National Oceanic and Atmospheric Administration [NOAA], <http://www.weather.gov/dtx/firstFallfreeze>; verified 02 Aug. 2017). In one Michigan study, winter wheat and cereal rye CCs preceding a corn cash crop were used to "tighten" the N cycle where early-season N availability preceding the corn cash crop was immobilized until N uptake by the corn in August (McSwiney et al., 2010). In the

same study, no differences in corn yields or total N uptake were observed between CC and no CC plots. On an irrigated loamy sand Andraski et al. (2005) increased corn grain yield (1.4 Mg ha⁻¹) utilizing oat, winter triticale, and winter rye CCs, reduced risk of N loss on sandy soils, and reduced mean EONR 32 kg N ha⁻¹ as opposed to winter fallow.

Cover crop kill date and incorporation method preceding a corn cash crop may affect in-season corn N availability (McSwiney et al., 2010). Delaying CC kill date can decrease the amount of available N for early corn N uptake (Crandall et al., 2005). A cereal rye CC killed 7 d before corn planting in May in Illinois resulted in reduced corn biomass, N uptake, and yield (Crandall et al., 2005). Grassy CCs similar to cereal rye may need to be killed no sooner than 14 d preceding a corn cash crop to avoid yield reductions and indicate corn N strategies that include the use of delayed N applications may have to be implemented to account for reduced N availability due to CC (Crandall et al., 2005). Cover crop residue decomposition and N mineralization and immobilization are dependent on the carbon (C) to nitrogen ratio (C:N) and lignin concentrations (Wagger et al., 1998; Jahanzad et al., 2016). Growers may need to consider CC biomass quality and quantity when deliberating between fertilizer N strategies to synchronize corn N demand and N availability.

Radish cover crops

Radish (*Raphanus sativus* L.) is a non-leguminous *Brassicaceae* CC utilized as a N scavenger in the non-growing season to sequester available soil NO₃-N (Mutch and Thelen, 2003; Warncke et al., 2009). Radishes are a large-taproot annual that grow quickly in late summer and autumn and have been observed to reduce soil compaction and suppress winter annual weeds (Williams and

Weil, 2004; Clark, 2007; Lawley et al., 2011). Radish winterkills when air temperatures drop below -4 °C eliminating the need for chemical termination (Thomas et al., 2017). Corn grain yields following an autumn radish cover crop have varied showing both no effect (Vyn et al., 2000; Gieske et al., 2016a; 2016b) and increased yield (Dapaah and Vyn, 1998; Vyn et al., 1999; O'Reilly et al., 2012) compared to no cover crop. Decreased C:N associated with a radish cover (14 to 31:1) relative to a grassy cover (15 to 38:1) (e.g., oats, triticale, or ryegrass) may help explain the 43 and 24% greater soil NO₃-N concentrations with a radish cover crop at mid-May corn planting and mid-June N sidedress application in Ontario, respectively (Vyn et al., 1999; Andraski and Bundy, 2005).

Oat cover crops

Oat (*Avena sativa* L.) is a non-leguminous cover crop also used to sequester residual soil N and prevent wind and water erosion (Warncke et al., 2009). Similar to radish, oats are an annual cover crop and that is subject to winterkill (Johnson et al., 1998; Snapp et al., 2005). Unlike radish, oat has a dense, fibrous root system characteristic of cereal cover crops (Thorup-Kristensen, 2001). Increased lignin and cellulose may reduce decomposition rates and dry matter loss in relation to a radish CC (Jahanzad et al., 2016). The C:N ratio of oat may exceed 30:1 which can immobilize soil N and reduce availability for corn uptake (Andraski and Bundy, 2005; Dean and Weil, 2009). Corn yield following oat cover crops have been generally unaffected or decreased in rainfed studies (Johnson et al., 1998; Vyn et al., 2000; O'Reilly et al., 2012). However, Andraski and Bundy (2005) suggested corn yield gains due to oat cover crops in an irrigated study were an effect of rotation and not N contributions as evidenced by similar May soil NO₃-N contents and reduced EONR (32 ± 8 kg N ha⁻¹). Variable corn yield response to

antecedent CCs suggest corn N fertilizer strategies may have to account for N availability affected by CC decomposition rates.

Plant, Soil, and Microbial Community Interactions

The soil is a living system composed of biological, chemical, and physical fractions (Doran and Zeiss, 2000). Plants, soils, and soil microbes function simultaneously to influence plant health and productivity (Chaparro et al., 2012). Plant-microbe interactions can enhance plant growth through ecosystem services such as nutrient cycling, plant pathogen control (biocontrol agents), and phytohormone production (e.g. auxin) (Nihorimbere et al., 2011). Soil microorganisms mediate transformations of N P, K, S, Fe, Zn, Cu, and Mn to soluble forms while plants can stimulate microbes through rhizodeposition of C-rich root exudates and residues, root morphology, and genotype (Berg and Smalla, 2009; Lambers et al., 2009; Aira et al., 2010; Prashar et al., 2014; Coyne and Mikkelsen, 2015). Greater than 95% of soil N is in organic form and requires microbial conversion for plant uptake (Havlin et al., 2014). Microflora can also impact soil structure. Plant roots and fungal hyphae can “enmesh” larger aggregates, while secretions from roots, fungal hyphae, bacteria, and fauna (i.e. earthworms) are involved in stabilizing smaller aggregates (Oades, 1993). In a barley rhizosphere, Cytophaga-like bacteria produced exopolysaccharides to decompose soil OM and contributed to soil aggregation (Johansen and Binnerup, 2002). A recent study conducted at four locations across Michigan determined the interaction of location, plant species, and soil attributes explained more variation (27%) in structure of bacterial communities than when each factor was considered alone (3.5%, 2.1%, and 0.7%, respectively) ($P \leq 0.15$) and illustrated the complex relationship between bacterial populations and their environment (da C. Jesus et al., 2010).

Increased soil bacterial diversity (e.g. functional, taxonomic) may enhance functional resiliency and increase overall microbial community stability against stress (i.e., environmental variation or disturbance) (Eisenhauer et al., 2012; Awasthi et al., 2014). The removal of crop residues, tillage, and reduced cropping system diversity may affect C substrate quality and reduce soil bacterial community diversity and abundance (Ceja-Navarro et al., 2010; Figuerola et al., 2015; Sun et al., 2016). Nitrogen fertilization strategies may select for distinct microbial communities or reduce community diversity while CCs may provide opportunities to increase cropping system diversity and affect microbial communities that can benefit cash crop health (Berg and Smalla, 2009; Li et al., 2016; Soman et al., 2017).

Effect of Plant Diversity on Soil Microbial Communities

Plant species diversity can influence the structure and function of microbial communities (Berg and Smalla, 2009). In greenhouse studies, introduction of new plant species to soils which had been in corn monoculture over 50 years resulted in increased microbial community richness, diversity, and new microbial species within eight weeks (Maul and Drinkwater, 2010). Addition of hairy vetch (*Vicia villosa*) and cereal rye (*Secale cereal*) to a tomato production system increased the absolute microbial biomass but increased available C from the cover crops reduced Gram-positive bacteria (Buyer et al., 2010). In the same study the microbial biomass under hairy vetch was greater than that of bare soil in May and illustrated the effect of root exudation and turnover (Buyer et al., 2010). Using phospholipid fatty acid (PLFA) analysis Zak et al. (2003) observed increased plant diversity altered heterotrophic microbial community composition, and increased the supply of soil N (Zak et al., 2003). Some plant species such as Velvetleaf (*Abutilon*

theophrasti), Common Lambsquarter (*Chenopodium album*), Redroot Pigweed (*Amaranthus retroflexus*), and Green Foxtail (*Setaria viridis*) can be unwanted in corn systems, and when growth is left uncontrolled in the spring can have a deleterious effect on the abundance of arbuscular mycorrhizal fungi (AMF) (Wortman et al., 2013). Plant roots can also serve as a habitat to soil microbes in addition to plant anchorage and nutrient acquisition. Root surface area has been found as a good predictor of microbial diversity and community structure using terminal restriction fragment length polymorphism targeting the 16S rRNA gene (Maul and Drinkwater, 2010).

Influence of Soil Properties on Microbial Community Composition

Soil physiochemical factors such as pH can influence microbial community composition and function. Using PCR-DGGE Wakelin et al. (2008) observed decreasing catabolism of C-substrates with declining pH levels between fine and coarse-textured soils. Plants utilizing NH_4^+ release H^+ in the process, which can decrease the pH of the rhizosphere (Abbott and Murphey, 2003). Nutrient management practices that influence pH may therefore impact microbial dynamics. In contrast, Buyer et al. (2010) did not find soil pH to be a significant factor in microbial community composition in bulk soil. A difference of 1.1 and 0.6 pH units in different years explained only 2.7% and 1.5% of microbial community composition variance in bulk soil (Buyer et al., 2010). Soil nutrient levels may also affect soil microbial community composition. Redundancy analysis has indicated that soil pH, P, K, Ca, $\text{NH}_4\text{-N}$, and OM concentrations have correlated to soil bacterial community structure in soybean, canola, sunflower, and switchgrass rhizosphere soils at four Michigan locations (da C. Jesus et al., 2010).

Rhizosphere Soil

The rhizosphere is an area of soil in the vicinity and influence of plant roots consisting of the endorhizosphere, rhizoplane, and ectorhizosphere and extends 1-2 mm the root surface (Prashar et al., 2014). Bulk soil is the volume not influenced by plant roots. Plant roots are able to congregate microbial communities through the exudation of rhizodeposits, growth, respiration, and nutrient exchange making the rhizosphere a “hot spot” of microbial activity in relation to the surrounding bulk soil (Mahmood et al., 2005; Prashar et al., 2014). Rhizodeposits are carbon-(C) rich allowing microbes to utilize as a source of energy or be inhibited by them (Mahmood et al., 2005; Neumann and Römheld, 2007). Rhizodeposits include sloughed root cells and lysates, border cells, and root exudates in the form of sugars, amino acids and amines, aliphatic-, aromatic-, and fatty acids, sterols, enzymes, miscellaneous vitamins, and plant growth regulators (Neumann and Römheld, 2007). As opposed to bulk soil the rhizosphere may contain bacteria, actinomycetes, and fungi in 2-20, 5-10, and 10-20 proportions, respectively (Morgan et al., 2005). Actinomycetes are a member of the Actinobacteria phylum and contain subgroups of *Streptomyces* associated with crop disease suppressiveness (Wiggins and Kinkel, 2005; Mendes et al., 2011). Microbial activity may be up to 50 times higher in rhizosphere soil as compared to the bulk soil (Prasher et al., 2014). A corn rhizosphere soil was found to contain 3.6x more microbial biomass and 1.2x higher bacterial population density relative to surrounding bulk soil demonstrating higher C-substrate availability (Mahmood et al., 2005). Rhizosphere bacterial phyla found most abundant in a canola, corn, soybean, sunflower rotation were Acidobacteria, Proteobacteria, Actinobacteria, and Verrucomicrobia (da C. Jesus et al., 2010). Bacterial population density was found to be higher on corn and wheat roots relative to the corresponding rhizosphere soil and lowest in the bulk soil which suggests similar soil samples contaminated

with unequal amounts of root biomass may have contrasting amounts of bacterial populations (Mahmood et al., 2005).

Plant Growth Promoting Rhizobacteria

Bacteria beneficial to plant growth and productivity have been termed Plant Growth Promoting Rhizobacteria (PGPR) and can colonize plant root surfaces in the rhizosphere, rhizoplane, and within plant tissue (endophytes) (Tilak et al., 2005). Colonization of PGPRs can be symbiotic or non-symbiotic (includes free-living, associative, or endophytic) and both groups are capable of N fixation (Tilak et al., 2005). Symbiotic nitrogen fixing PGPRs include cyanobacteria of the genera *Rhizobium*, *Bradyrhizobium*, *Azorhizobium*, *Allorhizobium*, *Sinorhizobium*, and *Mesorhizobium* (Hayat et al., 2010). Free-living nitrogen fixing PGPRs have included bacterial species of *Azospirillum*, *Enterobacter*, *Klebsiella*, and *Pseudomonas* (Hayat et al., 2010). Plant growth promotion by symbiotic and non-symbiotic PGPRs can be direct through the fixation of atmospheric N, phosphorus solubilization, and production of plant growth hormones such as auxins, cytokinins, gibberellins, ethylene, and abscisic acid (Tilak et al., 2005). Indirectly, PGPRs enhance plant growth by siderophore production for Fe uptake and provide protection from deleterious phytopathogenic organisms (Hayat et al., 2010). In vitro studies have found *Pseudomonas* spp. produced more hydrogen cyanide, a phytopathogen inhibitor, than *Bacillus* spp. in addition to an increase in siderophore production of 2.87 mm 5 day⁻¹ (Jarak et al., 2012). Symbiotic PGPR reside inside plant cells and produce nodules often in association with leguminous plants although non-legumes (i.e., other crops in rotation) may benefit from fixed N (Hayat et al., 2010).

Non-leguminous plants such as corn, wheat, rice, sugarcane, and cotton can benefit from PGPR effects of non-symbiotic diazotrophs through increased vegetative growth and yield (Kennedy et al., 2004). For example corn dry weight was increased 49.4-81.8% ($P < 0.05$) in the presence of fertilizer N after 6 weeks of growth in greenhouse studies using seed inoculated with *H. seropedicae* in non-sterilized soils as compared to an un-inoculated control. However, when N was withheld dry weight increased an average 15.6%, illustrating the potential of diazotrophic bacteria to increase NUE for biomass production in corn (Riggs et al., 2001). Field studies in Africa utilizing corn seed inoculated with a combination of PGPR's including *Azotobacter chroococcum*, *Pseudomonas* spp., and *Bacillus* spp. increased V3-6 corn dry weight 61.3% and yield $> 1000 \text{ kg ha}^{-1}$ as compared to a non-inoculated control and concluded that co-inoculation increased dry matter production over a single inoculation of each PGPR (Jarak et al., 2012).

LITERATURE CITED

LITERATURE CITED

- Abbott, L.K., and D.V. Murphy. 2003. Soil biological fertility: A key to sustainable land use in agriculture. Kluwer Academic Publ., The Netherlands.
- Aira, M., M. Gómez-Brandón, C. Lazcano, E. Bååth, and J. Domínguez. 2010. Plant genotype strongly modifies the structure and growth of maize rhizosphere microbial communities. *Soil Biol. Biochem.* 42:2276-2281.
- Andraski, T.W., and L.G. Bundy. 2005. Cover crop effects on corn yield response to nitrogen on an irrigated sandy soil. *Agron. J.* 97:1239-1244.
- Andraski, T.W., L.G. Bundy, and K.R. Brye. 2000. Crop management and corn nitrogen rate effects on nitrate leaching. *J. Environ. Qual.* 29:1095-1103.
- Asghari, M., and R.G. Hanson. 1984. Climate, management, and N effect on corn leaf N, yield, and grain N. *Agron. J.* 76:911-916.
- Awasthi, A., M. Singh, S.K. Soni, R. Singh, and A. Kalra. 2014. Biodiversity acts as insurance of productivity of bacterial communities under abiotic perturbations. *ISME J.* 8:2445-2452.
- Bender, R.R., J.W. Haegerle, M.L. Ruffo, and F.E. Below. 2013. Nutrient uptake, partitioning, and remobilization in modern, transgenic, insect-protected maize hybrids. *Agron. J.* 105:161-170.
- Berg, G., and K. Smalla. 2009. Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *FEMS Microb. Ecol.* 68:1-13.
- Bermudez, M., and A.P. Mallarino. 2002. Yield and early growth responses to starter fertilizer in no-till corn assessed with precision agriculture technologies. *Agron. J.* 94:1024-1033.
- Bermudez, M., and A.P. Mallarino. 2004. Corn response to starter fertilizer and tillage across and within fields having no-till management histories. *Agron. J.* 96:776-785.
- Blackmer, T.M., and J.S. Schepers. 1995. Use of a chlorophyll meter to monitor nitrogen status and schedule fertigation for corn. *J. Prod. Agric.* 8:56-60.
- Binder, D.L., D.H. Sander, and D.T. Walters. 2000. Maize response to time of nitrogen application as affected by level of nitrogen deficiency. *Agron. J.* 92:1228-1236.
- Bollero, G.A., D.G. Bullock, and S.E. Hollinger. 1996. Soil temperature and planting date effects on corn yield, leaf area, and plant development. *Agron. J.* 88:385-390.

- Brouder, S.M., D.B. Mengel, and B.S. Hofmann. 2000. Diagnostic efficiency of the blacklayer stalk nitrate and grain nitrogen tests for corn. *Agron. J.* 92:1236-1247.
- Bundy, L.G., and T.W. Andraski. 1999. Site-specific factors affecting corn response to starter fertilizer. *J. Prod. Agric.* 12:664-670.
- Bundy, L.G. 2006. How can we improve nitrogen use efficiency? In: *Proc. Wis. Fert. Aglime Pest Mgmt. Conf.* Madison, WI. 17-19 Jan. 2006. p. 54-60.
- Buyer, J.S., J.R. Teasdale, D.P. Roberts, I.A. Zasada, and J.E. Maul. 2010. Factors affecting soil microbial community structure in tomato cropping systems. *Soil Bio. and Biochem.* 42:831-841.
- Cassman, K.G., A.R. Dobermann, and D.T. Walters. 2002. Agroecosystems, nitrogen-use efficiency, and nitrogen management. *Ambio.* 31:132-140.
- Ceja-Navarro, J.A., F.N. Rivera-Orduña, L. Patiño-Zúñiga, A. Vila-Sanjurjo, J. Crossa, B. Govaerts, and L. Dendooven. 2010. Phylogenetic and multivariate analyses to determine the effects of different tillage and residue management practices on soil bacterial communities. *App. Environ. Micro.* 76:3685-3691.
- Chaparro, J.M., A.M. Sheflin, D.K. Manter, and J.M. Vivanco. 2012. Manipulating the soil microbiome to increase soil health and plant fertility. *Biol. Fertil.* 48:489-499.
- Clark, A. 2007. *Managing cover crops profitably*. 3rd ed. Sustainable Agric. Network, Beltsville, MD.
- Congreves, K.A., L.L. Van Eerd. 2015. Nitrogen cycling and management in intensive horticultural systems. *Nutr. Cycl. Agroecosyst.* 102:299-318.
- Coyne, M.S., and R. Mikkelsen. 2015. Soil microorganisms contribute to plant nutrition and root health. *Better Crops* 99:18-20.
- Crandall, S.M., M.L. Ruffo, and G.A. Bollero. 2005. Cropping system and nitrogen dynamics under a cereal winter cover crop preceding corn. *Plant and Soil* 268:209-219.
- Da C. Jesus, E., E. Susilawati, S.L. Smith, Q. Wang, B. Chai, R. Farris, J.L.M. Rodrigues, K.D. Thelen, and J.M. Tiedje. 2010. Bacterial communities in the rhizosphere of biofuel crops grown on marginal lands as evaluated by 16S rRNA gene pyrosequences. *Bioenerg. Res.* 3:20-27.

- Dapaah, H.K., and T.J. Vyn. 1998. Nitrogen fertilization and cover crop effects on soil structural stability and corn performance. *Commun. Soil Sci. Plant Anal.* 29:2557-2569.
- Dinnes, D.L., D.L. Karlen, D.B. Jaynes, T.C. Kaspar, J.L. Hatfield, T.S. Colvin, and C.A. Cambardella. 2002. Nitrogen management strategies to reduce nitrate leaching in tile-drained Midwestern soils. *Agron. J.* 94:153-171.
- Doran, J.W., and M.R. Zeiss. 2000. Soil health and sustainability: Managing the biotic component of soil quality. *Appl. Soil. Ecol.* 15:3-11.
- Dove, A., and S.C. Chapra. 2015. Long-term trends of nutrients and trophic response variables for the Great Lakes. *Limnol. Oceanogr.* 60:696-721.
- Eisenhauer, N., S. Scheu, and A. Jousset. 2012. Bacterial diversity stabilizes community productivity. *PLoS ONE* 7:e34517.
- Evans, L.T., and R.A. Fischer. 1999. Yield potential: Its definition, measurement, and significance. 39:1544-1551.
- Fernández, F. G., Nafziger, E. D., Ebelhar, S. A., & Hoef, R.G. 2009. Managing nitrogen. p. 113-132. In *Illinois agronomy handbook*. Univ. Illinois Coop. Ext. Serv., Urbana-Champaign.
- Figuerola, E.L.M., L.D. Guerrero, D. Türkowsky, L.G. Wall, and L. Erijman. 2015. Crop monoculture rather than agriculture reduces the spatial turnover of soil bacterial communities at a regional scale. *Environ. Micro.* 17:678-688.
- Forrestal, P.J., R.J. Kratochvil, and J.J. Meisinger. 2012. Late-season corn measurements to assess soil residual nitrate and nitrogen management. *Agron. J.* 104:148-157.
- Fowler, D.B., and J. Brydon. 1989. No-till winter wheat production on the Canadian prairies: placement of urea and ammonium nitrate fertilizers. *Agron. J.* 81:518-524.
- Fox, R.H., and L.D. Hoffman. 1981. The effect of N fertilizer source on grain yield, N uptake, soil pH, and lime requirements in no-till corn. *Agron. J.* 73:891-895.
- Fox, R.H., W.P. Piekielek, and K.E. Macneal. 2001. Comparison of late-season diagnostic tests for predicting nitrogen status of corn. *Agron. J.* 93:590-597.
- Gagnon, B., and N. Ziadi. 2010. Grain corn and soil nitrogen responses to sidedress nitrogen sources and application. *Agron. J.* 102:1014-1022.

- Gagnon, B., N. Ziadi, and C. Grant. 2012. Urea fertilizer forms affect grain corn yield and nitrogen use efficiency. *Can. J. Soil Sci.* 92:341-351.
- Gehl, R.J., J.P. Schmidt, L.D. Maddux, and W.B. Gordon. 2005. Corn yield response to nitrogen rate and timing in sandy irrigated soils. *Agron. J.* 97:1230-1238.
- Gerwing, J., R. Gelderman, and A. Bly. 1996. Effects of seed placed P studied. *Fluid J.* 4:14-15.
- Gieske, M.F., D.L. Wyse, and B.R. Durgan. 2016a. Spring- and Fall-seeded radish cover-crop effects on weed management in corn. *Weed Tech.* 30:559-572.
- Gieske, M.F., V.J. Ackroyd, D.G. Bass, D.R. Mutch, D.L. Wyse, and B.R. Durgan. 2016b. Brassica cover crop effects on nitrogen availability and oat and corn yield. *Agron. J.* 108:151-161.
- Guillard, K., T.F. Morris, and K.L. Kopp. 1999. The pre-sidedress soil nitrate test and nitrate leaching from corn. *J. Environ. Qual.* 28:1845-1852.
- Hayat, R, S. Ali, U. Amara, R. Khalid, and I. Ahmed. 2010. Soil beneficial bacteria and their role in plant growth promotion: a review. *Ann. Microbiol* 60:579–598.
- Havlin, J.L., S.L. Tisdale, W.L. Nelson, and J.D. Beaton. 2014. *Soil Fertility and Fertilizers. An Introduction to Nutrient Management.* 8th ed. Pearson Education Inc., Upper Saddle River, NJ.
- Hayhoe, K., C. Wake, T. Huntington, L. Luo, M. Schwartz, J. Sheffield, E. Wood, B. Anderson, J. Bradbury, A. Degaetano, T. Troy, and D. Wolfe. 2007. Past and future changes in climate and hydrological indicators in the U.S. *Clim. Dynam.* 28:381–407.
- Hatfield, J.L., and T.B. Parkin. 2014. Enhanced efficiency fertilizers: Effect on agronomic performance of corn in Iowa. *Agron. J.* 106:771-780.
- Hawkins, J.A., J.E. Sawyer, D.W. Barker, and J.P. Lundvall. 2007. Using relative chlorophyll meter values to determine nitrogen application rates for corn. *Agron. J.* 99:1034-1040.
- Hong, N., P.C. Scharf, J.G. Davis, N.R. Kitchen, and K.A. Sudduth. 2007. Economically optimal nitrogen rate reduces soil residual nitrate. *J. Environ. Qual.* 36:354-362.
- Jahanzad, E., A.V. Barker, M. Hashemi, T. Eaton, A. Sadeghpour, and S.A. Weis. 2016. Nitrogen release dynamics and decomposition of buried and surface cover crop residues. *Agron. J.* 108:1735-1741.

- Jarak, M., N. Mrkovački, D. Bjelić, D. Jošić, T. Hajnal-Jafari, and D. Stamenov. 2012. Effects of plant growth promoting rhizobacteria on maize in greenhouse and field trial. *African J. of Micro. Res.* 6:5683-5690.
- Johansen, J. E., and S. J. Binnerup. 2002. Contribution of Cytophaga-like bacteria to the potential of turnover of carbon, nitrogen, and phosphorus by bacteria in the rhizosphere of barley (*Hordeum vulgare* L.). *Microb. Ecol.* 43:298–306.
- Johnson, T.J., T.C. Kaspar, K.A. Kohler, S.J. Corak, and S.D. Logsdon. 1998. Oat and rye overseeded into soybean as fall cover crops in the upper Midwest. *Soil Water Conserv.* 53:276-279.
- Jokela, W.E., and G.W. Randall. 1997. Fate of fertilizer nitrogen as affected by time and rate of application in corn. *Soil Sci. Soc. Am. J.* 61:1695-1703.
- Kaiser, D.E., A.P. Mallarino, and M. Bermudez. 2005. Corn grain yield, early growth, and early nutrient uptake as affected by broadcast and in-furrow starter fertilization. *Agron. J.* 97:620-626.
- Kaiser, D.E., and J.C. Rubin. 2013. Corn nutrient uptake as affected by in-furrow starter fertilizer for three soils. *Agron. J.* 105:1199-1210.
- Karl, T.R., J.M. Melillo, and T.C. Peterson (ed.). 2009. *Global climate change impacts in the United States.* Cambridge Univ. Press, New York.
- Karlen, D.L., L.A. Kramer, and S.D. Logsdon. 1998. Field-scale nitrogen balances associated with long-term continuous corn production. *Agron. J.* 90:644-650.
- Karlen, D.L., R.L. Flannery, and E.J. Sadler. 1988. Aerial accumulation and partitioning of nutrients by corn. *Agron. J.* 80:232-242.
- Kaspar, T.C., and J.W. Singer. 2011. The use of cover crops to manage soil. p. 321-327. In Hatfield and Sauer (ed.) *Soil management: building a stable base for agriculture.* American Society of Agronomy, Madison, WI.
- Kennedy, I.R., A.T.M.A. Choudhury, and M.L. Kecskés. 2004. Non-symbiotic bacterial diazotrophs in crop-farming systems; can their potential for plant growth promotion be better explained? *Soil Biol. Biochem.* 36: 1229–1244.
- Kyveryga, P.M., H. Tao, T.F. Morris, and T.M. Blackmer. 2010. Identification of nitrogen management categories by corn stalk nitrate sampling guided by aerial imagery. *Agron. J.* 102:858-867.

- Ladha, J.K., H. Pathak, T.J. Krupnik, J. Six, and C. van Kessel. 2005. Efficiency of fertilizer nitrogen in cereal production: Retrospects and prospects. *Advances in Agron.* 87:85-176.
- Lambers, H., C. Mougel, B. Jaillard, and P. Hinsing. 2009. Plant-microbe-soil interactions in the rhizosphere: an evolutionary perspective. *Plant Soil* 321:83-115.
- Lauer, J.G., P.R. Carter, T.M. Wood, G. Diezel, D.W. Wiersman, R.E. Rand, and M.J. Mlynarek. 1999. Corn hybrid response to planting date in the Northern Corn Belt. *Agron. J.* 91:834–839.
- Lawley, Y.E., R.R. Weil, and J.R. Teasdale. 2011. Forage radish cover crop suppresses winter annual weeds in fall and before corn planting. *Agron. J.* 103:137-144.
- Li, Q., X. Song, H. Gu, and F. Gao. 2016. Nitrogen deposition and management practices increase soil microbial biomass carbon but decrease diversity in Moso bamboo plantations. *Scientific Reports* 6:1-11.
- Lohry, R.D. 1989. Effect of N fertilizer rate and nitrapyrin on leaf chlorophyll, leaf N concentration, and yield of three irrigated maize hybrids in Nebraska. Ph.D. diss. Univ. of Nebraska, Lincoln.
- Mahmood, T., W.M. Kaiser, R.Ali, M. Ashraf, A. Gulnaz, and Z. Iqbal. 2005. Ammonium versus nitrate nutrition of plants stimulates microbial activity. *Plant Soil* 277:233-243.
- Maul, J., and L. Drinkwater. 2010. Short-term plant species impact on microbial community structure in soils with long-term agricultural history. *Plant Soil* 330:369-382.
- McSwiney, C.P., S.G. Snapp, and L.E. Gentry. 2010. Use of immobilization to tighten the N cycle in conventional agroecosystems. *Ecol. Appl.* 20:648-662.
- Mendes, R., M. Kruijt, I. de Bruijn, E. Dekkers, M. van der Voort, J.H.M. Schneider, Y.M. Piceno, T.Z. DeSantis, G.L. Andersen, P.A.H.M. Bakker, and J.M. Raaijmakers. 2011. Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science* 332:1097-1100.
- Mikkelsen, R.L. 2011. The “4R” nutrient stewardship framework for horticulture. *Hort. Tech.* 21:658-662.
- Millar, N., G.P. Robertson, P.R. Grace, R.J. Gehl, and J.P. Hoben. 2010. Nitrogen fertilizer management for nitrous oxide (N₂O) mitigation in intensive corn (Maize) production: An emissions reduction protocol for US Midwest agriculture. *Mitig. Adapt. Strateg. Glob. Change* 15:185-204.

- Morgan, J.A.W., G.D. Bending, and P.J. White. 2005. Biological costs and benefits to plant–microbe interactions in the rhizosphere. *J. Exp. Bot.* 56: 1729–1739.
- Mutch, D.R., and K.D. Thelen. 2003. No-till drilling cover crops after wheat harvest and their influence on next season’s corn. Michigan State University Ext. Bull. E-2897.
- Mulvaney, R.L., S.A. Khan, and T.R. Ellsworth. 2006. Need for soil-based approach in managing nitrogen fertilizers for profitable corn production. *Soil Sci. Am. J.* 70:172-182.
- Nelson, D.W. and Huber, D. 2001. Nitrification inhibitors for corn production. Iowa State University Extension. NCH 55.
- Neumann, G., and V. Römheld. 2007. The release of root exudates as affected by the plant physiological status. p. 23-72. In R. Pinton et al. (ed.) *The rhizosphere: Biochemistry and organic substances at the soil-plant interface*. 2nd ed. Taylor and Francis Group, LLC, Florida.
- Niehues, B.J., R.E. Lamond, C.B. Godsey, and C.J. Olsen. 2004. Starter nitrogen fertilizer management for continuous no-till corn production. *Agron J.* 96:1412-1418.
- Nihorimbere, V., M. Ongena, M. Smargiassi, and P. Thonart. 2011. Beneficial effect of the rhizosphere microbial community for plant growth and health. *Biotechnol. Agron. Soc. Environ.* 15:327-337.
- Oades, J.M. 1993. The role of biology in the formation, stabilization, and degradation of soil structure. *Geoderma* 56:377-400.
- Oberle, S.L., and D.R. Keeney. 1990. Factors influencing corn fertilizer N requirements in the northern U.S. Corn Belt. *J. Prod. Agric.* 3:527-534.
- O'Reilly, K.A., J.D. Lauzon, R.J. Vyn, and L.L. Van Erd. 2012. Nitrogen cycling, profit margins and sweet corn yield under fall cover crop systems. *Can. J. Soil Sci.* 92:353-365.
- Piekielek, W.P., R.H. Fox, J.D. Toth, and K.E. Macneal. 1995. Use of a chlorophyll meter at the early dent stage of corn to evaluate nitrogen sufficiency. *Agron. J.* 87:403-408.
- Prashar, P., N. Kapoor, and S. Sachdeva. 2014. Rhizosphere: its structure, bacterial diversity and significance. *Rev. Environ. Sci. Biotechnol* 13:63-77.
- Randall, G.W., and R.G. Hoefl. 1988. Placement methods for improved efficiency of P and K fertilizers: A review. *J. Prod. Agric.* 1:70-79.

- Randall, G.W., and D.J. Mulla. 2001. Nitrate nitrogen in surface waters as influenced by climatic conditions and agricultural practices. *J. Environ. Qual.* 30:337-344.
- Randall, G.W., and J. Sawyer. 2008. Nitrogen application timing, forms, and additives. In *Gulf Hypoxia and Local Water Quality Concerns Workshop*, Ames, IA, 27-28 Sept. 2005, p. 73-84.
- Randall, G.W., and J.A. Vetsch. 2005. Corn production on a subsurface-drained Mollisol as affected by fall versus spring application of nitrogen and nitrapyrin. *Agron. J.* 97:472-478.
- Randall, G.W., J.A. Delgado, and J.S. Schepers. 2008. Nitrogen management to protect water resources. p. 911-945. In *Nitrogen in Agriculture, Monograph 49; Nitrogen in Agricultural Systems*. J.S. Schepers (ed.). SSSA, Madison, WI.
- Raun, W.R., D.H. Sander, and R.A. Olson. 1986. Emergence of corn as affected by source and rate of solution fertilizers applied with the seed. *J. Fert. Issues* 3:18-24.
- Raun, W.R., and G.V. Johnson. 1999. Improving nitrogen use efficiency for cereal production. *Agron. J.* 91:357-363.
- Rehm, G.W., and J.A. Lamb. 2009. Corn response to fluid fertilizers placed near the seed at planting. *Soil Sci. Soc. Am. J.* 73:1427-1434.
- Richardson, A.D., S.P. Duigan, and G.P. Berlyn. 2002. An evaluation of noninvasive methods to estimate foliar chlorophyll content. *New Phytol.* 153:185-194.
- Riggs, P.J., M.K. Chelious, A.L. Inguez, S.M. Kaeppler, and E.W. Triplett. 2001. Enhanced maize productivity by inoculation with diazotrophic bacteria. *Aust. J. Plant Physiol.* 28:8-14.
- Ritchie, K.B., R.G. Hoefl, E.D. Nafziger, L.C. Gonzini, and J.J. Warren. 1996. Starters perform more consistently than seed-placed fertilizers. *Fluid J. Winter* 1996:1-3.
- Roberts, T.L. 2007. Right product, right rate, right time, and right place...the foundation of BMP's for fertilizer. *Better Crops* 91:14-15.
- Robertson, G.P., and P.M. Vitousek. 2009. Nitrogen in agriculture: balancing an essential resource. *Annual Rev. Energy Environ.* 34:97-125.
- Rorie, R.L., L.C. Purcell, M. Mozaffari, D.E. Karcher, C.A. King, M.C. Marsh, and D.E. Longer. 2011. Association of "greenness" in corn with yield and leaf nitrogen concentration. *Agron. J.* 103:529-535.

- Samborski, S.M., N. Tremblay, and E. Fallon. 2009. Strategies to make use of plant sensors-based diagnostic information for nitrogen recommendations. *Agron. J.* 101:800-816.
- Sanchez, C.A., and A.M. Blackmer. 1988. Recovery of anhydrous ammonia-derived nitrogen-15 during three years of corn production in Iowa. *Agron. J.* 80:102-108.
- Sawyer, J., E. Nafziger, G. Randall, L. Bundy, G. Rehm, and B. Joern. 2006. Concepts and rationale for regional nitrogen rate guidelines for corn. Publ. PM2015, Iowa State University Extension, Ames, IA.
- Scavia, D., J.D. Allan, K.K. Arend, S. Bartell, D. Beletsky, N.S. Bosch, S.B. Brandt, R.D. Briland, I. Daloglu, J. V. DePinto, D.M. Dolan, M.A. Evans, T.M. Farmer, D. Goto, H. Han, T.O. Höök, R. Knight, S.A. Ludsin, D. Mason, A.M. Michalak, R.P. Richards, J.J. Roberts, D.K. Rucinski, E. Rutherford, D.J. Schwab, T.M. Sesterhenn, H. Zhang, and Y. Zhou. 2014. Assessing and addressing the re-eutrophication of Lake Erie: Central basin hypoxia. *J. Great Lakes Res.* 40, 226–246.
- Scharf, P.C., W.J. Wiebold, and J.A. Lory. 2002. Corn yield response to nitrogen fertilizer timing and deficiency level. *Agron. J.* 94:435-441.
- Schepers, J.S., D.D. Francis, M. Vigil, and F.E. Below. 1992. Comparison of corn leaf nitrogen concentration and chlorophyll meter readings. *Commun. Soil Sci. Plant Anal.* 23:2173-2187.
- Schröder, J.J., J.J. Neeteson, O. Oenema, and P.C. Struik. 2000. Does the crop of the soil indicate how to save nitrogen in maize production? Reviewing the state of the art. *Field Crops Res.* 66:151-164.
- Schultze, S.R., P. Sabbatini, and L. Luo. 2016. Effects of a warming trend on cool climate viticulture in Michigan, USA. *SpringerPlus* 5:1119.
- Schwartz, M.D., R. Ahas, and A. Aasa. 2006. Onset of spring starting earlier across the Northern Hemisphere. *Glob. Change Biol.* 12:343–351.
- Silva, G. 2011. End of season corn stalk nitrate test [Online]. Michigan State University Extension. (http://msue.anr.msu.edu/news/end_of_season_corn_stalk_nitrate_test). (posted 24 Aug. 2011; verified 12 Jan. 2015).
- Snapp, S.S., S.M. Swinton, R. Labarta, D. Mutch, J.R. Black, R. Leep, J. Nyiraneze, and K. O'Neill. 2005. Evaluating cover crops for benefits, costs and performance within cropping system niches. *Agron. J.* 97:322–332.
- Soman, C. D. Li, M.M. Wander, and A.D. Kent. 2017. Long-term fertilizer and crop-rotation treatments differentially affect soil bacterial community structure. *Plant Soil* 413:145-159.

- Stanford, G. 1973. Rationale for optimum nitrogen fertilization in corn production. *J. Environ. Qual.* 2:159-166.
- Steinke, K., and A. Chomas. 2014. Corn response to polymer-coated urea application timing, nitrogen rate, and blending ratio. Poster presented at: Slow/controlled release fertilizer technology. ASA, CSSA, and SSSA Annual Meetings, Long Beach, CA. 2-5 Nov. Poster 1240.
- Stevens, W.B., R.G. Hoefl, and R.L. Mulvaney. 2005a. Fate of nitrogen-15 in a long-term nitrogen rate study: I. Interactions with soil nitrogen. *Agron. J.* 97:1037-1045.
- Stevens, W.B., R.G. Hoefl, and R.L. Mulvaney. 2005b. Fate of nitrogen-15 in a long-term nitrogen rate study: II. Nitrogen uptake efficiency. *Agron. J.* 97:1046-1053.
- Subedi, K.D., and B.L. Ma. 2005. Effects of N-deficiency and timing of N supply on the recovery and distribution of labeled ¹⁵N in contrasting maize hybrids. *Plant and Soil* 273:189-202.
- Sun, B., S. Jia, S. Zhang, N.B. McLaughlin, A. Liang, X. Chen, S. Liu, and X. Zhang. 2016. No tillage combined with crop rotation improves soil microbial community composition and metabolic activity. *Environ. Sci. Pollut. Res.* 23:6472-6482.
- Teal, R.K., B. Tubana, K. Girma, K.W. Freeman, D.B. Arnall, O. Wash, and W.R. Raun. 2006. In-season prediction of corn grain yield potential using normalized difference vegetation index. *Agron. J.* 98:1488-1494.
- Tebaldi, C., K. Hayhoe, J.M. Arblaster, and G.E. Meehl. 2006. Going to the extremes: An intercomparison of model simulated historical and future changes in extreme events. *Clim. Change* 79:185-211.
- Thomas, B.W., X. Hao, F.J. Larney, C. Goyer, M.H. Chantigny, and A. Charles. 2017. Non-legume cover crops can increase non-growing season nitrous oxide emissions. *Soil Sci. Soc. Am. J.* 81:189-199.
- Thorup-Kristensen, K. 2001. Are differences in root growth of nitrogen catch crops important for their ability to reduce soil nitrate-N content, and how can this be measured? *Plant and Soil* 230:185-195.
- Tilak, K.V.B.R., N. Ranganayaki, K.K. Pal, R. De, A.K. Saxena, C.S. Nautiyal, S. Mittal, A.K. Tripathi, and B.N. Johri. 2005. Diversity of plant growth and soil health supporting bacteria. *Current Sci.* 89:136-150.

- Tremblay, N., Y.M. Bouroubi, C. Bélec, R.W. Mullen, N.R. Kitchen, W.E. Thomason, S. Ebelhar, D.B. Mengel, W.R. Raun, D.D. Francis, E.D. Vories, and I. Ortiz-Monasterio. 2012. Corn response to nitrogen is influenced by soil texture and weather. *Agron. J.* 104:1658-1671.
- Tucker, C.J. 1979. Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sens. Environ.* 8:127-150.
- U.S. Environmental Protection Agency. 2017. State of the Great Lakes 2017: Highlights Report. <https://binational.net/2017/06/19/sogl-edgl-2017/>. Accessed 14 Nov. 2017.
- USDA – ERS. 2016. Fertilizer use and price. <https://www.ers.usda.gov/data-products/fertilizer-use-and-price.aspx> (accessed 03 Feb. 2017).
- USDA – NASS. 2017. Crop production 2016 summary. <http://usda.mannlib.cornell.edu/MannUsda/viewDocumentInfo.do?documentID=1047> (accessed 03 Feb. 2017).
- Vanotti, M.B., and L.G. Bundy. 1994. An alternative rationale for corn nitrogen fertilizer recommendations. *J. Prod. Agric.* 7:243-249.
- Varvel, G.E., J.S. Schepers, and D.D. Francis. 1997. Ability for in-season correction of nitrogen deficiency in corn using chlorophyll meters. *Soil Sci. Soc. Am. J.* 61:1233-1239.
- Vetsch, J.A., and G.W. Randall. 2000. Enhancing no-tillage systems for corn with starter fertilizers, row cleaners, and nitrogen placement methods. *Agron. J.* 92:309-315.
- Vetsch, J.A., and G.W. Randall. 2002. Corn production as affected by tillage system and starter fertilizer. *Agron. J.* 94:532-540.
- Vetsch, J.A., and G.W. Randall. 2004. Corn production as affected by nitrogen application timing and tillage. *Agron. J.* 96:502-509.
- Vyn, T.J., J.G. Faber, K.J. Janovicek, and E.G. Beauchamp. 2000. Cover crop effects on nitrogen availability to corn following wheat. *Agron. J.* 92:915-924.
- Vyn, T.J., K.J. Janovicek, M.H. Miller, and E.G. Beauchamp. 1999. Soil nitrate accumulation and corn response to preceding small-grain fertilization and cover crops. *Agron. J.* 91:17-24.
- Wagger, M.G., M.L. Cabrera, and N.N. Ranells. 1998. Nitrogen and carbon cycling in relation to cover crop residue quality. *Soil Water Conserv.* 53:214–218.

- Wakelin, S.A., L.M. Macdonald, S.L. Rogers, A.L. Gregg, T.P. Bolger, and J.A. Baldock. 2008. Habitat selective factors influencing the structural composition and functional capacity of microbial communities in agricultural soils. *Soil Biol. & Biochem.* 40:803-813.
- Walsh, O., W. Raun, A. Klatt, and J. Solie. 2012. Effect of delayed nitrogen fertilization on maize (*Zea mays* L.) grain yields and nitrogen use efficiency. *J. of Plant Nut.* 35:538-555.
- Warncke, D., J. Dahl, and L. Jacobs. 2009. Nutrient recommendations for field crops in Michigan. Bull. E2904, Michigan State University Extension, East Lansing, MI.
- Wiggins, B.E., and L.L. Kinkel. 2005. Green manures and crop sequences influence alfalfa root rot pathogen inhibitory activity among soil-borne streptomycetes. *Plant and Soil* 268:271-283.
- Williams, S.M., and R.R. Weil. 2004. Crop cover root channels may alleviate soil compaction effects on soybean crop. *Soil Sci. Soc. Am. J.* 68:1403-1409.
- Wolkowski, R.P. 2000. Row-placed fertilizer for maize grown with an in-row crop residue management system in southern Wisconsin. *Soil and Tillage Research* 54:55-62.
- Wortman, C.S., S.A. Xerinda, M. Mamo, and C.A. Shapiro. 2006. No-till crop response to starter fertilizer in eastern Nebraska: I. Irrigated and rainfed corn. *Agron. J.* 98:156-162.
- Wortman, S.E., R.A. Drijber, C.A. Francis, and J.L. Lindquist. 2013. Arable weeds, cover crops, and tillage drive soil microbial community composition in organic cropping systems. *Applied Soil Ecol.* 72:232-241.
- Zak, D.R., W.E. Holmes, D.C. White, A.D. Peacock, and D. Tilman. 2003. Plant diversity, soil microbial communities, and ecosystem function: Are they links? *Ecology* 84:2042-2050.

CHAPTER 2

**CORN GROWTH AND YIELD RESPONSES TO PRE-PLANT AND IN-SEASON
NITROGEN COMBINATIONS**

Abstract

In-season nitrogen (N) applications offer flexibility to synchronize availability with corn (*Zea mays* L.) uptake but deliberating between early (V4) and late sidedress [SD] (V11) require further validation due to increased climate variability. A six site-year study investigated N-placement, timing, and source combinations on corn yield, profitability, and growth. Three strategies were investigated involving pre-plant incorporated (PPI), in-furrow [IF] (7.8 kg N ha⁻¹), or sub-surface banded N fertilizer below the furrow [5x5] (44.8 kg N ha⁻¹). Treatment combinations within the IF and 5x5 starter N strategies included sidedress (SD) at V4, V11, or 50/50 (split) V4 and V11. Pre-plant incorporated (PPI) strategies involved 100% urea, 25/75 mix of urea with polymer-coated urea, and poultry manure applied at 2.2 Mg ha⁻¹ plus SD N V11. With the exception of a poultry manure application in wet soils, there was no yield benefit to late season N (i.e., V11) application with similar profitability trends. The IF strategy required reduced N rates at planting and sometimes reduced yield potential when SD was delayed from V4 to V11, unlike the 5x5 strategy. Splitting N application (i.e., multi-pass) increased yield 4.4 to 16.1% compared with a one-pass PPI strategy in 4 of 6 site years. Canopy NDVI at V6 indicated success of SD time may depend on the ability of N strategies to supply N until SD time in order to maintain yield potential. Risk of reduced yield potential late (i.e., V11) was greater than the risk of reduced yield potential early (i.e., V4) suggesting the V11 timing should only be utilized as a rescue application when growers miss a V4 timing to obtain yields ≥ 11.4 Mg ha⁻¹.

Introduction

In the Northern Corn Belt spring preplant N applications are considered a best-management practice in conventionally tilled, poorly drained, medium- to fine-textured soils (Vetsch and Randall, 2004). Corn yield potential is affected by interacting agronomic practices [e.g., fertility management], cultivars, and the environment (Evans and Fischer, 1999). However, recent studies project more heat waves (5°C above climatic normal), increasing air temperatures (1.5 to 2.0°C) over the next 30 years, changing early spring frost dates (1 wk) in the northern hemisphere, and increasing precipitation frequency in winter and spring months with more intense events [> 50.8 mm in 48 hours] (Schwartz et al., 2006; Tebaldi et al., 2006; Hayhoe et al., 2007; Karl et al., 2009). Warming spring temperatures are likely to move corn planting dates earlier than currently recommended [1 and 7 May] (Lauer et al., 1999). Weather variability may increase risk of early applied N loss (immobilization, leaching, denitrification, and clay fixation) and suggests N management strategies should be revisited in response to climate change (Scharf et al., 2002; Scavia et al., 2014; Dove and Chapra, 2015). Optimizing fertility management includes adjusting strategies to account for the right placement, time, source, and rate (e.g., '4R') of nitrogen (N) adapted to site or region-specific environments since corn response to N is mediated by soil texture and weather (Snyder et al., 2009; Tremblay et al., 2012; Norton, 2014; Venterea and Coulter, 2015). Since rapid corn N uptake does not occur until V10, early planting dates combined with spring weather volatility suggest delayed N applications (up to V10) would mitigate the time lapse between N availability and uptake but further research is needed to refine current grower practices (Bender et al., 2013).

One N management strategy used by growers are pre-plant incorporated (PPI) applications. The PPI strategy represents a one-pass system where 100% of the N inputs are applied up to planting time and incorporated. Other one-pass systems include pre-emergent applications (PRE) applied 100% at-plant or within 1 to 3-d after planting (e.g., weed 'n' feed) using urea or urea ammonium nitrate (UAN), and spring pre-plant N applications. In MI, the majority of corn acreage is located on calcareous soils with a soil pH > 7.2 which can potentially increase NH₃ volatilization losses, potential for immobilization, and reduce the efficiency of surface applied urea containing N sources (Vetsch and Randall, 2000; Rawluk et al., 2001; Havlin et al., 2014). In MI studies, urea blended with polymer coated urea (PCU) (75/25 percent PCU/urea blend ratio) have improved corn yields up to 1.38 Mg ha⁻¹ relative to 100% urea PPI (2-4 weeks) when Apr. and May rainfall were above avg (Franzen, 2017). In the same studies a PCU/urea blend (75/25) extended N activity in dry soils which increased corn grain yield 1.07 Mg ha⁻¹ relative to a V4-6 urea surface banded sidedress (SD) application. Despite the recommended use of spring preplant N applications in the Northern Corn Belt, early weather can be volatile with above-average rainfall in April, May, and June, increasing the risk of N loss conditions (Vetsch and Randall, 2004). Single spring pre-plant N applications have reduced yield 0.39 Mg ha⁻¹ relative to split applications in Minnesota and 2.45 Mg ha⁻¹ relative to a single at-plant application in MI (Randall et al., 2003; Franzen, 2017). Reduced efficiency of early applied N has prompted some MI growers to utilize a multi-pass system to improve N recovery but limited MI data utilizing the PPI strategy warrant further research including in-season N applications (Warncke et al., 2009).

In an attempt to improve synchrony between N applications and plant N uptake, grower interest in late N (e.g., post-V11) applications has increased. In-furrow (IF) nutrient placement (i.e.,

"popup" or "seed-placed" starter) is one strategy to provide immediate nutrient access to emerging corn roots as fertilizer is placed with the seed at planting (Niehues et al., 2004). Due to less planter equipment, the IF placement is popular in the Northern Corn Belt (Kaiser et al., 2016). However to prevent yield reductions due to seedling injury or delayed emergence, reduced fertilizer rates and non-urea containing forms of N are used to avoid excessive salt concentrations or ammonia toxicity (Raun et al., 1986; Laboski et al., 2008; Rehm and Lamb, 2009). Michigan IF fertilizer recommendations to corn include $\leq 5.6 \text{ kg N+K}_2\text{O ha}^{-1}$ where CEC is $< 7 \text{ cmol(+) kg}^{-1}$ and $\leq 9.0 \text{ kg N+K}_2\text{O ha}^{-1}$ where CEC is $\geq 8 \text{ cmol(+) kg}^{-1}$ (Steinke, 2013). In-furrow starter fertilizer has increased early season plant height and kernel mass, while decreasing days to silking and grain moisture (Kaiser et al., 2016). Despite increased early season plant growth IF applications have not corresponded well to grain yield ($r = 0.44$) (Bermudez and Mallarino, 2002; 2004). In-furrow starters often contain phosphorus (P) and may contain an N, P, and K combination. In soils testing low in phosphorus (STP) ($< 16 \text{ mg P kg}^{-1}$, Bray-1 test) positive grain yields using IF starter fertilizer were attributed to a P-component as opposed to the N-component when positive yield responses were noted in high STP soils ($> 23 \text{ mg kg}^{-1}$) in Iowa (Bermudez and Mallarino, 2002, 2004). In corn production regions with compressed growing seasons and thus shorter maturity length hybrids (i.e., MI), data on fertilizer strategies that provide minimal early-season N in favor of later applications are minimal and require further investigation.

To encourage early season plant growth and nutrient uptake, sub-surface banded N applications 5 cm below and 5 cm laterally (5x5) relative to the seed furrow is another option for at-plant N applications. The 5x5 placement requires installation of an extra planting coulter that may reduce

planting speeds and can be more affected by soil moisture than IF due to bringing more soil to the surface. However, the 5x5 placement can utilize multiple N sources, and allow growers more flexibility for N rate selection relative to reduced N rates required by IF placements (Niehues et al., 2004). Corn V6 plant biomass and grain yield have increased more consistently with no-till 5x5-placed starter relative to IF placements in Illinois while 5x5 placement (N-P) increased yield 0.82 Mg ha^{-1} relative to a no 5x5 control in six no-tillage strip trials in Missouri (Ritchie et al., 1996; Scharf, 1999). In a Wisconsin study 5x5 placed starter fertilizer increased yields up to 0.7 Mg ha^{-1} relative to a no starter control when combined across multiple tillage regimes (Wolkowski, 2000). When compared to the IF placement positive yield responses to the 5x5 placement have been attributed to corn stand maintenance and increased 5x5 N rates [16.3 to $27.2 \text{ kg N ha}^{-1}$] (Lamond and Gordon, 2001; Bermudez and Mallarino, 2002). Corn response to 5x5 starter may also depend upon the interaction between growing season length and hybrid maturity class. On Missouri silt-loam soils increased early corn growth (6-weeks after planting) and reduced days to silking were influenced by hybrid maturity class, tillage, and 5x5 starter (N-P) application but yield was not affected (Cromley et al., 2003, 2006). Bundy and Andraski (1999) observed that a positive yield and economic response to 5x5 starter was linearly related ($R^2=0.51$) to the sum of hybrid relative maturity (RM) and planting date in Julian days (PD) and most likely when $RM+PD > 235$. Corn responded 33% and 53% of the time to starter fertilizers with relative maturity classes $<$ and $>$ 100 days, respectively, with a mean yield increase of 251 kg ha^{-1} relative to a no 5x5 control (Bundy and Andraski, 1999). When using the 5x5 strategy in a split-application approach (i.e., 5x5 + SD) few data exist to validate deliberation among SD times and require further investigation for use in the northern corn belt.

Growers who choose to split apply often utilize SD applications up until the V6 growth stage but interest in delayed (i.e., post-V6) N applications has increased as a strategy to better synchronize N availability and uptake while reducing N loss (Binder et al., 2000; Scharf et al., 2002). Corn N uptake does not begin to accelerate until V6 to V8 (Bender et al., 2013). However weather, planting date, and fertilizer application strategy can affect the N uptake ability of a maturing corn plant and the potential for plant N stress prior to SD time resulting in unrealized yield potential (Bundy and Andraski, 1999; Scharf et al., 2002). When compared to at-plant N applications, N applied at V8 was found to increase N recovery 11% at N rates between 75 - 150 kg N ha⁻¹ (Jokela and Randall, 1997). In Missouri, Scharf et al. (2002) did not observe a yield loss when N was delayed until V11, but noted a 0-3% reduction with later applied N (i.e., V12 to V16). However delayed N applications may have less effect on plant growth and development in the Midwestern U.S. Corn Belt as opposed to northern production regions. Midwest states tend to have protracted growing seasons as compared to northern production regions, where fewer suboptimal growing days due to a compressed growing season and use of shorter relative maturity class corn hybrids may influence N synchrony (Scharf et al., 2002). Nitrogen stress due to late N application emphasizes the importance of satisfying early-season corn N requirements, and the need to identify corn response to delayed sidedress N applications. The objective of this study was to investigate the effects of early- and late-season sidedress N applications in combination with other grower N strategies on early corn growth, corn grain yield, and profitability.

Materials and Methods

Field trials were conducted from 2014-2016 at the Saginaw Valley Research and Extension Center (SVREC) (43°23'58.2"N, -83°41'52.7994"W) near Richville, MI on a Tappan-Londo loam soil (fine-loamy, mixed, active, calcareous, mesic Typic Epiaquolls) and at the South Campus Research Farm (SCRF) (42°40'24.24"N, -84°29'13.1994"W) in Lansing, MI on a Capac loam soil (fine-loamy, mixed, active, mesic Aquic Glossudalf). Fields were autumn chisel plowed following soybean [*Glycine max* (L.) Merr.] harvest, and followed with spring tillage using a soil finisher to a 10-cm depth. Soil samples were collected to a depth of 20 cm prior to fertilizer application, air-dried, and ground to pass through a 2-mm sieve. Soil characteristics at SVREC included 7.6-7.7 pH (1:1 soil/water) (Peters et al., 2015), 19-24 mg kg⁻¹ P (Bray-P1) (Frank et al., 2015), 138-164 mg kg⁻¹ K (ammonium acetate method) (Warncke and Brown, 2015), and 27-28 g kg⁻¹ soil organic matter (loss-on-ignition) (Combs and Nathan, 2015). Soil characteristics at SCRF included 6.5-6.8 pH, 25-47 mg kg⁻¹ P, 91-114 mg kg⁻¹ K, and 28-34 g kg⁻¹ soil OM. Broadcast P and K fertilizer were pre-plant incorporated (10-cm depth) prior to planting as monoammonium phosphate (MAP) (11-52-0 N-P-K) and muriate of potash (MOP) (0-0-62) based on soil tests. Weed control at SVREC consisted of S-metolachlor [Acetamide, 2-chloro-N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl)-,(S)] and glyphosate [N-(phosphonomethyl)glycine] (6 June 2014, 28 May 2015 [glyphosate only], and 9 June 2016) followed by a second application of glyphosate (20 June 2014 and 22 June 2015). Weed control at SCRF consisted of S-metolachlor and glyphosate (10 June 2014), acetochlor [2-chloro-N-ethoxymethyl-N-(2-ethyl-6-methylphenyl) acetamide] and glyphosate (defined previously) (2 June 2015 and 10 June 2016) followed by a second application of glyphosate (26 June 2014, 26 June 2015, and 27 June 2016). Environmental data were collected throughout the year using the

Michigan Automated Weather Network (<http://www.agweather.geo.msu.edu/mawn/>, Michigan State University, East Lansing, Michigan; verified 24 Jul. 2017) from an on-site weather station.

Ten treatments (including a non-fertilized control) were arranged in a randomized complete block with four replications. Plot width was 4.5 m by 12.1 m in length. Treatments were equalized to a total N rate based on the site-specific maximum return to nitrogen rate (MRTN) (202 and 157 kg N ha⁻¹ for SVREC and SCRF, respectively) (Sawyer et al., 2006). Treatments were grouped into three strategies including 1) broadcast pre-plant incorporated (PPI) N, 2) in-furrow (IF) starter N (7.8 kg N ha⁻¹ ammonium polyphosphate [10-34-0]), and 3) starter N sub-surface banded 5 cm beside and 5 cm below the furrow (5x5) (44.8 kg N ha⁻¹ urea ammonium nitrate [UAN; 28-0-0]). Combinations within the IF and 5x5 N strategies included sidedress (SD) at V4, V11, or a 50/50 split at V4 and V11. Pre-plant incorporated N strategies included 100% urea (46-0-0), 25/75 mix of urea with polymer-coated urea (PCU) (44N-0P-0K, Agrium Inc., Calgary, Alberta, Canada), and dried poultry litter (PL) (4-3-2) applied at 2.2 Mg ha⁻¹ plus SD N V11. Corn V4 N was UAN coulter injected 5 cm deep and 38 cm to the side of each row (Ritchie et al., 1997). Corn V11 N was UAN mixed with a urease inhibitor (CO(NH₂)₂ + n-(n-butyl) thiophosphoric triamide] (Koch Agronomic Services, LLC, Wichita, KS) to prevent N volatilization and banded 10 – 15 cm to the side of each row. A non-limiting N reference (280 kg N ha⁻¹) was included to normalize canopy sensor data. An untreated control which did not receive N fertilizer was also included. A Monosem planter (Monosem Inc., Kansas City, Kansas) equipped with Yetter floating planter-unit mounted row cleaners (Yetter Manufacturing, Colchester, IL) and liquid fertilizer applicators was used to apply IF and 5x5 starter. At all sites

Dekalb DKC48-12 RIB (98 d relative maturity) (mid-season hybrid) (Monsanto Co., St. Louis, MO) was planted in 76-cm rows at 84,510 seeds ha⁻¹ (Table 1).

Corn plant density (V3) was determined as number of corn plants in both harvest rows (9.3 m² total area) expressed as plants per ha⁻¹. Canopy normalized difference vegetation index (NDVI) [V6 and V11] was collected using a GreenSeeker® Model 505 handheld red-band optical sensor (Trimble Agriculture Div., Westminster, CO). Nitrogen status using the ear leaf (R1 and R4) was determined with a Minolta SPAD 502 chlorophyll meter (CM) (Konica Minolta, Tokyo, Japan) and normalized to the non-limiting N reference per replicate (Shapiro et al., 2013).

The center two rows of each plot were harvested with a research plot combine to determine grain yield, moisture, and test weight. Yield data were reported at 155 g kg⁻¹ moisture.

Treatment profitability was calculated as net return = gross return from yield – input costs (Table 2).

Data were subject to analysis of variance using the GLIMMIX procedure in SAS assuming fixed effects of site, year, and treatment, and random block effects (SAS Institute, 2011). The UNIVARIATE procedure was used to test for normality of residuals ($P \leq 0.05$). Levene's test was used to investigate homogeneity of error variances using squared- and absolute-values of residuals ($P \leq 0.05$). The LINES option of the slice statement was used to separate lsmeans when ANOVA indicated a significant interaction ($P \leq 0.10$). Multiple degree of freedom (df) contrasts were constructed using the mean of three treatment combinations within each N strategy (e.g., the 5x5 strategy included a mean of three sidedress treatments which received 5x5 starter

fertilizer). Pearson product-moment correlations were used to investigate the relationship of canopy sensor data and weather with grain yield ($P \leq 0.05$) using the CORR procedure.

Results and Discussion

Growing conditions

Total growing season (Apr. – Sept.) precipitation differed by -9, 20 and 1% and 4, -19, and -26% from the 30-yr mean during 2014-2016 at Lansing and Richville, respectively (Table 3). May and June 2015 Lansing precipitation was 24 and 103 mm greater than the 30-yr mean, but Richville 2014-2016 May and June precipitation was deficient by 6-44 mm and 20-52 mm, respectively. August precipitation was 2 – 79 mm above the 30-yr mean across all site years. Mean April air temperatures were below the 30-yr mean across all years and locations while May and June air temperatures were within 0.3 – 1.6 °C at each location. In July 2014 air temperatures were 3.1 – 3.3 °C below the 30-yr mean at each location. August and September mean air temperatures were within 0.5 – 2.2 °C of the 30-yr mean across all site years.

Corn plant density

Means were combined across treatments within each strategy (i.e., IF, PPI, and 5x5) as no SD N applications occurred prior to V3 stand counts. In 2014 the IF strategy reduced plant density up to 2.6% relative to other strategies including the untreated control (Table 4). At the Lansing location dry soil conditions (i.e., 8.9 mm rainfall over 14 d following planting) may have limited diffusion of fertilizer into the soil solution while at the Richville location cool soil temperatures (4.4 °C cooler than Lansing with 5 d of consecutive minimums near 7.2 °C) may have decreased root growth. Either scenario may have increased the amount of time germinating corn roots were

exposed to the fertilizer band and resulted in injury (Laboski et al., 2008). Stand reductions were not observed in other years. Plant density was evaluated as a covariate in the statistical analysis of grain yield, found not significant ($P \leq 0.05$), and was not included in grain yield analysis.

Grain yield

Pearson correlations indicated mean grain yields of treatments receiving N fertilizer increased in association with mean minimum June air temperatures ($r = 0.94$; $P=0.005$) and mean maximum July air temperatures ($r = -0.85$; $P=0.030$). Other studies have reported the influence of air temperature on post-silking dry matter production and growth rate which affect grain yield (Zhou et al., 2016). Weather correlation to yield and treatments receiving N fertilization but not the untreated control emphasizes the impact in-season weather variability may have on N strategy and corn growth.

Grain yield was influenced by the interaction of year, site, and treatment and a significant grain yield response to treatment was observed in 4 of 6 site years (Table 5 and 6). Treatment means were presented separately for each site year. Mean grain yields (i.e., treatments receiving N, 2014-2016) were 11.8 – 14.7 and 11.1 – 15.0 Mg ha⁻¹ at the SCRF and SVRF locations, respectively, with the greatest yields occurring in 2014 at each location. In the two non-significant site years (i.e., SVRF 2015 and SCRF 2016), deficit spring rainfall (i.e., Apr.-June) resulted in dry soils leading to water-stressed corn, reduced mobility of urea-N, and no response to N placement and timing combinations (Venterea and Coulter, 2015; Maharian et al., 2016). For the four significant site years, treatments are discussed by near normal to deficit rainfall (i.e., 3 site years) and above normal rainfall (i.e., 1 site year).

When Apr. – June rainfall was near normal (i.e., SCRF and SVRF 2014) to deficit (i.e., SVRF 2016) with near normal air temperatures, no yields gains occurred with a V11 SD timing (Table 6). Precipitation timing at SVRF in 2016 coincided with SD applications and provided ≥ 6.9 mm cumulative rainfall within 3 d of each SD N application assisting N movement to corn roots and indicated that yield gains with SD still depend upon rainfall. Within the IF strategy, delaying N application from V4 to V11 resulted in similar or 7.6 – 10.1% yield reductions. No statistical differences were observed using the 5x5 strategy with a V4 or V11 SD timing, indicating a more consistent yield response across variable weather conditions and concurs with previous findings (Ritchie et al., 1996). A lack of differences indicated the additional 37 kg N ha⁻¹ provided by the 5x5 strategy was sufficient for attaining similar yields with V4 or V11 SD relative to the IF strategy and emphasizes the importance of considering starter N strategies for unconstrained corn growth when deliberating between early and late SD timings. Relative to a split (50/50) application, SD delayed until V11 resulted in similar or 5.6 – 6.9% yield reductions across both the IF and 5x5 strategies. Yield reductions indicate that starter N (8 to 45 kg N ha⁻¹) was likely exhausted and reduced yield potential due to deficit N supplies by V11 SD. Scharf et al. (2002) suggested that a protracted growing season (e.g., Michigan) may reduce the duration of corn N uptake and may explain full yield potential realization with a V11 SD application in longer growing season climates (e.g., Missouri). Relative to a V4 timing, full SD at V11 was too late to increase yield in near normal to deficit Apr. – Jun. rainfall conditions. Split SD N (50/50) applications did not provide additional yield gains compared to full V4 SD. Reduced yield potential suggests the V11 timing should be utilized as a rescue application only to maintain yields ≥ 11.4 Mg ha⁻¹ when growers miss a V4 timing.

Under the same near normal to deficit rainfall conditions discussed above, pre-plant incorporated PL followed by V11 SD produced similar yields as IF and V4 SD, indicating yield potential was not increased with a slowly mineralizable N source (60.5 kg N ha⁻¹ first year mineralizable-N) (Table 6). Poultry litter was able to extend N activity relative to other early applied N treatments (i.e., IF or 5x5 strategies) and resulted in similar or increased yields (6.6 to 9.7%) when full SD was delayed until V11. When compared to a urea or PCU and urea blends applied PPI, a PL plus V11 SD application achieved similar or increased yields (8.1 to 19.8%), respectively, across site years while the PCU did not provide yield gains as compared to urea alone. The above average June rainfall (32 mm) at the SCRF (2014) and near-normal to deficit May and June rainfall at SVRF (2014, 2016) may have reduced N loss opportunities and provided no benefit to using PCU.

Above normal May – June rainfall (127 mm excess) occurred at SCRF in 2015 resulting in wet soils. In this scenario, pre-plant incorporated PL followed by V11 SD increased yield 8.6 to 13.9% relative to IF or 5x5 starter and V4 SD (Table 6). Moist soil conditions but a lack of heavy rainfall events in May (< 21 mm) may have prevented some degree of early PPI N loss and hindered the effectiveness of PCU. Yield results with PCU were in contrast to Gagnon et al. (2012) where corn yield gains (0.8 to 1.6 Mg ha⁻¹) due to pre-plant incorporated PCU fertilizer in Canada were observed relative to urea fertilizer in a wet spring similar to the current study. However average air temperatures in Lansing, MI were 5 to 6 °C warmer than the Canadian study which may have increased the rate of N release (Franzen, 2017). Although delaying the

majority of N application until V11 may have reduced the opportunity for N losses in wet years, this practice did not increase the opportunity for yield gains in other years.

Multiple degree of freedom (df) contrast statements were constructed to compare means across strategies (e.g., PPI strategy is a mean of urea, PCU and urea, and PL and V11 SD treatments) as well as one pass (i.e., urea and PCU with urea) and multi-pass N (i.e., treatments with SD N) application systems. In 5 of 6 site years the IF and 5x5 strategies achieved similar yields (Table 7). Relative to the PPI strategy, in dry years (i.e., SVRF 2015 and 2016; SCRF 2016) the 5x5 strategy increased yield 5.0 to 9.2% and illustrated the difficulty associated with PPI N uptake in dry soils. Similar yields were obtained with the IF and 5x5 strategy except at the SCRF (2016) where the 5x5 strategy increased yields 5.8%. However, no differences among strategies were observed when May – June rainfall was above normal to excessive (i.e., SCRF 2014 and 2015). Multi-pass N application systems are a university recommended best management practice in MI to improve N recovery (Warncke et al., 2009). In 4 of 6 sites years a multi-pass system increased yields 4.4 to 16.1% relative to a one-pass system. In variable weather conditions yield gains in multi-pass systems suggests improved synchrony of N application with corn uptake. Increased starter N rates with the 5x5 strategy may offer more consistent yield response from increased N supply at V6 as opposed to low N rates applied IF.

Profitability

When total treatment costs were subtracted from treatment gross returns there was a treatment x site x year interaction (Tables 2, 5). Treatment means are presented separately by site and year (Table 7). Across years profitability was correlated with grain yield ($r \geq 0.82$; $P < 0.01$) and total

treatment cost ($r \leq -0.29$; $P \geq 0.01$). A negative correlation with total treatment costs suggests that growers should consider N and application costs in addition to yields when maximizing profitability.

Full SD N applied at V11 did not increase profitability (Table 7). When cumulative Apr. – June rainfall was near normal to deficit, SD delayed from V4 to V11 using 8 kg N ha⁻¹ with the IF strategy resulted in similar or reduced profitability (192 to 195 \$ ha⁻¹) (3 of 6 site years). A similar reduction in profit was observed (99 \$ ha⁻¹) (1 of 6 site years) with the 5x5 strategy under reduced N loss conditions (i.e., SVRF 2014). A reduced frequency of profit loss indicated the 45 kg N per ha⁻¹ applied at planting stabilized profit variability among SD timings. Increased N rates at planting with the 5x5 strategy may allow greater flexibility to achieve similar profitability when deliberating between a V4 and V11 SD application. Despite similar yields split-applied SD (50/50) resulted in similar or reduced (152 \$ ha⁻¹) profitability relative to IF or 5x5 starter plus full V4 SD and suggested no profit benefit to a second SD application. Over two years of optimal growth conditions (i.e., SCRF and SVRF, 2014) a full V11 SD application reduced profitability relative to split-applied SD with starter IF (95 to 117 \$ ha⁻¹) or 5x5 (100 to 105 \$ ha⁻¹). Although similar profits were often achieved with IF and 5x5 plus V4 SD, growers using a V11 SD rescue application were more likely to maintain profitability if preceded by increased N rates in the 5x5 strategy.

Under the same near normal to deficit Apr. – June rainfall conditions discussed above, a PL and V11 SD application resulted in similar or reduced profitability (154 to 252 \$ ha⁻¹) relative to PPI urea or PCU and urea (Table 7). Likewise, PCU and urea resulted in similar or reduced

profitability (98 \$ ha⁻¹) relative to a urea PPI application. Reduced profitability and increased PCU N costs (\$0.42 to 0.54 kg⁻¹ N) relative to urea N emphasizes the risk associated with enhanced efficiency N when N loss conditions are not present (i.e., SVRF 2014). Profitability with a single V11 SD application was most affected in dry years (deficit Apr. – June rainfall) (i.e., SVRF 2015 and 2016; SCRF 2016) and emphasized the importance of considering N costs when deliberating N strategies. For instance, despite similar yields V11 SD N applications that utilized PL sometimes reduced profitability compared to an IF strategy (129 \$ ha⁻¹) and 5x5 strategy (145 to 278 \$ ha⁻¹). At the same time a 5x5 strategy and V11 SD increased profitability (200 to 228 \$ ha⁻¹) relative to the IF strategy and illustrates the increased capacity of an additional 37 kg N ha⁻¹ at planting in a 5x5 strategy to increase profits as compared to 8 kg N ha⁻¹ in the IF strategy. In dry years a PL with V11 SD reduced profitability relative to a V4 SD with IF (190 to 245 \$ ha⁻¹) or 5x5 strategy (186 to 346 \$ ha⁻¹) with fewer apparent differences during near normal rainfall conditions. The PL-N source cost up to 5.7x more than urea- and UAN-N sources, which were both reduced 26 to 38% after 2014 and contributed to the reduced PL profitability in dry years. Despite greater yields in a wet year, similar profits with PL and V11 SD compared to IF or 5x5 starter with V4 SD indicated N savings through improved synchrony of N application and uptake was not sufficient to offset the increased cost of the PL treatment. A lack of profitability gains with a V11 SD suggested post-V4 SD increased risks of reduced profitability as a result of cost and reduced yield.

Profitability between IF, PPI, and 5x5 strategies corresponded to yield data as indicated by multiple df contrasts. When Apr. – June rainfall was near normal to excessive, strategies often resulted in similar profitability (Table 7). However under deficit Apr. – June rainfall conditions,

PPI profitability was reduced relative to the 5x5 (155 to 220 \$ ha⁻¹) and IF strategies (88 to 177 \$ ha⁻¹) and illustrates the difficulty associated with PPI N uptake, grain production, and subsequent profitability in dry soils. In two instances an IF strategy reduced profitability (60 to 140 \$ ha⁻¹) as compared to the 5x5 strategy. In 4 of 6 site years no differences were observed between one and multi-pass N application systems indicating weather variability impacted profitability less than yield. In two contrasting years reduced N loss opportunities increased profitability of a one pass system (122 \$ ha⁻¹) while apparently reduced PPI N uptake in dry soils reduced profitability (201 \$ ha⁻¹) relative to the multi-pass system. Growers often perceive yield loss as a larger risk than profit loss. One pass systems increased the frequency of yield loss while a 5x5 strategy achieved similar or increased profitability relative to other strategies. Results suggest a multi-pass system with a 5x5 strategy may reduce risk of profit loss where weather variability influences yield response.

Plant characteristics

NDVI: Pearson correlations indicated significant relationships between V6 NDVI and yield across years at the Lansing ($r=0.42$; $P<.01$) and Richville ($r=0.76$; $P<.01$) locations, and similar to an r value of 0.46 used to predict grain yield with V8 NDVI in another study (Liu and Wiatrak, 2011). In the current study, positive relationships suggest corn yield increases in relation to NDVI and emphasizes the importance of N management strategies to sufficiently supply N to maintain yield potential until SD time. Active canopy sensing at V6 corn indicated a significant year x treatment and location x treatment interaction and data are presented separately for year and location (Tables 5, 8). NDVI is an indicator of green biomass and has been used to compare plant growth response and N management (Tucker, 1979). When cumulative Apr. –

Jun. rainfall was deficit at both sites (i.e., 2016) or near normal to deficit across years (i.e., SVRF) treatment comparisons were not significant and suggests that deficit rainfall reduces corn growth response to N strategies. No significant differences between SD timings within the IF or 5x5 strategies indicated that corn NDVI had not yet responded to V4 SD N applied 4 to 11-d prior to V6 measurements. In optimal growing conditions with reduced N loss opportunities (i.e., 2014), NDVI increased in corn receiving a full N rate at planting as urea (9.2 to 11.7%) or PCU (6.3 to 8.8%) relative to starter N plus V4 or V11 SD. Multiple df contrast statements showed that NDVI increased with PPI or IF N strategies relative to a 5x5 N placement and suggested increased positional N availability when soils were not dry. Compared to corn receiving no N (i.e., untreated), the IF, 5x5, and PPI N increased corn NDVI (7.8 to 19.2%). Increased NDVI in corn receiving N at planting suggested native soil N supplies were insufficient to achieve similar growth in unfertilized corn. Corn requires nearly 15% of the total N uptake by V6 which may be supplied by soil N mineralization (Bender et al., 2013). Although corn yield potential is set prior to V8, N deficiencies 10 – 42 days after emergence have decreased stem elongation, leaf area, and net photosynthetic rate resulting in less dry matter accumulation and unrealized yield potential emphasizing the importance of strategies to supply sufficient N early in the growing season when photosynthetic rates are high (i.e., $\mu\text{mol CO}_2/\text{m}^2/\text{s}$) (Varvel et al., 1997; Binder et al., 2000; Zhao et al., 2003; Yu et al., 2016).

Active canopy sensing at V11 was not influenced by site or year and data are presented by treatment (Tables 5, 9). When full SD rates were delayed from V4 to V11 the IF strategy reduced NDVI 6.6% as compared to 1.8% with the 5x5 strategy. The larger NDVI reduction with the IF strategy indicated that the 8 kg N ha⁻¹ was not sufficient to supply the required N for both green

biomass and chlorophyll production and suggested less potential to capture photosynthetically active radiation (PAR) (Zhao et al., 2003). However the PL application which had not yet received V11 SD increased NDVI (3.9 to 6.6%) relative to the IF and 5x5 strategies across all site years and suggested increased potential to maintain yield until the V11 SD timing. All strategies increased V11 NDVI relative to unfertilized corn. Increased NDVI in corn receiving PPI N (5.1%) relative to IF N but not 5x5 N suggests N rates > 8 kg N ha⁻¹ were required to improve NDVI but then may affect stand density via saltation.

Chlorophyll content: Chlorophyll meter indices normalized to a non-limiting N control (280 kg N ha⁻¹) at R1 were used to indicate corn N status. Values were influenced by the interaction of year, site, and treatment and data are presented by treatment for each site year (Tables 5, 10). Except for a wet year (i.e., SCRF 2015), relative CM indices indicated reduced plant chlorophyll when SD was delayed from V4 to V11 within both the IF (2.0 to 22%) and 5x5 (4.1 to 9.2%) strategies. Decreased chlorophyll reduced photosynthesis for corn grain production (Hatfield and Parkin, 2014; Yu et al., 2016). Reduced CM values when delaying SD from V4 to V11 indicated less N was sequestered and assimilated into canopy tissues for photosynthesis and corresponded to yield reductions. When May – June rainfall was below normal (i.e., SVRF 2014 – 2016; SCRF 2016), PL with V11 SD often reduced CM values relative to IF or 5x5 starter and V4 SD (5 to 10%) and other PPI's (4.3 to 6.3%) likely due to reduced moisture required for mineralization. In the same years, PL and IF strategies receiving V11 SD resulted in CM values below criterion previously used to trigger N application in MI (≤ 0.96) and indicated N stress (Elwadie et al., 2005). However, studies have indicated that 35% of total corn N uptake occurs post-silking in modern hybrids so despite CM values below a 0.96 threshold, statistically higher grain yields in

all years with PL applications suggested that a slowly mineralizable N source may continue to provide sufficient N for uptake and assimilation during corn reproductive stages (Bender et al., 2013). Across treatment strategies and one- versus multi-pass systems, multiple df contrasts indicated CM values were generally ≥ 0.96 , with few biologically significant differences.

Relative CM indices at R4 were not affected by year or location and treatment means are presented across factors (Tables 5, 11). Studies by Hatfield and Parkin (2014) suggest corn N strategies which increased green leaf area (i.e., stay-green) at the onset of grain-fill enhanced the ability of the plant to capture and convert PAR into greater yield. When a split or full V4 SD was delayed until V11, CM values decreased (6.1 to 8.8%) regardless of IF or 5x5 strategy and corresponded to similar reductions at R1. Despite similar or reduced greenness with the PL and V11 SD relative to the IF and 5x5 starter and V4 or split SD N treatments, similar greenness at R4 (and R1) was not required to attain similar yield. Rather a CM threshold (0.96) corresponded more to reduced yield associated with V11 SD (i.e., IF or 5x5) and urea or PCU/urea applied PPI which indicated N deficiency at R4. A PL application increased greenness between R1 and R4 (numerically) likely due to post-silking N mineralization and uptake. The IF or 5x5 strategies increased CM values (3.2 to 4.3%) in regard to a PPI strategy while a multi-pass system increased relative CM content (4.3%) as compared to a one-pass system. Despite few differences at R1, reduced greenness at R4 with PPI strategies applied in a one pass system suggested less N was sequestered post-silking and reduced opportunities for increased dry matter accumulation (i.e., yield).

Conclusions

When Apr. – June rainfall was near normal to deficit, full SD delayed from V4 to V11 did not result in yield gains. Reduced N rates required by the IF strategy resulted in similar or reduced yields 7.6 – 10.1% and profitability 192 – 195 \$ ha⁻¹ when SD was delayed from V4 to V11. The 5x5 strategy resulted in no yield differences and reduced profitability (99 \$ ha⁻¹) in only 1 of 6 site years when SD was delayed from V4 to V11 which indicated that increased N rates afforded by the 5x5 strategy may stabilize profit and yield variability. Regardless of 5x5 or IF strategy, similar or reduced yields 5.6 – 6.9% with delayed SD (V11) relative to a split (50/50) SD indicated that starter N (8 to 45 kg N ha⁻¹) may have been exhausted due to deficit N supplies, but reduced profitability 152 \$ ha⁻¹ due to the split SD indicate no benefit to a second SD application. Poultry litter did not increase yield potential relative to other strategies (i.e., IF or 5x5 plus V4 SD), but when full SD was delayed until V11 increased yields 6.9 – 9.7% due to PL were offset by the increased N cost and reduced profitability 129 to 278 \$ ha⁻¹. Substituting urea with PCU did not affect yields in relation to 100% urea but reduced profitability 98 \$ ha⁻¹ and emphasized the risk of enhanced efficiency N use when N loss conditions are not present. In wet soils (above normal May – June rainfall) PL followed by V11 SD increased yield 8.6 to 13.9% relative to IF or 5x5 strategies and V4 SD. Relative to one-pass systems yield gains 4.4 – 16.1% of multi-pass systems in 4 of 6 site years suggest improved synchrony N application with corn N uptake. Delaying the majority of N application until V11 may have reduced the opportunity for N losses in wet years, this practice did not increase the opportunity for yield gains in other years. Since SD N applications are often dependent on seasonal weather patterns growers must leverage risk when deliberating between early and late SD. The ability of N strategies to sufficiently supply N to corn until SD remain critical in order to maintain yield

potential. Canopy sensory data suggest the IF, PPI, and 5x5 strategies can increase plant vigor at V6 and V11 relative to an unfertilized strategy, but reduced N rates required by the IF strategy may reduce the capacity to maintain yield potential when full SD is delayed from V4 to V11. Weather forecast models can vary in their predictive accuracy and reduced yield potential and economic response to V11 SD N suggest less risk is associated with the V4 timing. However, growers missing the V4 application window can utilize the V11 timing as a rescue application to achieve yields ≥ 11.2 Mg ha⁻¹. Growers utilizing a rescue application are more likely to maintain yield and profitability if preceded by increased N rates in the 5x5 strategy.

As models predict warmer temperatures, the use of relative maturity class corn hybrids greater than 98 d during a lengthened growing season may create greater opportunities for late season N applications to have a significant impact on grain yield in northern climates. Future studies that include similar treatments and multiple corn hybrids replicated across Corn Belt regimes may provide evidence to substantiate this idea as well as additional data and tools to address N management in a changing 21st century climate.

Acknowledgements

The authors would like to thank the USDA National Institute of Food and Agriculture, Michigan AgBioResearch, the Michigan State University's College of Agriculture and Natural Resources, and Corn Marketing and Promotion of Michigan for partial funding and support of this research.

APPENDICES

APPENDIX A

CHAPTER 2 TABLES

Table 2.01. Corn planting date, V4, and V11 sidedress N application dates for SCRF (Lansing, MI) and SVRF (Richville, MI), 2014-2016.

	-----SCRF-----			-----SVRF-----		
	Planting	V4	V11	Planting	V4	V11
2014	19 May	09 June	07 July	08 May	04 June	30 June
2015	01 May	02 June	29 June	28 April	28 May	25 June
2016	18 May	06 June	05 July	09 May	03 June	29 June

Table 2.02. Prices received and variable input costs utilized used for profitability analysis, 2014-2016.

Years		2014	2015	2016
	Returns	-----US \$ Mg ⁻¹ -----		
Prices Received [†]	Corn	138.18	142.91	131.88
	Costs	-----US \$ kg ⁻¹ -----		
Fertilizer [‡]	Urea-N	1.20	1.01	0.74
	28% urea ammonium nitrate (UAN)	1.26	1.14	0.81
	10-34-0	5.68	7.19	6.17
	Polymer coated urea (PCU)	1.63	1.54	1.23
	Poultry Litter (PL)	3.78	4.23	4.23
		-----US \$ L ⁻¹ -----		
Nitrogen stabilizer [§]	Urease Inhibitor	19.01	19.01	19.01
		-----US \$ ha ⁻¹ -----		
Application [¶]	5x5 starter applicator	5.24	5.12	4.99
	Urea broadcast application	17.27	14.65	15.59
	Urea incorporation	15.96	17.54	21.62
	UAN injection application (V4)	27.63	31.88	30.32
	UAN high-clearance application (V11)	19.00	25.03	22.76
	Grain hauling (farm to mkt or elv.)	0.32	0.32	0.32

[†]Fall grain prices for each year (USDA – NASS 2014; 2015; 2016).

[‡]Spring urea and UAN prices for each year (USDA – AMS 2014; 2015; 2016); PL, 10-34-0, and PCU prices obtained from local vendors.

[§]Urease inhibitor price obtained from local vendors and applied at label rates.

[¶]Application costs obtained from Michigan State University Extension custom machine and work rate estimates for each year (http://msue.anr.msu.edu/topic/farm_management/firm_publication_archive, Michigan State University, East Lansing, Michigan; verified 23 May 2017).

Table 2.03. Growing season (April – September) and 30-yr mean temperature[†] (°C) and precipitation (mm) data for SCRF (Lansing, MI) and SVRF (Richville, MI), 2014 – 2016.

Site	Year	Apr.	May	Jun.	Jul.	Aug.	Sept.	Total
°C								
SCRF	2014	8.0	14.4	20.0	18.8	20.2	15.4	--
	2015	8.2	16.3	19.2	20.9	20.3	19.1	--
	2016	7.5	14.8	20.3	23.0	22.9	17.4	--
	30-yr	8.7	14.7	20.0	22.1	21.3	16.9	--
SVRF	2014	7.4	14.3	20.2	19.0	19.7	15.5	--
	2015	7.4	15.7	18.5	20.7	20.0	18.6	--
	2016	5.6	14.8	19.7	22.6	22.4	18.2	--
	30-yr [‡]	7.7	14.2	19.7	22.1	20.9	16.8	--
mm								
SCRF	2014	22	83	123	61	86	85	460
	2015	23	109	192	61	123	95	604
	2016	75	52	18	96	163	106	510
	30-yr	73	85	89	83	84	92	505
SVRF	2014	101	78	70	106	99	77	531
	2015	50	73	68	56	100	67	413
	2016	33	40	38	88	131	52	382
	30-yr [‡]	81	84	90	79	82	98	513

[†]Air temperature and precipitation data were collected from Michigan Automated Weather Network (<http://www.agweather.geo.msu.edu/mawn/>) and determined as a mean of the average monthly high and low.

[‡]30-yr mean source for air temperature and precipitation data: NOAA (<https://www.ncdc.noaa.gov/cdo-web/datatools/normals>).

Table 2.04. Impact of N strategy on V3 corn plant density prior to in-season sidedress applications using multiple degree of freedom contrasts[†], across sites, 2014 - 2016.

N strategy	2014	2015	2016
	plants ha ⁻¹		
In-furrow starter (IF) [‡]	79,115 b [#]	81,806 a	80,729 a
Pre-plant, incorporated (PPI) [§]	81,268 a	81,268 a	80,729 a
5x5 starter [¶]	80,729 a	80,729 a	81,806 a
Untreated control	81,268 a	80,191 a	78,038 b
<i>P > F</i>	<0.001	0.612	0.042

[†]Contrasts consisted of 3-treatment means which utilized each strategy.

[‡]Corn received 8 kg N ha⁻¹ with the IF strategy.

[§]Corn received 2.2 Mg ha⁻¹ poultry litter [61 kg N ha⁻¹ first-year mineralizable N] or 157 – 202 kg N ha⁻¹ PPI.

[¶]Corn received 45 kg N ha⁻¹ with the 5x5 strategy.

[#]Treatment means within each column followed by the same letter are not significantly different at $P \leq 0.10$.

Table 2.05. Analysis of variance for V3 stand count, V6 and V11 NDVI, R1 and R4 rel. chlorophyll meter (SPAD), and grain yield in Lansing, and Richville, MI, 2014 - 2016.

Observation	Source of variation ($P > F$)						
	Location (L)	Treatment (T)	Year (Y)	L x T	L x Y	T x Y	Y x L x T
NDVI V6	<0.001	0.026	<0.001	<0.001	<0.001	0.037	0.158
NDVI V11	0.088	0.007	0.001	0.122	0.074	0.870	0.240
Rel. SPAD R1	0.921	<0.001	0.090	<0.001	0.143	<0.001	<0.001
Rel. SPAD R4	0.644	<0.001	0.165	0.479	0.944	0.165	0.313
Grain yield	0.957	<0.001	<0.001	0.786	0.113	0.013	0.002
Net return	0.1723	<0.001	<0.001	0.721	0.165	0.010	0.001

Table 2.06. Corn grain yield[†] as affected by in-furrow (IF), pre-plant incorporated (PPI), and 5x5 N strategies in combination with sidedress timings, at SCRf (Lansing, MI) and SVRF (Richville, MI)[‡] 2014 - 2016.

N strategy	2014				2015				2016			
	SCRf		SVRF		SCRf		SVRF		SCRf	SVRF		
	Mg ha ⁻¹											
IF + V4	14.5	ab [§]	14.0	bc	12.2	bc	12.7	a	12.2	a	13.8	a
IF + V11	13.4	d	13.6	c	12.7	bc	11.4	a	11.8	a	12.4	b
IF + V4/V11	14.4	abc	14.5	ab	13.2	ab	12.4	a	12.0	a	12.9	ab
PPI: urea	13.6	cd	15.0	a	11.9	c	11.2	a	11.8	a	11.2	c
PPI: PCU/urea	13.3	d	14.7	ab	12.0	c	11.5	a	12.4	a	11.1	c
PPI: PL + V11	14.7	a	14.5	ab	13.9	a	12.0	a	12.1	a	13.3	ab
5x5 + V4	13.9	bcd	14.4	abc	12.8	bc	11.9	a	13.1	a	13.0	ab
5x5 + V11	13.4	d	13.7	c	13.1	ab	12.6	a	12.4	a	12.8	b
5x5 + V4/V11	14.2	abc	14.6	ab	12.7	bc	12.3	a	12.7	a	13.3	ab
<i>P</i> > <i>F</i>	0.010		0.049		0.046		0.133		0.349		<0.001	
Untreated [¶]	7.2		6.0		5.9		7.2		8.5		5.8	
Multiple <i>df</i> contrasts												
IF strategy [#]	14.1	a	14.0	b	12.7	a	12.1	a	12.0	b	13.0	a
PPI strategy	13.9	a	14.7	a	12.6	a	11.5	b	12.1	b	11.9	b
5x5 strategy	13.8	a	14.2	b	12.9	a	12.3	a	12.7	a	13.0	a
<i>P</i> > <i>F</i>	0.633		0.019		0.712		0.065		0.066		0.004	
One pass ^{††}	13.5	b	14.9	a	11.9	b	11.3	b	12.1	a	11.2	b
Multi-pass	14.1	a	14.2	b	12.9	a	12.2	a	12.3	a	13.0	a
<i>P</i> > <i>F</i>	0.035		0.006		0.001		0.011		0.487		<0.001	

[†]Grain yield at 155 g kg⁻¹ moisture.

[‡]Total maximum return to N rate used at Lansing and Richville locations was 157 and 202 kg N ha⁻¹, respectively, across years.

[§]Treatment means within each column followed by the same letter are not significantly different at *P* ≤ 0.10.

[¶]Untreated control not included in statistical analysis.

[#]Contrasts consisted of 3-treatment means that utilized each strategy.

^{††}One pass system multiple degree of freedom contrast was mean of urea and PCU and urea treatments. Multi-pass system was a mean of all other treatments combined.

Table 2.07. Corn profitability as affected N strategy at SCRF (Lansing, MI) and SVRF (Richville, MI)[†] 2014 - 2016.

N strategy	2014		2015		2016							
	SCRF	SVRF	SCRF	SVRF	SCRF	SVRF						
	Net Returns [‡] (\$ per ha)											
IF + V4	1663	a [§]	1544	bcd	1412	a	1451	ab	1349	bcd	1521	a
IF + V11	1501	d	1475	d	1470	a	1256	de	1288	d	1329	cde
IF + V4/V11	1618	abc	1570	bc	1514	a	1372	abcd	1300	cd	1369	bcd
PPI: urea	1593	abcd	1719	a	1450	a	1321	bcde	1349	bcd	1244	ef
PPI: PCU/urea	1499	d	1621	b	1395	a	1288	cde	1386	abcd	1167	f
PPI: PL + V11	1558	bcd	1467	d	1498	a	1206	e	1159	e	1276	def
5x5 + V4	1620	abc	1623	b	1550	a	1393	abcd	1505	a	1463	ab
5x5 + V11	1539	cd	1524	cd	1586	a	1484	a	1418	abc	1421	abc
5x5 + V4/V11	1639	ab	1629	ab	1495	a	1416	abc	1435	ab	1464	ab
<i>P</i> > <i>F</i>	0.064		0.001		0.376		0.025		0.002		<0.001	
Untreated [¶]	959		798		813		990		1080		741	
Multiple <i>df</i> contrasts												
IF strategy [#]	1595	a	1532	b	1466	a	1360	a	1313	b	1406	a
PPI strategy	1550	a	1602	a	1448	a	1272	b	1298	b	1229	b
5x5 strategy	1599	a	1592	a	1544	a	1431	a	1453	a	1449	a
<i>P</i> > <i>F</i>	0.338		0.062		0.145		0.005		0.002		<0.001	
One pass [#]	1546	a	1670	a	1422	a	1305	a	1367	a	1205	b
Multi-pass	1591	a	1548	b	1504	a	1368	a	1351	a	1406	a
<i>P</i> > <i>F</i>	0.218		<0.001		0.109		0.152		0.683		<0.001	

[†]Total maximum return to N rate used at Lansing and Richville locations was 157 and 202 kg N ha⁻¹, respectively, in all years.

[‡]Net returns calculated as gross return [yield x corn price] minus total costs [N + N protectant + application cost] ha⁻¹ per respective year.

[§]Treatment means within each column followed by the same letter are not significantly different at *P* ≤ 0.10.

[¶]Untreated control not included in statistical analysis.

[#]Contrasts consisted of 3-treatment means that utilized each strategy.

^{††}One pass system multiple degree of freedom contrast was mean of urea and PCU and urea treatments. Multi-pass system was a mean of all other treatments combined.

Table 2.08. Mean V6 NDVI readings as affected by N strategy across locations at SCRFF (Lansing, MI) and SVRF (Richville, MI) 2014 – 2016.

Treatment combinations	Years			Location						
	2014	2015	2016	SCRFF	SVRF					
	NDVI									
IF + V4	0.4620	cd [‡]	0.3974	ab	0.3296	a	0.4678	a	0.3248	a
IF + V11 [†]	0.4696	cd	0.3965	ab	0.3182	a	0.4647	a	0.3248	a
IF + V4/V11	0.4768	bcd	0.4087	a	0.3187	a	0.4701	a	0.3327	a
PPI: urea	0.5132	a	0.3703	bcd	0.3085	a	0.4358	bc	0.3589	a
PPI: PCU/urea	0.4998	ab	0.3799	bc	0.3233	a	0.4488	ab	0.3531	a
PPI: PL + V11	0.4886	abc	0.3909	ab	0.3176	a	0.4573	ab	0.3408	a
5x5 + V4	0.4700	cd	0.3560	cd	0.3162	a	0.4197	cd	0.3418	a
5x5 + V11	0.4595	d	0.3502	d	0.3072	a	0.4071	d	0.3376	a
5x5 + V4/V11	0.4860	abcd	0.3599	cd	0.3083	a	0.4242	cd	0.3454	a
<i>P</i> > <i>F</i>	0.021		0.002		0.935		<0.001		0.168	
Untreated [§]	0.4263		0.3350		0.3133		0.4174		0.2977	
Multiple <i>df</i> contrasts										
IF starter only [¶]	0.4696	b	0.3965	a	0.3182	a	0.4647	a	0.3248	b
PPI strategy [#]	0.5005	a	0.3804	a	0.3164	a	0.4473	a	0.3509	a
5x5 starter only	0.4595	b	0.3502	b	0.3072	a	0.4071	b	0.3376	ab
Untreated control	0.4263	c	0.3350	b	0.3061	a	0.4174	b	0.2943	c
<i>P</i> > <i>F</i>	<0.001		<0.001		0.856		0.001		<0.001	

[†]Not all treatments at V6 have received total seasonal N application (i.e., V11 N applications).

[‡]Treatment means within each column followed by the same letter are not significantly different at $P \leq 0.10$.

[§]Untreated not included in analysis of pairwise comparisons.

[¶]The IF and 5x5 strategies indicated as ‘starter’ as these contrasts only contained those treatments not receiving SD N to enable comparison with PPI N and the untreated control.

[#]‘strategy’ indicates a mean of all three treatments utilizing the PPI strategy.

Table 2.09. Mean V11 NDVI readings as affected by N strategy across locations and years[†] in Lansing and Richville, MI, 2015 - 2016.

N strategy	NDVI	
IF + V4	0.7309	a [§]
IF + V11 [‡]	0.6823	d
IF + V4/V11	0.7223	ab
PPI: urea	0.7071	bc
PPI: PCU/urea	0.7157	abc
PPI: PL + V11	0.7274	ab
5x5 + V4	0.7132	abc
5x5 + V11	0.7003	cd
5x5 + V4/V11	0.7190	abc
<i>P</i> > <i>F</i>	0.050	
Untreated [¶]	0.6526	
Multiple <i>df</i> contrasts		
IF starter only [#]	0.6823	b
PPI strategy ^{††}	0.7168	a
5x5 starter only	0.7003	ab
Untreated control	0.6526	c
<i>P</i> > <i>F</i>	<0.001	

[†]Data were not collected in 2014.

[‡]Not all treatments at V11 have received total seasonal N application (i.e., V11 N applications).

[§]Treatment means followed by the same letter are not significantly different at $P \leq 0.10$.

[¶]Untreated not included in analysis of pairwise comparisons.

[#]The IF and 5x5 strategies indicated as 'starter' as these contrasts only contained those treatments not receiving SD N to enable comparison with PPI N and the untreated control.

^{††}'strategy' indicates a mean of all three treatments utilizing the PPI strategy.

Table 2.10. Mean R1 relative[†] chlorophyll (CM) meter measurements as affected by N strategy at SCRF (Lansing, MI) and SVRF (Richville, MI), MI[‡], 2014 - 2016.

N strategy	2014		2015		2016							
	SCRF	SVRF	SCRF	SVRF	SCRF	SVRF						
	Relative CM index											
IF + V4	0.99	a [§]	0.97	cd	0.95	bc	1.00	a	1.00	a	1.04	a
IF + V11	0.97	bcd	0.91	e	0.94	c	0.78	e	0.92	d	1.00	bc
IF + V4/V11	0.99	abc	1.02	ab	0.99	a	0.97	ab	1.00	a	1.01	abc
PPI: urea	0.99	abc	1.00	abc	0.98	ab	0.96	bc	0.98	abc	0.99	c
PPI: PCU/urea	0.96	d	1.00	bc	0.95	bc	0.94	c	1.00	a	1.00	bc
PPI: PL + V11	0.99	ab	0.95	d	0.99	a	0.90	d	0.95	c	0.95	d
5x5 + V4	0.97	cd	1.03	a	0.99	a	0.98	ab	1.01	a	1.02	ab
5x5 + V11	0.93	e	0.97	d	0.98	a	0.89	d	0.96	bc	0.99	bc
5x5 + V4/V11	0.97	abcd	1.01	abc	0.99	a	0.97	abc	0.99	ab	1.03	a
<i>P</i> > <i>F</i>	0.052		<0.001		0.019		<0.001		<0.001		<0.001	
Untreated [¶]	0.74		0.71		0.72		0.74		0.84		0.73	
	Multiple <i>df</i> contrasts											
IF strategy [#]	0.98	a	0.97	b	0.96	a	0.92	b	0.97	a	1.02	a
PPI strategy	0.98	a	0.98	ab	0.97	a	0.93	ab	0.98	a	0.98	b
5x5 strategy	0.96	b	1.00	a	0.99	a	0.95	a	0.98	a	1.01	a
<i>P</i> > <i>F</i>	0.013		0.074		0.119		0.078		0.523		<0.001	
One pass ^{††}	0.97	a	1.00	a	0.96	a	0.95	a	0.99	a	0.99	a
Multi-pass	0.97	a	0.98	a	0.97	a	0.93	b	0.97	a	1.00	a
<i>P</i> > <i>F</i>	0.818		0.124		0.410		0.074		0.129		0.135	

[†]Readings normalized to non-limiting N control (280 kg N ha⁻¹).

[‡]Total maximum return to N rate used at Lansing and Richville locations was 157 and 202 kg N ha⁻¹, respectively, in all years.

[§]Treatment means within each column followed by the same letter are not significantly different at *P* ≤ 0.10.

[¶]Untreated control not included in statistical analysis.

[#]Contrasts consisted of 3-treatment means that utilized each strategy.

^{††}One pass system multiple degree of freedom contrast was mean of urea and PCU and urea treatments. Multi-pass system was a mean of all other treatments combined.

Table 2.11. Mean R4 relative[†] chlorophyll (CM) meter measurements as affected by N strategy across locations[‡] and years[§], 2015-2016.

Treatment combinations	Rel. CM index	
IF + V4	0.99	ab [¶]
IF + V11	0.93	cd
IF + V4/V11	0.99	ab
PPI: urea	0.93	d
PPI: PCU/urea	0.95	cd
PPI: PL + V11	0.96	bc
5x5 + V4	1.02	a
5x5 + V11	0.94	cd
5x5 + V4/V11	0.99	ab
<i>P</i> > <i>F</i>	<0.001	
Untreated [#]	0.57	
Multiple <i>df</i> contrasts		
IF strategy ^{††}	0.97	a
PPI strategy	0.94	b
5x5 strategy	0.98	a
<i>P</i> > <i>F</i>	0.01	
One pass ^{‡‡}	0.94	b
Mutli-pass	0.98	a
<i>P</i> > <i>F</i>	0.001	

[†]Readings normalized to non-limiting N control (280 kg N ha⁻¹).

[‡]Total maximum return to N rate used at Lansing and Richville locations was 157 and 202 kg N ha⁻¹, respectively, in all years.

[§]Data not collected in 2014.

[¶]Treatment means followed by the same letter are not significantly different at $P \leq 0.10$.

[#]Untreated control not included in statistical analysis.

^{††}Contrasts consisted of 3-treatment means that utilized each strategy.

^{‡‡}One pass system multiple degree of freedom contrast was mean of urea and PCU and urea treatments. Multi-pass system was a mean of all other treatments combined.

APPENDIX B

CHAPTER 2 DATA COLLECTED AND NOT USED FOR PUBLICATION

Table B.01. Total vegetative N uptake[†] at physiological maturity as affected by in-furrow (IF), pre-plant incorporated (PPI), and 5x5 N strategies in combination with sidedress timings, at SCRf (Lansing, MI) and SVRF (Richville, MI)[‡] 2014 - 2016.

N strategy	2014				2015				2016			
	SCRf		SVRF		SCRf		SVRF		SCRf		SVRF	
	kg ha ⁻¹											
IF + V4	61.5	ab [§]	38.1	bc	33.3	a	46.3	a	42.5	a	37.7	a
IF + V11	53.7	bc	40.4	bc	51.7	a	26.7	e	39.9	a	39.6	a
IF + V4/V11	66.4	a	54.7	a	48.0	a	38.2	abc	44.2	a	33.7	a
PPI: urea	52.9	bc	46.3	ab	42.5	a	36.4	bcd	42.8	a	40.6	a
PPI: PCU/urea	48.3	c	46.2	b	42.3	a	33.4	cde	49.2	a	39.4	a
PPI: PL + V11	65.7	a	39.1	bc	45.7	a	32.9	cde	38.9	a	41.4	a
5x5 + V4	48.8	c	46.9	ab	40.4	a	43.1	ab	45.9	a	36.3	a
5x5 + V11	50.2	c	35.6	c	42.6	a	28.3	de	39.7	a	34.1	a
5x5 + V4/V11	51.8	bc	41.8	bc	36.6	a	35.6	bcd	41.9	a	38.3	a
<i>P</i> > <i>F</i>	0.0172		0.0181		0.1718		0.0051		0.8244		0.8105	
Untreated [¶]	22.6		19.2		17.5		15.1		22.2		17.1	
Multiple <i>df</i> Contrasts[#]												
IF strategy	60.6	a	44.3	a	44.3	a	37.1	a	42.2	a	37.0	a
PPI strategy	55.7	ab	43.9	a	43.5	a	34.2	a	43.6	a	40.5	a
5x5 strategy	50.3	b	41.1	a	39.9	a	35.6	a	42.5	a	36.1	a
<i>P</i> > <i>F</i>	0.0308		0.5201		0.5380		0.6978		0.8973		0.2415	
PPI: urea	53.1	a	46.3	a	42.5	a	36.4	a	42.8	a	40.6	a
PPI: PCU/urea	48.3	a	46.2	a	42.3	a	33.4	a	49.2	a	39.4	a
Split-N	56.9	a	42.2	a	42.6	a	35.9	a	41.9	a	37.2	a
<i>P</i> > <i>F</i>	0.1758		0.3533		0.9978		0.8351		0.2175		0.5536	

[†]Determined as total vegetative biomass multiplied by percent total N.

[‡]Total maximum return to N rate used at Lansing and Richville locations was 157 and 202 kg N ha⁻¹, respectively, across years.

[§]Treatment means within each column followed by the same letter are not significantly different at *P* ≤ 0.10.

[¶]Untreated control not included in statistical analysis.

[#]Contrasts consisted of 3-treatment means that utilized each strategy.

Table B.02. Soil residual nitrate[†] at corn harvest as affected by in-furrow (IF), pre-plant incorporated (PPI), and 5x5 N strategies in combination with sidedress timings, at SCRf (Lansing, MI) and SVRF (Richville, MI)[‡] 2014 - 2016.

N strategy	2014				2015				2016			
	SCRf		SVRF		SCRf		SVRF		SCRf		SVRF	
	kg N ha ⁻¹											
IF + V4	13.6	a [§]	14.0	cd	11.3	a	20.3	a	19.8	a	10.7	a
IF + V11	26.6	a	41.2	a	13.5	a	11.1	bcde	11.8	a	9.2	a
IF + V4/V11	15.1	a	27.9	a	13.8	a	19.1	ab	12.2	a	10.1	a
PPI: urea	16.7	a	21.0	bc	12.8	a	8.6	cde	9.6	a	10.3	a
PPI: PCU/urea	15.1	a	27.0	b	13.6	a	8.0	e	13.5	a	12.5	a
PPI: PL + V11	25.5	a	18.6	bc	11.5	a	14.3	bc	10.5	a	8.6	a
5x5 + V4	13.8	a	11.5	d	12.4	a	18.7	a	12.1	a	7.8	a
5x5 + V11	15.2	a	20.9	bc	11.7	a	12.1	bcd	8.5	a	8.8	a
5x5 + V4/V11	16.7	a	16.2	bcd	13.6	a	8.1	de	13.2	a	8.8	a
<i>P</i> > <i>F</i>	0.2792		0.0003		0.9905		0.0002		0.1860		0.6226	
Untreated [¶]	13.6		9.1		9.2		7.4		8.3		8.6	
Multiple <i>df</i> Contrasts[#]												
IF strategy	18.4	a	27.7	a	12.9	a	16.8	a	14.6	a	10.0	a
PPI strategy	19.1	a	22.9	ab	12.6	a	10.3	b	11.2	a	10.5	a
5x5 strategy	15.2	a	16.2	b	12.6	a	13.0	ab	11.3	a	8.4	a
<i>P</i> > <i>F</i>	0.6019		0.0757		0.9629		0.0395		0.1080		0.1363	
PPI: urea	16.7	a	22.4	a	12.8	a	8.6	b	9.6	a	10.3	ab
PPI: PCU/urea	15.1	a	27.8	a	13.6	a	8.0	b	13.5	a	12.5	a
Split-N	18.1	a	21.5	a	12.5	a	14.8	a	12.6	a	9.1	b
<i>P</i> > <i>F</i>	0.8909		0.6860		0.8391		0.0243		0.2987		0.0597	

[†]Determined at the 0 – 30 centimeter depth.

[‡]Total maximum return to N rate used at Lansing and Richville locations was 157 and 202 kg N ha⁻¹, respectively, across years.

[§]Treatment means within each column followed by the same letter are not significantly different at $P \leq 0.10$.

[¶]Untreated control not included in statistical analysis.

[#]Contrasts consisted of 3-treatment means that utilized each strategy.

Table B.03. Corn stalk nitrate[†] at corn physiological maturity as affected by in-furrow (IF), pre-plant incorporated (PPI), and 5x5 N strategies in combination with sidedress timings, at SCRf (Lansing, MI) and SVRF (Richville, MI)[‡] 2014 - 2016.

N strategy	2014				2015				2016			
	SCRf		SVRF		SCRf		SVRF		SCRf		SVRF	
	mg NO ₃ -N kg ⁻¹											
IF + V4	447	a [§]	359	abc	445	a	1579	a	748	a	84	c
IF + V11	268	a	195	bc	138	a	91	c	121	bc	326	b
IF + V4/V11	467	a	226	abc	426	a	592	b	211	abc	53	cd
PPI: urea	99	a	220	abc	432	a	636	b	347	ab	934	a
PPI: PCU/urea	217	a	504	a	482	a	867	b	544	a	1238	a
PPI: PL + V11	516	a	391	ab	529	a	82	c	49	c	62	c
5x5 + V4	179	a	213	c	233	a	2227	a	625	a	35	cd
5x5 + V11	240	a	156	c	366	a	325	b	145	bc	40	cd
5x5 + V4/V11	548	a	357	ab	655	a	744	b	267	abc	7	d
<i>Pr</i> > <i>F</i>	0.4327		0.0886		0.8351		<.0001		0.0439		<.0001	
Untreated [¶]	1.3		0.0		1.9		3.4		1.1		0.3	
	Multiple <i>df</i> Contrasts[#]											
IF strategy	393	a	264	a	336	a	754	b	359	a	155	b
PPI strategy	281	a	372	a	481	a	528	b	313	a	744	a
5x5 strategy	322	a	248	a	418	a	1099	a	351	a	27	c
<i>Pr</i> > <i>F</i>	0.3364		0.1936		0.6187		0.0046		0.6493		<.0001	
PPI: urea	109	b	220	a	432	a	636	a	347	a	934	a
PPI: PCU/urea	217	ab	504	a	482	a	867	a	544	a	1238	a
Split-N	380	a	275	a	399	a	806	a	311	a	87	b
<i>Pr</i> > <i>F</i>	0.0631		0.2207		0.9262		0.7033		0.2046		<.0001	

[†]Determined using 20 centimeter stalk segments collected 15 centimeters above the soil surface.

[‡]Total maximum return to N rate used at Lansing and Richville locations was 157 and 202 kg N ha⁻¹, respectively, across years.

[§]Treatment means within each column followed by the same letter are not significantly different at $P \leq 0.10$.

[¶]Untreated control not included in statistical analysis.

[#]Contrasts consisted of 3-treatment means that utilized each strategy.

LITERATURE CITED

LITERATURE CITED

- Bender, R.R., J.W. Haegele, M.L. Ruffo, and F.E. Below. 2013. Nutrient uptake, partitioning, and remobilization in modern, transgenic, insect-protected maize hybrids. *Agron. J.* 105:161-170.
- Bermudez, M., and A.P. Mallarino. 2002. Yield and early growth responses to starter fertilizer in no-till corn assessed with precision agriculture technologies. *Agron. J.* 94:1024-1033.
- Bermudez, M., and A.P. Mallarino. 2004. Corn response to starter fertilizer and tillage across and within fields having no-till management histories. *Agron. J.* 96:776-785.
- Binder, D.L., D.H. Sander, and D.T. Walters. 2000. Maize response to time of nitrogen application as affected by level of nitrogen deficiency. *Agron. J.* 92:1228-1236.
- Bundy, L.G., and T.W. Andraski. 1999. Site-specific factors affecting corn response to starter fertilizer. *J. Prod. Agric.* 12:664-670.
- Cromley, S.M., W.J. Wiebold, and P.C. Scharf. 2003. Tillage effects on corn response to starter fertilizer. Online. *Crop Management* doi:10.1094/CM-2003-0623-01-RS.
- Cromley, S.M., W.J. Wiebold, P.C. Scharf, and S.P. Conley. 2006. Hybrid and planting date effects on corn response to starter fertilizer. Online. *Crop Management* doi:10.1094/CM-2006-0906-01-RS.
- Combs, S.M., and M.V. Nathan. 2015. Soil organic matter. In: M.V. Nathan and R. Gelderman, editors, *Recommended chemical soil test procedures for the North Central Region*. North Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 12.1-12.6.
- Dove, A., and S.C. Chapra. 2015. Long-term trends of nutrients and trophic response variables for the Great Lakes. *Limnol. Oceanogr.* 60:696-721.
- Elwadie, M.E., F.J. Pierce, and J. Qi. 2005. Remote sensing of canopy dynamics and biophysical variables estimation of corn in Michigan. *Agron. J.* 97:99-105.
- Evans, L.T., and R.A. Fischer. 1999. Yield potential: Its definition, measurement, and significance. 39:1544-1551.
- Frank, K., D. Beegle, and J. Denning. 2015. Phosphorus. In: M.V. Nathan and R. Gelderman, editors, *Recommended chemical soil test procedures for the North Central Region*. North

- Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 6.1-6.6.
- Franzen, D.W. 2017. Nitrogen extenders and additives. NDSU Extension publication SF 1581. North Dakota State University Extension Service, Fargo, ND.
www.ag.ndsu.edu/publications/crops/nitrogen-extenders-and-additives-for-field-crops/sf1581.pdf. Accessed Apr. 04, 2017.
- Gagnon, B., Z. Ziadi, and C. Grant. 2012. Urea fertilizer forms affect corn grain yield and nitrogen use efficiency. *Can. J. Soil Sci.* 92:341-351.
- Hatfield, J.L., and T.B. Parkin. 2014. Enhanced efficiency fertilizers: Effect on agronomic performance of corn in Iowa. *Agron. J.* 106:771-780.
- Havlin, J.L., S.L. Tisdale, W.L. Nelson, and J.D. Beaton. 2014. *Soil Fertility and Fertilizers. An Introduction to Nutrient Management*. 8th ed. Pearson Education Inc., Upper Saddle River, NJ.
- Hayhoe, K., C. Wake, T. Huntington, L. Luo, M. Schwartz, J. Sheffield, E. Wood, B. Anderson, J. Bradbury, A. Degaetano, T. Troy, and D. Wolfe. 2007. Past and future changes in climate and hydrological indicators in the U.S. *Clim. Dynam.* 28:381-407.
- Jokela, W.E., and G.W. Randall. 1997. Fate of fertilizer nitrogen as affected by time and rate of application on corn. *Soil Sci. Soc. Am. J.* 61:1695-1703.
- Kaiser, D.E., J.A. Coulter, and J.A. Vetsch. 2016. Corn hybrid response to in-furrow starter fertilizer as affected by planting date. *Agron. J.* 108:2493-2501.
- Karl, T.R., J.M. Melillo, and T.C. Peterson (ed.). 2009. *Global climate change impacts in the United States*. Cambridge Univ. Press, New York.
- Laboski, C.A.M. 2008. Understanding salt index of fertilizers. In: *Proceedings of the Wisconsin Fertilizer, Aglime, and Pest Management Conference*, Madison, WI. 15-17 Jan. 2008. Vol. 47. Univ. of Wisconsin Coop. Ext., Madison.
- Lamond, R.E. and W.B. Gordon. 2001. NPK starters best N-only starters. *Fluid J.* Fall 2001.
- Lauer, J.G., P.R. Carter, T.M. Wood, G. Diezel, D.W. Wiersma, R.E. Rand, and M.J. Mlynarek. 1999. Corn hybrid response to planting date in the Northern Corn Belt. *Agron. J.* 91:834-839.
- Liu, K., and P. Wiatrak. 2011. Corn production and plant characteristics response to N fertilization management in dry-land conventional tillage system. *Int. J. Pl. Prod.* 5:405-416.

- Maharjan, B., C.J. Rosen, J.A. Lamb, and R.T. Venterea. 2016. Corn response to nitrogen management under fully-irrigated vs. water-stressed conditions. *Agron. J.* 108:1-10.
- Niehues, B.J., R.E. Lamond, C.B. Godsey, and C.J. Olsen. 2004. Starter nitrogen fertilizer management for continuous no-till corn production. *Agron. J.* 96:1412-1418.
- Norton, R. 2014. Combating climate change through improved agronomic practices and input-use efficiency. *J. Crop Improv.* 28:575-618.
- Peters, J.B., M.V. Nathan, and C.A.M. Laboski. 2015. pH and lime requirement. In: M.V. Nathan and R. Gelderman, editors, *Recommended chemical soil test procedures for the North Central Region*. North Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 4.1-4.7.
- Randall, G.W., J.A. Vetsch, and J.R. Huffman. 2003. Corn production on a subsurface-drained Mollisol as affected by time of nitrogen application and nitrapyrin. *Agron. J.* 95:1213-1219.
- Raun, W.R., D.H. Sander, and R.A. Olson. 1986. Emergence of corn as affected by source and rate of solution fertilizers applied with the seed. *J. Fert. Issues* 3:18-24.
- Rawluk, C.D.L, C.A. Grant, and G.J. Racz. 2001. Ammonia volatilization from soils fertilized with urea and varying rates of urease inhibitor NBPT. *Can. J. Soil Sci.* 81:239-246.
- Rehm, G.W., and J.A. Lamb. 2009. Corn response to fluid fertilizers placed near the seed at planting. *Soil Sci. Soc. Am. J.* 73(4):1427-1434.
- Ritchie, K.B., R.G. Hoelt, E.D. Nafziger, L.C. Gonzini, and J.J. Warren. 1996. Starters perform more consistently than seed-placed fertilizers. *Fluid J.* Winter 1996:1-3.
- Ritchie, S.W., J.J. Hanway, and G.O. Benson. 1997. How a corn plant develops. Spec. Rep. 48. Iowa State Univ. of Sci. and Technol., Coop. Ext. Serv., Ames.
- SAS Institute. 2011. *The SAS system for windows*. Version 9.4. SAS Inst., Cary, NC.
- Sawyer, J., E. Nafziger, G. Randall, L. Bundy, G. Rehm, and B. Joern. 2006. Concepts and rationale for regional nitrogen rate guidelines for corn. Publ. PM2015, Iowa State University Extension, Ames, IA.
- Scavia, D., J.D. Allan, K.K. Arend, S. Bartell, D. Beletsky, N.S. Bosch, S.B. Brandt, R.D. Briland, I. Daloğlu, J. V. DePinto, D.M. Dolan, M.A. Evans, T.M. Farmer, D. Goto, H. Han, T.O. Höök, R. Knight, S.A. Ludsins, D. Mason, A.M. Michalak, R.P. Richards, J.J. Roberts, D.K. Rucinski, E. Rutherford, D.J. Schwab, T.M. Sesterhenn, H. Zhang, and Y. Zhou. 2014.

- Assessing and addressing the re-eutrophication of Lake Erie: Central basin hypoxia. *J. Great Lakes Res.* 40, 226–246.
- Scharf, P.C. 1999. On-farm starter fertilizer response to no-till corn. *J. Prod. Agric.* 12:692-695.
- Scharf, P.C., W.J. Wiebold, and J.A. Lory. 2002. Corn yield response to nitrogen fertilizer timing and deficiency level. *Agron. J.* 94:435-441.
- Schwartz, M.D., R. Ahas, and A. Aasa. 2006. Onset of spring starting earlier across the Northern Hemisphere. *Glob. Change Biol.* 12:343–351.
- Shapiro, C.A., D.D. Francis, R.B. Ferguson, G.W. Hergert, T.M. Shaver, and C.S. Wortmann. 2013. Using a chlorophyll meter to improve N management. Available at extensionpublications.unl.edu/assets/pdf/g1632.pdf [cited 03 Mar. 2016; verified 24 Jul. 2017]. Univ. of Nebraska, Lincoln.
- Snyder, C.S., T.W. Bruulsema, T.L. Jensen, and P.E. Fixen. 2009. Review of greenhouse gas emissions from crop production systems and fertilizer management effects. *Agric. Ecosyst. Environ.* 133:247-266.
- Steinke, K. 2013. Look beyond product rates when considering pop-up fertilizers. Michigan State University Ext. Online. Available at http://msue.anr.msu.edu/news/look_beyond_product_rates_when_considering_pop_up_fertilizers (posted 25 Apr. 2013; verified 30 June 2017).
- Tebaldi, C., K. Hayhoe, J.M. Arblaster, and G.E. Meehl. 2006. Going to the extremes: An intercomparison of model-simulated historical and future changes in extreme events. *Clim. Change* 79:185–211.
- Tremblay, N., Y.M. Bouroubi, C. Bélec, R.W. Mullen, N.R. Kitchen, W.E. Thomason, S. Ebelhar, D.B. Mengel, W.R. Raun, D.D. Francis, E.D. Vories, and I. Ortiz-Monasterio. 2012. Corn response to nitrogen is influenced by soil texture and weather. *Agron. J.* 104:1658-1671.
- Tucker, C.J. 1979. Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sensing of Environ.* 8:127-150.
- Varvel, G.E., J.S. Schepers, and D.D. Francis. 1997. Ability of in-season correction of nitrogen deficiency in corn using chlorophyll meters. *Soil Sci. Soc. Am. J.* 61:1233-1239.
- Venterea, R.T., and J.A. Coulter. 2015. Split application of urea does not decrease and may increase nitrous oxide emissions in rainfed corn. *Agron. J.* 107:337-348.

- Vetsch, J.A., and G.W. Randall. 2000. Enhancing no-tillage systems for corn with starter fertilizers, row cleaners, and nitrogen placement methods. *Agron. J.* 92:309-315.
- Vetsch, J.A., and G.W. Randall. 2004. Corn production as affected by nitrogen application timing and tillage. *Agron. J.* 96:502-509.
- Warncke, D., J. Dahl, and L. Jacobs. 2009. Nutrient recommendations for field crops in Michigan. Bull. E2904, Michigan State University Extension, East Lansing, MI.
- Warncke, D., and J.R. Brown. 2015. Potassium and other basic cations. In: M.V. Nathan and R. Gelderman, editors, Recommended chemical soil test procedures for the North Central Region. North Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 7.1-7.3.
- Wolkowski, R.P. 2000. Row-placed fertilizer for maize grown with an in-row crop residue management system in southern Wisconsin. *Soil & Till. Res.* 54:55-62.
- Yu, C.L., D. Hui, Q. Deng, J. Wang, K.C. Reddy, and S. Dennis. 2016. Responses of corn physiology and yield to six agricultural practices over three years in middle Tennessee. *Nature* 6:27504.
- Zhao, D., K.R. Reddy, V.G. Kakani, J.J. Read, and G.A. Carter. 2003. Corn (*Zea mays* L.) growth, leaf pigment concentration, photosynthesis and leaf hyperspectral reflectance properties as affected by nitrogen supply. *Plant and Soil* 257:205-217.
- Zhou, B., Y. Yue, X. Sun, X. Wang, Z. Wang, W. Ma, and M. Zhao. 2016. Maize grain yield and dry matter production responses to variations in weather conditions. *Agron. J.* 108:196-204.

CHAPTER 3

NITROGEN CYCLING, CORN GROWTH AND GRAIN YIELD RESPONSE TO NON- LEGUMINOUS COVER CROPS AND FERTILIZER

Abstract

Cover crops (CC) seeded prior to a corn (*Zea mays* L.) cash crop may influence nitrogen (N) availability but little validation has occurred when combined with corn sidedress (SD) N applications. Field studies were conducted in 2015 and 2016 to evaluate the effects of a Daikon radish [*Raphanus sativus* (L.) var. The Buster], Forage oat [*Avena sativa* (L.) var. Magnum], or no CC established following wheat (*Triticum aestivum* L.) on soil NO₃-N levels, corn growth, N uptake and grain yield. Nitrogen management strategies included 179 kg N ha⁻¹ applied as urea pre-plant incorporated (PPI), poultry litter (PL) PPI (61 kg N ha⁻¹) plus SD N V11, starter N (52 kg N ha⁻¹) applied 5 cm beside and 5 cm below the furrow (5x5) followed by SD at V4, V11, or V4 plus V11, and a zero N control. Cover crops reduced autumn soil NO₃-N levels 78 to 84% compared with the no cover control but did not increase N availability in corn. When 5x5 starter was followed by full SD at V11 radish and oats reduced yield 3.2 – 3.9% and profitability 11.8 – 13.2% from the no CC control and indicated CCs reduced the effectiveness of 5x5 starter to maintain yield potential until SD time. In the zero N strategy, CCs reduced grain yield 11.8 – 14.2% and increased yield response to N 63 – 79%, suggesting CCs reduced N availability. In years where excessive rainfall increases N loss opportunities CCs may exacerbate soil NO₃-N reduction and increase grain yield reductions. Regardless of CC, growers may reduce risk of yield and profitability loss when full SD at V4 follows 5x5 starter N.

Introduction

Degradation of Great Lakes water quality may require further improvement upon corn nitrogen (N) management strategies to simultaneously increase nitrogen use efficiency (NUE) and reduce opportunities for N loss (Dove and Chapra, 2015; USEPA, 2017). Michigan, located in the northern corn belt and the central basin of the Great Lakes, is characterized by a temperate, humid growing season. Winter wheat (*Triticum aestivum* L.) often precedes corn in Michigan crop rotations and is harvested in July leaving approx. a 90 d fallow period prior to autumn soil freezing (-2.2°C) (National Oceanic and Atmospheric Administration [NOAA], <http://www.weather.gov/dtx/firstFallfreeze>; verified 02 Aug. 2017).

Climate change has prolonged the period when autumn soil temperatures remain above 50 degrees F which may extend organic matter mineralization and increase soil NO₃-N availability (Schwartz et al., 2006; USEPA, 2016; Schultze et al., 2016). In fallow soils, residual soil NO₃-N combined with residue mineralization and no crop N uptake can increase the potential for N loss (Weinert et al., 2002; Derby et al., 2009). To assimilate available N (i.e., post-harvest autumn residual N, inorganic N from soil organic matter (SOM) mineralization) and ameliorate N loss in humid climates, cover crops (CC) are sometimes established in the non-growing season (Shipley et al., 1992; Macdonald et al., 2005; Tonitto et al., 2006; McSwiney et al., 2010). Cover crop biomass production and ensuing N assimilation may improve N cycling and use efficiency (NUE) in maize, maintain and support corn yield, and improve profitability (Decker et al., 1994; McSwiney et al., 2010; O'Reilly et al., 2012). However, corn yields following non-leguminous CCs may be affected by the rate of residue decay and resulting N availability (Dapaah and Vyn, 1998; Vyn et al., 2000; O'Reilly et al., 2012; Gieske et al., 2016a). Corn yield gains and profitability may depend upon the synchrony of CC N availability and N fertilizer timing.

Radish (*Raphanus sativus* L.) is a non-leguminous Brassicaceae CC utilized as a nutrient scavenger in the non-growing season to sequester nutrients including available soil NO₃-N (Mutch and Thelen, 2003; Warncke et al., 2009). Radishes are a large-taproot annual that grow quickly in late summer and autumn and reduce soil compaction and suppress winter annual weeds (Williams and Weil, 2004; Clark, 2007; Lawley et al., 2011). Radish winterkills when air temperatures drop below -4 °C eliminating the need for chemical termination (Thomas et al., 2017). Autumn radish biomass production in Michigan has ranged from 2.9 to 6.1 Mg ha⁻¹ (herbage and roots) after 2 to 3 mo. of growth while N uptake was 10 to 164 kg ha⁻¹ and reduced soil inorganic N 24 to 42% compared to no CC (Hill et al., 2016). However, dry autumn soils may limit plant growth and reduce N uptake (Gieske et al., 2016b). Corn grain yields following an autumn radish CC have varied showing both no effect (Vyn et al., 2000; Gieske et al., 2016a; 2016b) and increased yield (Dapaah and Vyn, 1998; Vyn et al., 1999; O'Reilly et al., 2012) compared to no CC. Variable corn yield response may be due to effects of biomass production, chemical composition, and residue decomposition on subsequent N availability. Cover crop residue decomposition and N mineralization and immobilization are dependent on the carbon (C) to nitrogen ratio (C:N) and lignin concentrations (Wagger et al., 1998; Jahanzad et al., 2016). Decreased C:N associated with a radish cover (14 to 31:1) relative to a grassy cover (15 to 38:1) (e.g., oats, triticale, or ryegrass) may help explain the 43 and 24% greater soil NO₃-N concentrations following a radish CC at mid-May corn planting and mid-June N sidedress application in Ontario, respectively (Vyn et al., 1999; Andraski and Bundy, 2005). However when compared to a no CC control, radishes may increase the risk of N loss (i.e., leaching or denitrification) during freeze-thaw cycles in the non cash-crop growing season (Dean and Weil, 2009; Thomas et al., 2017) and may not increase N availability to a subsequent corn crop (Vyn et

al., 2000; Gieske et al., 2016a). Adjusting N fertilizer strategies may be needed to synchronize radish N availability with corn N uptake and improve grain yield.

Oat (*Avena sativa* L.) is a non-leguminous CC also used to sequester residual soil N and prevent wind and water erosion (Warncke et al., 2009). Similar to radish, oats are an annual CC subject to winterkill (Johnson et al., 1998; Snapp et al., 2005). Unlike radish, oat has a dense, fibrous root system characteristic of cereal CCs (Thorup-Kristensen, 2001). Cereal CC residues often contain more lignin and cellulose that may slow decomposition rates and extend N release when compared to radish (Jahanzad et al., 2016). Slower decay of cereal residue may affect synchrony N availability with corn uptake (Wagger, 1989). The C:N ratio of oat may exceed 30:1 which can immobilize soil N and reduce availability for uptake (Andraski and Bundy, 2005; Dean and Weil, 2009). Corn yield following oat CCs have been generally decreased or not affected in non-irrigated studies (Johnson et al., 1998; Vyn et al., 2000; O'Reilly et al., 2012). However, Andraski and Bundy (2005) suggested corn yield gains due to oat CCs in an irrigated study were an effect of rotation and not N contributions as evidenced by similar May soil NO₃-N contents. Vyn et al. (2000) suggested oat CCs may reduce corn N availability as indicated by greater corn yield response with added fertilizer N. Reduced corn yield following an oat CC corresponded to reduced soil NO₃-N concentrations at corn planting (36%) and at mid-June sidedress (20%) (Vyn et al., 2000). Cereal CCs have also increased gaseous N₂O emissions 76% relative to no CC during the non-growing season and may help explain reduced N availability to the subsequent corn crop (Thomas et al., 2017).

Michigan corn growers utilize multiple strategies to optimize corn N management. Some growers prefer one-pass application systems where 100% of the N inputs are applied prior to or within 1 – 3 d after spring planting but this can reduce corn yield fine-textured soils (Franzen, 2017). Volatile spring weather has prompted grower adoption of multi-pass N application systems for improved N recovery (Vetsch and Randall, 2004; Warncke et al., 2009). Multi-pass N application systems may involve starter N (up to 67 kg N ha⁻¹) applied in a sub-surface band placed 5 cm below and 5 cm laterally (5x5) relative to the seed furrow followed by sidedress (SD) N at V4 corn. Recent studies identified maximum corn N uptake between V10 and V14 and suggested delayed in-season N SD applications may improve synchrony between N availability and uptake while reducing losses from early applied N (Binder et al., 2000; Scharf et al., 2002; Bender et al., 2013). However V6 N deficiency has reduced corn yield potential and highlighted the importance of N management strategies that maintain yield potential until SD time (Binder et al., 2000). Although radish and oat CCs can tighten the N cycle by sequestering autumn soil N, additional research is needed to investigate the impact of CCs on corn response to N management. The objective of this study was to evaluate the effect of N placement and timing on corn growth and grain yield in response to a preceding daikon radish or forage oat cover crop.

Materials and Methods

Site characteristics and cultural practices

A two-year field study was conducted (2014 – 2016) at the Michigan State University South Campus Research Farm in Lansing, MI on a Capac loam (fine-loamy, mixed, active, mesic Aquic Glossudalf). Soil samples (20-cm depth) were collected prior to August CC establishment and May corn planting, air-dried, and ground to pass through a 2-mm sieve. Soil characteristics

5.9-6.2 pH (1:1 soil/water) (Peters et al., 2015), 27-51 mg kg⁻¹ P (Bray-P1) (Frank et al., 2015), 105-163 mg kg⁻¹ K (ammonium acetate method) (Warncke and Brown, 2015), and 2.4-3.4 g kg⁻¹ soil organic matter (loss-on-ignition) (Combs and Nathan, 2015). Fields were chisel plowed after July wheat [*Triticum aestivum* L.] harvest and disk harrowed (10-cm depth) prior to CC establishment. Calcitic lime was spring-applied at 2.2 Mg ha⁻¹ prior to corn planting to a target pH of 6.5. Prior to corn planting fields received disk tillage (10-cm depth) followed by a soil finisher (10-cm depth). Broadcast P and K fertilizer were pre-plant incorporated (PPI) (10-cm depth) as triple super phosphate (0-45-0 N-P-K) and muriate of potash (0-0-62) based on soil test. Corn weed control consisted of acetochlor [2-chloro-N-ethoxymethyl-N-(2-ethyl-6-methylphenyl) acetamide] and glyphosate [N-(phosphonomethyl) glycine] followed by a second application of glyphosate 17-24 d later. Environmental data were collected during the growing season using the Michigan Automated Weather Network (<http://www.agweather.geo.msu.edu/mawn/>, Michigan State University, East Lansing, Michigan; verified 24 Jul. 2017) from an on-site weather station.

Experimental design and treatment application

Eighteen treatments were arranged in a split-plot, randomized complete block design with four replications. The main plot factor was CC while the subplot factor was N strategy. Three CC treatments included a no cover control, 'The Buster' daikon radish, or 'Magnum' forage oat (Weaver Seed of Oregon, Crabtree, OR) and measured 55 m by 12 m in length. The radish cover was drill-planted at 11.2 kg ha⁻¹ and forage oat cover at 28.0 kg ha⁻¹ in August preceding the ensuing corn crop using a Gandy orbit-Air Seeder coupled with John Deere double disk openers in 19-cm rows (Table 1). The no CC treatment received a single, glyphosate application in the

autumn to remain free of vegetative ground cover. To ensure winterkill, CCs were terminated with glyphosate in November (79-83 days of growth).

Subplots consisted of six N placement by N timing strategies applied to corn plots measuring 4.6 m by 12.2 m in length. Nitrogen strategies included a zero N control, pre-plant incorporated (PPI) urea (46-0-0), dried poultry litter (4-3-2) applied PPI at 2.2 Mg ha⁻¹ plus sidedress (SD) N at V11 corn, and starter N subsurface banded 5 cm beside and 5 cm below the seed furrow (5x5) (45 kg N ha⁻¹ urea ammonium nitrate [UAN; 28-0-0]) followed by SD N at V4, V11, or split (50/50) V4 and V11. Corn V4 N was UAN coulter injected 5 cm deep and 38 cm to the side of each row while V11 SD was UAN mixed with a urease inhibitor (CO(NH₂)₂ + n-(n-butyl) thiophosphoric triamide] (Koch Agronomic Services, LLC, Wichita, KS) to prevent N volatilization and banded 10 – 15 cm to the side of each row. Treatment total N rates were equalized to the maximum return to nitrogen rate (179 kg N ha⁻¹) (Steinke, 2015). Corn seeding and starter N application utilized a Monosem planter (Monosem Inc., Kansas City, Kansas) equipped with Yetter floating planter-unit mounted row cleaners (Yetter Manufacturing, Colchester, IL) and liquid fertilizer applicators. Corn was seeded in 0.76 m rows at 84,510 seeds ha⁻¹ using Dekalb DKC48-12 RIB (98 d relative maturity) (Monsanto Co., St. Louis, MO) (Table 1).

Data collection and statistical analysis

Soil inorganic nitrogen: Ten (0-30 cm) soil samples were collected from each replication at CC establishment, from whole plots at CC biomass collection, and at corn planting (Table 1). After corn planting two (0-30 cm) soil samples were collected monthly from May – October from

subplots of the zero N strategy. Samples were analyzed for nitrate-N (cadmium reduction) (Huffman and Barbarick, 1981).

CC measurements: Cover crop measurements included biomass production and total tissue N content at the time of termination (Table 1). Within each whole plot replicate radish (shoots and roots) and oat (shoots only) cover samples were collected within 5 d of termination in the autumn and again in mid-March (oats only) using three 0.25 m² quadrats. At sampling, radish roots were excavated and washed with water. Fresh weights of cover subsamples were recorded and tissues dried at 60°C and ground using a 1-mm mesh screen. Nitrogen uptake was calculated as the product of percent N and biomass (dry basis).

Corn measurements: Active canopy sensing was conducted to determine normalized difference vegetation index (NDVI) at V6 and V10 using a GreenSeeker® Model 505 handheld red-band optical sensor (Trimble Agriculture Div., Westminster, CO) (Table 1). Chlorophyll content was assessed at R1 to indicate N status using a Minolta SPAD 502 chlorophyll meter (CM) (Konica Minolta, Tokyo, Japan) (Scharf et al., 2006). Corn plant height was measured at V6 and R2 as an average of 20 randomly selected plants plot⁻¹. Measurements were taken from the soil surface to the top of the newest fully developed leaf (height of leaf curve apex) with visible collar (V6) or to the top of the tassel (R2) (Yin and McClure, 2013).

The center two rows of each plot were harvested with a Massey Ferguson 8XP research plot combine (Kincaid Equipment Manufacturing, Haven, KS) to determine grain yield, moisture, and test weight. Yield data were reported at 155 g kg⁻¹ moisture. Treatment profitability was

calculated as net return = gross return from yield – input costs. Grain prices and input costs were determined each year using spring N prices (USDA – AMS 2015; 2016), autumn grain prices (USDA – NASS 2015; 2016), and application costs (http://msue.anr.msu.edu/topic/farm_management/firm_publication_archive, Michigan State University, East Lansing, Michigan; verified 23 May 2017) while CC and manure prices were obtained from local vendors.

Data were analyzed in SAS (Ver. 9.4, SAS Institute, 2011) using the GLIMMIX procedure. Corn measurements assumed fixed effects of year, cover, and N management, and random effects of block and block x cover (SAS Institute, 2011). Cover crop influence on soil inorganic N data were analyzed at the whole plot level, assumed block as a random effect, and were compared at each sample time and not over time. Normality and homoscedasticity assumptions were validated with the UNIVARIATE procedure and Levene's test, respectively, using residuals ($P \leq 0.05$). The LINES option of the slice statement was used to separate treatment means when ANOVA indicated a significant interaction ($P \leq 0.10$). Pearson's correlation coefficients were used to investigate linear relationships between monthly soil NO₃-N data and R1 SPAD with grain yield ($P \leq 0.05$) using the CORR procedure.

Results and Discussion

Weather

Following cover crop (CC) seeding, mean Aug. – Nov. 2014 air temperatures were below the 30-yr mean but were above the 30-yr mean from Sept. – Nov. 2015 (Table 2). Cover crops received 27 – 35 mm rainfall within 1 week of planting both years followed by near to above normal

rainfall in August and September. Following May 2015 corn planting, mean June – Aug. air temperatures were 0.8 – 1.2 °C below normal while Sept. – Oct. was 0.8 – 2.2 °C above normal. Precipitation was 28, 116, and 46% above 30-yr means in May, June, and Aug. 2015, respectively, which may have increased N loss due to denitrification in saturated soils. Following May 2016 corn planting, mean monthly air temperatures were 0.1 – 1.6 °C above normal May through September. In contrast with 2015, 39 and 80% below normal precipitation in May and June, respectively, produced dry soil conditions while 15 – 94% above normal precipitation from July – September produced wet soils later into the growing season.

Autumn CC biomass and nitrogen uptake, and soil NO₃-N

Data were combined across years due to no interaction between CC and year for dry matter production (above-ground only for oats), N uptake, and November soil NO₃-N measurements ($P \leq 0.10$) (Table 3). Cover crop biomass production represented 74 – 81 d of growth each year. Following autumn termination, mean radish dry matter production (i.e., above- and belowground biomass) was 68% greater than aboveground oat CC biomass which increased total N uptake 56% relative to the oat CC. Radish CCs reached anthesis each year while oats did not and may explain greater radish CC biomass observed (O'Reilly et al., 2012). In contrast, radish and oat biomass (herbage only) or N uptake did not differ in Indiana research (Vyn et. al 2000), suggesting that the growth of non-leguminous CCs was limited by low soil NO₃-N (4.4 to 8.3 mg kg⁻¹) concentrations following wheat harvest. Soil NO₃-N concentrations at CC planting in the current study were 8.7 to 12.7 mg kg⁻¹ and supports the greater radish and oat biomass production and N uptake values observed. At autumn termination, soil NO₃-N concentrations (0 – 30 cm) were reduced 78 to 84% relative to no cover crop, when averaged across CCs. Despite

reduced biomass and N uptake compared to radish, oats reduced soil NO₃-N levels an additional 25% suggesting that a substantial amount of N remained in the oat roots which were not collected. Both CCs immobilized soil NO₃-N compared to no cover which suggests opportunities for tightening of the N cycle and reduced post-harvest N losses (Vyn et al., 2000; Weinert et al., 2002; O'Reilly et al., 2012).

Spring CC biomass and nitrogen uptake, and monthly soil NO₃-N

Radish residues decomposed prior to March biomass sampling (Dean and Weil, 2009; Hill et al., 2016). Insufficient radish residues suggested decomposition was rapid in Michigan and that radish N may have been subject to denitrification, leaching, or volatilization N loss mechanisms during the residue degradation process. Approximately 50% of oat residue remained in March (2.9 Mg ha⁻¹, 21% CV) with 38% of the autumn total N uptake remaining immobilized in the biomass (45.9 kg N ha⁻¹, 18% CV). Growers may need to consider different N management strategies to synchronize CC N availability with subsequent crop N uptake.

In June data indicated a significant CC x year interaction on soil NO₃-N levels likely due to contrasting precipitation and therefore data are presented by year (Table 4). In wet soils (i.e., 2015), radish and oat CCs reduced June soil NO₃-N (28 – 32%) relative to the no cover while in dry soils (i.e., 2016) a 19% reduction was observed with oats only. It is unclear whether dry soils in May and June (i.e., 2016) slowed CC biomass decomposition and reduced radish CC N immobilization or if dry soils reduced opportunities for N loss and offset N immobilization in the radish CC. In both years soil NO₃-N reductions indicate N immobilization due to CC decomposition may have contributed to reduced N availability (Ketterings et al., 2015). June soil

NO₃-N levels were the most correlated with corn grain yields relative to other months ($r=0.73$; $P < 0.001$) and corroborated previous findings (Vyn et al., 1999). Although rapid corn N uptake occurs V10 to V14 corn yield potential can be affected at V6 (Binder et al., 2000; Bender et al., 2013). Reduced N availability at V6 has resulted in unrealized yield potential (Binder et al., 2000). In the current study, reduced June nitrate levels in the no cover zero N control suggest N availability may not synchronize well with N needed for early corn growth.

Except for June, year did not interact with CC response and data were combined across years (Table 4). Soil NO₃-N levels (Mar., May, Jul., Aug., and Sept.) were not affected by planting a cover crop, suggesting that radish and oat CCs did not increase NO₃-N availability during the ensuing corn growing season (Vyn et al., 2000; O'Reilly et al., 2012; Gieske et al., 2016a). Reduced CC residue for biomass sampling in March indicate winter decomposition and may help explain no increase in CC soil NO₃-N availability to the ensuing corn cash crop. Between November and March a net loss of 2.9 mg kg⁻¹ soil NO₃-N occurred in the no CC control while an increase of 2.3 and 2.5 mg kg⁻¹ soil NO₃-N due to oat and radish CCs, respectively, was observed and suggested residue mineralization contributed to the N pool. Soils in Lansing, Michigan, may not freeze during the winter due to location among the Great Lakes. Studies indicate CC decomposition during freeze-thaw cycles in the winter can supply labile C and N and stimulate N₂O emissions (Thomas et al., 2017). Oats reduced soil NO₃-N levels 23, 29, and 12% in May, June (2016), and Sept., respectively, compared to radish indicating less N availability to corn. Studies suggest soil NO₃-N levels due to oat CC were likely influenced by a slower rate of residue decomposition due to increased lignin concentrations and recalcitrant N contained in root residues (Malpassi et al., 2000; Jahanzad et al., 2016). Soil NO₃-N data suggest

radish and oat cover crops did not provide an N credit to the ensuing corn crop and growers should not decrease N rates (Andraski and Bundy, 2005; O'Reilly et al., 2012).

Corn response to CCs and N strategies

Grain yield: A cover x N strategy interaction ($P=0.071$) indicated that N strategy may need to be adjusted for specific CC treatments (Table 5). Year was not a significant factor thus data were combined across years and presented for cover by N strategy (Table 6). Within N strategies, radish and oat CCs did not improve and sometimes reduced yield relative to the no CC control and suggested that N availability was not synchronized with corn N uptake. Few differences between those strategies receiving N compared to the zero N strategy indicated that the N rate used in the study (179 kg N ha^{-1}) was sufficient to offset N immobilization from the cover crops (O'Reilly et al., 2012). Cover crop effects on grain yield were most apparent for the zero N strategy ($P=0.025$) as radish and oat reduced grain yield 12 and 14%, respectively, compared to the no CC control. Vyn et al. (2000) observed 15 – 22% grain yield reductions from an oat CC compared to a no cover control receiving zero N fertilizer. Cover crop biomass production has explained some variability in subsequent grain yield in previous studies ($r= -0.24$; $P=0.05$) (Hill et al., 2016). In the current study aboveground radish biomass accounted for 58% (i.e., 5.8 Mg ha^{-1}) of total production compared to $0.89 - 1.27 \text{ Mg ha}^{-1}$ observed by Vyn et al. (2000) implying that total cover crop biomass production limits grain yield. Growers sometimes apply N to CCs seeded prior to a corn cash crop to maximize growth. Results of the current study suggest increased CC growth may reduce N to a corn cash crop and N applied to encourage CC growth may reduce corn yield gains. Reduced soil $\text{NO}_3\text{-N}$ levels at June soil sampling time (previously mentioned) suggest that CCs may have limited N availability during rapid N uptake and reduced

yield potential of the zero N strategy. When mean grain yield of plots not receiving N was subtracted from mean grain yield of plots receiving N, radish and oat CCs increased yield response to N fertilizer by 63 – 79%. Reduced yields in non-fertilized radish and oat and increased yield response to N suggest N availability to corn was reduced where CCs were present. Earlier termination of actively growing cover crops or later corn planting dates may lessen the negative effects of CCs on grain yield, but these practices may not be in agreement with conservation efforts if the goal of non-leguminous cover crops is to produce maximum autumn growth for reduced soil NO₃-N availability.

Corn yield was reduced 3 – 5% in the nitrogen PPI treatment in oats as compared to the no cover or radish treatments. When 52 kg N ha⁻¹ was applied at planting with full SD delayed until V11, radish and oat CCs reduced yield 3 – 4% compared to the no cover control. However no differences amongst CCs were observed in V11 SD followed PL or when strategies included V4 SD, indicating that CCs reduced longevity of the 5x5 placement required to supply N until V11 SD time. Success of delayed SD N (i.e., V11) may require a sufficient N supply to meet corn demands prior to SD timing to maintain yield potential (Binder et al., 2000). In Michigan 5x5 starter placement and SD N is a recommended two-pass fertility strategy to reduce risk of early N loss. Corn preceded by radish and oat cover crops may require increased starter N rates (> 52 kg N ha⁻¹) or slowly-available N sources (e.g., PL) when using delayed N applications (i.e., V11).

Across N strategies, N fertilizer increased grain yield ≥ 2.2 Mg ha⁻¹ and confirmed corn N response within each CC treatment. Within the no CC treatment no differences among N

strategies indicated no benefit to split-applied N although SD N is a recommended corn N strategy in Michigan to increase N recovery (Warncke et al., 2009). In the radish CC, yield was reduced 4.5% in the 5x5 starter with V11 SD reduced yield 4.5% from the PL and indicated that a slowly mineralizable N source may help maintain yield potential under variable weather conditions. Other than PPI N reducing yield 3.9% relative to the 3 pass system (i.e., 5x5 and split SD), few differences were observed within the oat cover crop. Nitrogen applied PPI following oats may have been partially immobilized by decaying oat residue explaining the greater yield decrease from this treatment. No yield difference between strategies within the no cover treatment yet differences following radish and oat suggests that CCs can impact N application method. The 5x5 plus V4 SD and PL plus V11 SD strategies consistently resulted in the greatest yields among CC treatments, but few differences when no CC was present.

Profitability: A CC x N strategy interaction occurred when total treatment costs were subtracted from treatment gross returns (Table 5). Data were combined across years due to no interaction with year. Across years net return was better correlated with grain yield ($r = 0.85$; $P < 0.01$) than treatment application costs ($r = -0.13$; $P = 0.11$) or total N costs ($r = 0.20$; $P = 0.01$). The significant correlation indicated that profitability was more dependent upon grain yield and suggests when N rate is held constant, strategies that maximize grain yield maximize profitability.

Within N strategies, radish and oat CCs did not increase corn profitability compared to the no cover treatment (Table 7). With the exception for PL in radish, net returns were reduced across strategies receiving N fertilizer with radish (\$129 to 214 ha⁻¹) and oat (\$163 to 275 ha⁻¹) relative

to the no cover. Within the PPI N strategy, oats reduced net return relative to radish by \$97 – 106 ha⁻¹. Seed costs of oats averaged \$40 ha⁻¹ greater than radish and contributed to net return reductions. Net return differences were most pronounced within the unfertilized strategy where CC reduced return \$347 - \$427 ha⁻¹. In intensively managed cropping systems inputs may be applied with little data to support agronomic decisions (Orlowski et al., 2016). Radish and oat CCs were not able to improve profitability of N strategies. Future studies should incorporate leguminous CC with N strategies to determine if N availability in corn and if yield and profitability increase. The PL and V11 SD strategy resulted in the lowest net return regardless of CC treatment. While yield consistencies with PL and V11 SD strategy did not correspond to profit consistencies within CC treatments, reduced net returns likely resulted from the increased costs associated with PL. The PL N source cost 333 and 384% more per kg N than urea and 28% UAN-N sources, respectively, while the treatment increased costs over other N fertilizer strategies (\$178 to 232 ha⁻¹). Growers with access to inexpensive manure sources similar to that used in this study may increase their potential to impact profitability. The urea-N and 5x5 starter plus V4 SD strategies consistently resulted in the highest net returns although sometimes results were similar to other strategies within each CC. The urea PPI strategy was the least expensive among N fertilizer strategies and reduced cost (\$20 ha⁻¹) compared to the 5x5 and V4 SD strategy. However reduced yields associated with the oats and urea PPI strategy may hinder adoption by growers and favor the 5x5 plus V4 SD as yield loss is perceived as a larger risk than N savings (Rutan and Steinke, 2017).

Normalized difference vegetation index

Active canopy sensing at V6 indicated both CC and N strategy affected corn normalized difference vegetation index (NDVI) and results were combined across years for N strategy

(Tables 5, 8). The PL (prior to SD) strategy increased NDVI 8.2% relative to zero N and was similar to other strategies that had received N (i.e., urea PPI or 5x5 and V4 SD). Red-NDVI is sensitive to green biomass and an indicator of plant vigor (Tucker, 1979). Despite contrasting soil moistures at sensing time each year (i.e., 0.23 and 0.09 cm³cm⁻³ in 2015 and 2016, respectively) increased V6 NDVI with PL may indicate that either mineralization was sufficient to increase plant vigor or trace amounts of other elements (i.e., 0.42, 0.57, 7.2% S, Mg, and Ca, respectively, and 30, 103, 592, 350, and 256 mg kg⁻¹ B, Cu, Fe, Mn, and Zn, respectively) were sufficient to increase NDVI and the capacity to maintain yield potential until SD time.

The radish CC differentially affected corn V6 NDVI response each year (P=0.01) (Table 9). In 2015 radish and oat CCs increased NDVI 22 – 26% relative to the no cover while in 2016 the oat cover increased NDVI 23 – 25% relative to the radish and no CC control. Plant heights at V7 were similar to V6 NDVI measurements across years in that radish and oat increased height 17 – 29% (P=0.01) relative to no cover in 2015 while oat increased height 17 – 22% (P=0.01) in 2016 (data not shown). Increased corn growth with both CCs may have resulted from a rotation effect rather than N contributions (Andraski and Bundy, 2005). In the 2016 dry spring, phosphorus deficiency was observed on V4 corn preceded by radish and the no CC control but not in oats. Phosphorus availability and uptake requires soil moisture for diffusion and indicate oat residues may have increased soil moisture thus limiting P deficiencies. Previous Michigan studies observed increased soil volumetric water content at planting due to previous radish (29%) and cereal rye (79%) residues (Hill et al., 2016). In the current study however, increased early-season corn growth following CCs did not translate into increased yield.

Chlorophyll content (SPAD)

A significant CC x N strategy interaction affected SPAD chlorophyll meter measurements across years (Table 5). Chlorophyll meter (SPAD) values observed at R1 were well correlated with grain yield ($r=0.89$; $P<0.01$) and suggested N status at tasseling impacts grain fill. Within N strategies the effect of CCs on R1 SPAD values were most pronounced in the zero N strategy where radish reduced SPAD values 4% from the no CC control while oats reduced values 6 and 10% from the radish and no CC treatments, respectively (Table 10). Reduced SPAD values in the zero N strategy suggested 1) both CCs augmented R1 N stress and oats reduced N availability to a greater extent compared with radish and 2) demonstrate the ability of N fertilizer to mask the effect of CCs on soil N fertility as suggested by Duiker and Curran (2005). The application of N fertilizer decreased the immobilization of N from radish and oat residues further diminishing any negative effects on SPAD values. Relative to radish, the oat CC reduced SPAD values 3.2% with PPI N applications which corresponded to reduced yield and indicated that the PPI strategy was not able to supply sufficient N to overcome reduced soil NO₃-N levels and limited R1 yield potential. Full SD delayed until V11 following oats reduced SPAD values 2.7 – 3.5% relative to the no cover. In order to maintain yield potential, success of the late SD timing (i.e., V11) may depend upon sufficient N supply up until SD. When a 5x5 starter and V11 SD strategy were utilized reduced SPAD (oats) values and yield (radish and oats) relative to the no cover control suggested CCs reduced the ability of 5x5 starter to supply N sufficient to maintain yield potential up to V11 SD. However, when PL was utilized as an N source a lack of yield differences suggested a slowly available N source may have continued to provide post-tasseling N to overcome R1 N stress observed in the oat treatment. The addition of N fertilizer increased SPAD values indicating response to N. When SD included a V4 timing, SPAD values were

consistently > 55.0 regardless of CC. Greater SPAD values suggest opportunities to reduce N stress regardless of CC selection and N stress up to R1 caused by CC could be overcome by N fertilizer applications.

Conclusions

Radish and oat covers were non-leguminous species effective at sequestering residual soil NO₃-N following a wheat cash crop. While radish covers had greater biomass production as measured by root and shoot growth, oats were more effective at reducing autumn soil NO₃-N levels.

Although cover crops reduced soil NO₃-N in the autumn it cannot be assumed N assimilated will be protected from loss or available to a following corn cash crop. Increased early corn growth due to covers did not correspond to soil N availability and did not translate into improved corn yields or profitability. In the Michigan corn growing season June soil NO₃-N levels were correlated to grain yield. However cover crops often reduced soil NO₃-N in June and increased yield response to N fertility which suggested reduced N availability to a corn cash crop. Corn yields were unaffected by covers when V4 SD followed 5x5 starter N (52 kg N ha⁻¹) but reduced when full SD at V11 followed 5x5 starter or when urea-N followed an oat cover. Despite few yield differences observed among N strategies within cover crop species reduced N availability suggest years of increased N loss opportunities may exacerbate soil NO₃-N reductions due to cover crops and increase grain yield reductions. Results of the current study suggests single species cover crops in winter wheat – corn rotations may be less appropriate as tools for corn N fertility management but more appropriate as tools for soil conservation and water quality management. Contrasting precipitation observed each spring illustrated the impact of climate variability on soil moisture and growing season conditions. Regardless of cover crop use and

weather variability growers are likely to increase opportunities for consistent yield and profitability gains when N strategies include the use of 5x5 starter N and full SD early (i.e., V4) as well as PL and V11 SD if the manure source is not cost prohibitive.

Two site-years of data which included wet and dry soils suggest further studies are needed to improve corn N fertility management when CCs are planted. Varying termination dates, seeding rates, and tillage methods may improve synchrony of CC N availability with corn N uptake.

Acknowledgements

The authors would like to thank the USDA National Institute of Food and Agriculture, Michigan AgBioResearch, the Michigan State University's College of Agriculture and Natural Resources, and Corn Marketing and Promotion of Michigan for partial funding and support of this research.

APPENDIX

Table 3.01. Dates of field operations and observations in 2014 – 2015 and 2015 – 2016, Lansing, MI.

Operation	2014 – 2015	2015 – 2016
	————chronological order————	
Tillage	08 Aug	29 Jul
Pre-cover soil sample	14 Aug	17 Aug
Cover planting	14 Aug	17 Aug
Fall soil sampling	03 Nov	30 Oct
Cover termination	05 Nov	04 Nov
March soil sampling	20 Mar	18 Mar
Lime application	23 Mar	05 Apr
Spring tillage	15 Apr	17 May
Corn planting and soil sampling	01 May	17 May
V4 sidedress	02 Jun	06 Jun
June soil sampling	03 Jun	14 Jun
V6 observations	10 Jun	15 Jun
V10 observations	28 Jun	01 Jul
V11 sidedress	29 Jun	05 Jul
July soil sampling	01 Jul	15 Jul
R1 observations	20 Jul	22-Jul
R2 observations	29 Jul	05 Aug
August soil sampling	03 Aug	15 Aug
September soil sampling	05 Sep	20 Sep
October soil sampling	02 Oct	11 Oct

Table 3.02. Mean monthly and 30-yr precipitation and temperature data for Lansing, MI, 2014 – 2016.

Month	precipitation [†]				air temperature [‡]			
	2014	2015	2016	30 yr	2014	2015	2016	30 yr
			---mm---				---°C---	
Jan.	na [§]	15	22	46	na	-6.9	-3.5	-4.6
Feb.	na	1	18	38	na	-12.3	-1.8	-3.2
Mar.	na	13	101	45	na	-0.1	4.9	1.9
Apr.	na	23	75	73	na	8.2	7.5	8.7
May	na	109	52	85	na	16.3	14.8	14.7
June	na	192	18	89	na	19.2	20.3	20.0
July	na	61	96	83	na	20.9	23.0	22.1
Aug.	86	123	163	84	20.2	20.3	22.9	21.3
Sept.	85	95	106	92	15.4	19.1	17.4	16.9
Oct.	57	48	82	70	9.2	11.2	9.2	10.4
Nov.	37	31	na	71	0.8	6.9	na	4.6
Dec	35	59	na	42	-0.5	4.0	na	-1.9

[†]Precipitation and air temperature data were collected from Michigan Automated Weather Network (<http://www.agweather.geo.msu.edu/mawn/>).

[‡]30-yr mean sources for precipitation and air temperatures: NOAA (<https://www.ncdc.noaa.gov/cdo-web/datatools/normals>; verified 17 Aug. 2017).

[§]Data not applicable. Monthly data start with August as cover crop planting began the trial period each year.

Table 3.03. Main effects of cover crop on dry matter biomass production, total N uptake, and soil residual NO₃-N concentrations following autumn growth, Lansing, MI, 2014 – 2015.

Cover crop	Dry matter	Total N uptake [‡]	soil NO ₃ -N
	Mg ha ⁻¹	kg ha ⁻¹	mg kg ⁻¹
No cover [¶]	---	---	6.88 [§] a
Radish	10.0 a [#]	189 a	1.50 b
Oats	5.9 b	121 b	1.13 c
CV (%)	34	37	98
Source of variation (<i>P</i> > <i>F</i>)			
Cover crop (C)	<.001	0.007	<.001
Year (Y)	0.013	0.011	0.564
C x Y	0.152	0.275	0.128

[†]Cover crop residues were sampled on 03 Nov 2014 and 30 Oct 2015.

[‡]Total N uptake relative to aboveground biomass only for oat covers.

[§]Soil NO₃-N data transformed to meet normality assumption using root function. Untransformed means are presented. CV represents untransformed data.

[¶]Radish biomass was collected above and belowground in contrast to oats which were aboveground only.

[#]Means followed by same letter in a column are not significant different at alpha=0.10.

Table 3.04. Impact of preceding cover crops not receiving N fertilizer on ensuing monthly[†] soil test nitrate (NO₃-N) (0 – 30 cm) concentrations and linear correlation to grain yield (*r*-value), Lansing, MI, 2015 – 2016.

Cover crop	Mar.	May	June	July	Aug. [‡]	Sept.
	mg kg ⁻¹					
No Cover	3.94 a [§]	6.56 ab	14.92 . [¶]	4.93 a	2.24 a	1.71 ab
Radish	3.95 a	7.81 a	12.90 .	5.38 a	1.80 a	1.73 a
Oats	3.42 a	6.01 b	10.81 .	4.08 a	0.77 a	1.52 b
CV (%)	33	29	32	76	153	23
<i>r</i> -value	0.55**	0.50**	0.73***	0.02	0.18	0.61**
	<i>P</i> > <i>F</i>					
Cover crop (C)	0.317	0.088	0.015	0.680	0.359	0.095
Year (Y)	0.004	0.002	0.001	0.001	0.010	0.036
C x Y	0.219	0.881	0.040	0.905	0.642	0.853
	June C x Y interaction					
			—2015—	—2016—		
No Cover			19.37	a [§]	10.48	a
Radish			13.97	b	11.84	a
Oats			13.18	b	8.44	b
CV (%)			26		21	
<i>P</i> > <i>F</i>			0.035		0.042	

*, **, *** Significant at the 0.05, 0.01, and 0.001 probability levels, respectively.

[†]March samples were collected prior to corn plot establishment.

[‡]August data were transformed using root functions. CV's represent untransformed data.

[§]Means followed by same letter in a column are not significant different at alpha=0.10.

[¶]No means separation due to significant cover x year interaction. Data presented separately.

Table 3.05. ANOVA of cover crop, N strategy, and year fixed effects and their interactions on corn V6 NDVI, R1 SPAD, R2 height, grain yield, and net return in Lansing, MI, 2015 – 2016.

Fixed effects	V6 NDVI	R1 SPAD	R2 height	Grain yield	Net return
	Source of variation ($P > F$)				
Cover crop (C)	0.001	0.006	0.352	0.035	<.001
N strategy (N)	0.014	<.001	<.001	<.001	<.001
Year (Y)	0.060	<.001	<.001	0.002	<.001
C x N	0.367	0.038	0.061	0.071	0.060
C x Y	0.077	0.124	0.150	0.198	0.188
N x Y	0.672	<.001	0.015	0.129	0.187
C x N x Y	0.282	0.728	0.521	0.628	0.553

Table 3.06. Cover crop and N strategy interaction on corn grain yield (155 g kg⁻¹ moisture) and impact of cover crop on response to N, Lansing, MI[†], 2015 – 2016.

N strategy	Cover crop						<i>P</i> > <i>F</i>
	No Cover		Radish		Oat		
			Mg ha ⁻¹				
Zero N	12.7	b [‡] A [§]	11.2	cB	10.9	cB	0.025
PPI	15.3	aA	15.1	abA	14.6	bB	0.077
PL [¶] + V11 SD	14.9	aA	15.5	aA	15.0	abA	0.280
5x5 starter [#] + V4 SD	15.0	aA	15.1	abA	14.8	abA	0.774
5x5 starter + V11 SD	15.4	aA	14.8	bB	14.9	abB	0.061
5x5 starter + V4/V11 SD	15.1	aA	15.1	abA	15.2	aA	0.903
<i>P</i> > <i>F</i>	<.001		<.001		<.001		--
Response to N ^{††}	2.4	B	3.9	A	4.3	A	0.034

[†]Strategies requiring N fertilizer received total maximum return to N rate of 179 kg N ha⁻¹ in all years.

[‡]Means within each column followed by same lower case letters are not significantly different at alpha ≤ 0.10.

[§]means within each row followed by same upper case letters are not significantly different at alpha ≤ 0.10.

[¶]PL = poultry litter applied at 2.2 Mg ha⁻¹ PPI

[#]5x5 starter was applied 5 cm beside and 5 cm below the corn seed furrow at planting (45 kg N ha⁻¹)

^{††}Response to N determined as mean yield from plots receiving N fertility minus yield from Zero N plots within each respective cover.

Table 3.07. Interaction of cover crop and N strategy on corn profitability, Lansing, MI†, 2015 – 2016.

N strategy	Cover						<i>P</i> > <i>F</i>
	No Cover		Radish		Oat		
	US\$ ha ⁻¹						
Zero N	1690	b‡A§	1343	dB	1263	cB	<.001
PPI	1834	aA	1665	aB	1559	aC	<.001
PL¶ + V11 SD	1555	cA	1489	cA	1392	bB	0.014
5x5 starter# + V4 SD	1773	abA	1644	abB	1574	aB	0.001
5x5 starter + V11 SD	1816	aA	1602	bB	1577	aB	<.001
5x5 starter + V4/V11 SD	1756	bA	1613	abB	1586	aB	0.001
<i>P</i> > <i>F</i>	<.001		<.001		<.001		--

†Strategies requiring N fertilizer received total maximum return to N rate of 179 kg N ha⁻¹ in all years.

‡Means within each column followed by same lower case letters are not significantly different at alpha ≤ 0.10.

§means within each row followed by same upper case letters are not significantly different at alpha ≤ 0.10.

¶PL = poultry litter applied at 2.2 Mg ha⁻¹ PPI

#5x5 starter was applied 5 cm beside and 5 cm below the corn seed furrow at planting (45 kg N ha⁻¹).

Table 3.08. Main effect of N strategy[†] on corn V6 NDVI in Lansing, MI, combined across 2015 – 2016.

N Strategy	V6 NDVI
Zero N	0.3741 bcd [‡]
PPI	0.3918 ab
PL [§] + V11 SD	0.4048 a
5x5 starter [¶] + V4 SD	0.3841 abc
5x5 starter + V11 SD	0.3625 cd
5x5 starter + V4/V11 SD	0.3605 d
<i>P</i> > <i>F</i>	0.014

[†]All strategies requiring N fertilizer received total maximum return to N rate of 179 kg N ha⁻¹ in all years.

[‡]Means followed by same lower case letters are not significantly different at alpha=0.10.

[§]PL = poultry litter applied at 2.2 Mg ha⁻¹ PPI

[¶]5x5 starter was applied 5 cm beside and 5 cm below the corn seed furrow at planting (45 kg N ha⁻¹).

Table 3.09. Corn V6 NDVI measurements as affected by previous cover crop, Lansing, MI, 2015 – 2016.

Cover crop	NDVI			
	2015		2016	
No Cover	0.3547	b†	0.3254	b
Radish	0.4311	a	0.3198	b
Oats	0.4457	a	0.4010	a
<i>P > F</i>	0.007		0.006	

†Means followed by same lower case letters are not significantly different at alpha=0.10.

Table 3.10. Interaction of cover crop and N strategy on SPAD R1 chlorophyll meter readings, Lansing, MI†, 2015 – 2016.

N strategy	Cover crop						<i>P</i> > <i>F</i>
	No Cover		Radish		Oats		
	SPAD index						
Zero N	49.8	d‡A§	48.0	bB	44.9	dC	<.001
PPI	54.7	bcAB	56.1	aA	54.3	bB	0.066
PL¶ + V11 SD	54.5	cA	55.0	aA	52.6	cB	0.009
5x5 starter# + V4 SD	56.1	aA	55.9	aA	55.5	aA	0.721
5x5 starter + V11 SD	55.8	abA	55.2	aAB	54.3	bB	0.091
5x5 starter + V4/V11 SD	56.3	aA	56.2	aA	55.2	abA	0.255
<i>P</i> > <i>F</i>	<.001		<.001		<.001		

†Strategies requiring N fertilizer received total maximum return to N rate of 179 kg N ha⁻¹ in all years.

‡Means within each column followed by same lower case letters are not significantly different at alpha < 0.10.

§Means within each row followed by same upper case letters are not significantly different at alpha ≤ 0.10.

¶PL = poultry litter applied at 2.2 Mg ha⁻¹ PPI

#5x5 starter was applied 5 cm beside and 5 cm below the corn seed furrow at planting (45 kg N ha⁻¹).

LITERATURE CITED

LITERATURE CITED

- Andraski, T.W., and L.G. Bundy. 2005. Cover crop effects on corn yield response to nitrogen on an irrigated sandy soil. *Agron. J.* 97:1239-1244.
- Bender, R.R., J.W. Haegele, M.L. Ruffo, and F.E. Below. 2013. Nutrient uptake, partitioning, and remobilization in modern, transgenic, insect-protected maize hybrids. *Agron. J.* 105:161-170.
- Binder, D.L., D.H. Sander, and D.T. Walters. 2000. Maize response to time of nitrogen application as affected by level of nitrogen deficiency. *Agron. J.* 92:1228-1236.
- Clark, A. 2007. *Managing cover crops profitably*. 3rd ed. Sustainable Agric. Network, Beltsville, MD.
- Combs, S.M., and M.V. Nathan. 2015. Soil organic matter. In: M.V. Nathan and R. Gelderman, editors, *Recommended chemical soil test procedures for the North Central Region*. North Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 12.1-12.6.
- Dapaah, H.K., and T.J. Vyn. 1998. Nitrogen fertilization and cover crop effects on soil structural stability and corn performance. *Commun. Soil Sci. Plant Anal.* 29:2557-2569.
- Dean, J.E., and R.R. Weil. 2009. Brassica cover crops for nitrogen retention in the Mid-Atlantic Coastal Plain. *J. Environ. Qual.* 38:520-528.
- Decker, A.M., A.J. Clark, J.J. Meisinger, F.R. Mulford, and M.S. McIntosh. 1994. Legume cover crop contributions to no-tillage corn production. *Agron. J.* 86:126-135.
- Derby, N.E., F.X.M. Casey, and R.E. Knighton. 2009. Long-term observations of Vadose zone and groundwater nitrate concentrations under irrigated agriculture. *Vadose Zone J.* 8:290-300.
- Dove, A., and S.C. Chapra. 2015. Long-term trends of nutrients and trophic response variables for the Great Lakes. *Limnol. Oceanogr.* 60:696-721.
- Duiker, S.W., and W.S. Curran. 2005. Rye cover crop management for corn production in the Northern Mid-Atlantic region. *Agron. J.* 97:1413-1418.
- Frank, K., D. Beegle, and J. Denning. 2015. Phosphorus. In: M.V. Nathan and R. Gelderman, editors, *Recommended chemical soil test procedures for the North Central Region*. North

- Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 6.1-6.6.
- Franzen, D.W. 2017. Nitrogen extenders and additives. NDSU Extension publication SF 1581. North Dakota State University Extension Service, Fargo, ND.
- Gieske, M.F., D.L. Wyse, and B.R. Durgan. 2016a. Spring- and Fall-seeded radish cover-crop effects on weed management in corn. *Weed Tech.* 30:559-572.
- Gieske, M.F., V.J. Ackroyd, D.G. Bass, D.R. Mutch, D.L. Wyse, and B.R. Durgan. 2016b. Brassica cover crop effects on nitrogen availability and oat and corn yield. *Agron. J.* 108:151-161.
- Hill, E.C., K.A. Renner, and C.L. Sprague. 2016. Cover crop impact on nitrogen availability and dry bean in an organic system. *Agron. J.* 108:329-341.
- Huffman, S.A. and K.A. Barbarick. 1981. Soil nitrate analysis by cadmium reduction. *Communications in Soil Science and Plant Analysis* 12:79-89.
- Jahanzad, E., A.V. Barker, M. Hashemi, T. Eaton, A. Sadeghpour, and S.A. Weis. 2016. Nitrogen release dynamics and decomposition of buried and surface cover crop residues. *Agron. J.* 108:1735-1741.
- Johnson, T.J., T.C. Kaspar, K.A. Kohler, S.J. Corak, and S.D. Logsdon. 1998. Oat and rye overseeded into soybean as fall cover crops in the upper Midwest. *Soil Water Conserv.* 53:276-279.
- Ketterings, Q.M., S.N. Swink, S.W. Duiker, K.J. Czymmek, D.B. Beegle, and W.J. Cox. 2015. Integrating cover crops for nitrogen management in corn systems on northeastern U.S. dairies. *Agron. J.* 107:1365-1376.
- Lawley, Y.E., R.R. Weil, and J.R. Teasdale. 2011. Forage radish cover crop suppresses winter annual weeds in Fall and before corn planting. *Agron. J.* 103:137-144.
- Macdonald, A.J., P.R. Poulton, M.T. Howe, K.W.T. Goulding, and D.S. Powlson. 2005. The use of cover crops in cereal-based cropping systems to control nitrate leaching in SE England. *Plant and Soil* 273:355-373.
- Malpassi, R.N., T.C. Kaspar, T.B. Parkin, C.A. Cambardella, and N.A. Nubel. 2000. Oat and rye root decomposition effects on nitrogen mineralization. *Soil Sci. Soc. Am. J.* 64:208-215.
- McSwiney, C.P., S.S. Snapp, and L.E. Gentry. 2010. Use of N immobilization to tighten the N cycle in conventional agroecosystems. *Eco. App.* 20:648-662.

- Mutch, D.R., and K.D. Thelen. 2003. No-till drilling cover crops after wheat harvest and their influence on next season's corn. Michigan State University Ext. Bull. E-2897.
- O'Reilly, K.A., J.D. Lauzon, R.J. Vyn, and L.L. Van Erd. 2012. Nitrogen cycling, profit margins and sweet corn yield under fall cover crop systems. *Can. J. Soil Sci.* 92:353-365.
- Orlowski, J.M., B.J. Haverkamp, R.G. Laurenz, D.A. Marburger, E.W. Wilson, S.N. Casteel, S.P. Conley, S.L. Naeve, E.D. Nafziger, K.L. Roozeboom, W.J. Ross, K.D. Thelen, and C.D. Lee. 2016. High-input management systems effect on soybean seed yield, yield components, and economic break-even probabilities. *Crop Sci.* 56:1988-2004.
- Pantoja, J.L., K.P. Woli, J.E. Sawyer, and D.W. Barker. 2015. Corn nitrogen fertilization requirement and corn-soybean productivity with a Rye cover crop. *Soil Sci. Soc. Am. J.* 79:1482-1495.
- Peters, J.B., M.V. Nathan, and C.A.M. Laboski. 2015. pH and lime requirement. In: M.V. Nathan and R. Gelderman, editors, Recommended chemical soil test procedures for the North Central Region. North Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 4.1-4.7.
- SAS Institute. 2011. The SAS system for windows. Version 9.4. SAS Inst., Cary, NC.
- Scharf, P.C., W.J. Wiebold, and J.A. Lory. 2002. Corn yield response to nitrogen fertilizer timing and deficiency level. *Agron. J.* 94:435-441.
- Scharf, P.C., S.M. Brouder, and R.G. Hoefl. 2006. Chlorophyll meter readings can predict nitrogen need and yield response of corn in the North-Central USA. *Agron. J.* 98:655-655.
- Schultze, S.R., P. Sabbatini, and L. Luo. 2016. Effects of a warming trend on cool climate viticulture in Michigan, USA. *SpringerPlus* 5:1119.
- Schwartz, M.D., R. Ahas, and A. Aasa. 2006. Onset of spring starting earlier across the Northern Hemisphere. *Glob. Change Biol.* 12:343-351.
- Shiple, P.R., J.J. Meisinger, and A.M. Decker. 1992. Conserving residual corn fertilizer nitrogen with winter cover crops. *Agron. J.* 84:869-876.
- Snapp, S.S., S.M. Swinton, R. Labarta, D. Mutch, J.R. Black, R. Leep, J. Nyiraneze, and K. O'Neill. 2005. Evaluating cover crops for benefits, costs and performance within cropping system niches. *Agron. J.* 97:322-332.
- Strock, J.S., P.M. Porter, and M.P. Russelle. 2004. Cover cropping to reduce nitrate loss through subsurface drainage in the U.S. Northern Corn Belt. *J. Environ. Qual.* 33:1010-1016.

- Thomas, B.W., X. Hao, F.J. Larney, C. Goyer, M.H. Chantigny, and A. Charles. 2017. Non-legume cover crops can increase non-growing season nitrous oxide emissions. *Soil Sci. Soc. Am. J.* 81:189-199.
- Thorup-Kristensen, K. 2001. Are differences in root growth of nitrogen catch crops important for their ability to reduce soil nitrate-N content, and how can this be measured? *Plant and Soil* 230:185-195.
- Tonitto, C., M.B. David, and L.E. Drinkwater. 2006. Replacing bare fallows with cover crops in fertilizer-intensive cropping systems: A meta-analysis of crop yields and N dynamics. *Agric., Ecosystems and Environ.* 112:58-72.
- Tremblay, N., Y.M. Bouroubi, C. Bélec, R.W. Mullen, N.R. Kitchen, W.E. Thomason, S. Ebelhar, D.B. Mengel, W.R. Raun, D.D. Francis, E.D. Vories, and I. Ortiz-Monasterio. 2012. Corn response to nitrogen is influenced by soil texture and weather. *Agron. J.* 104:1658-1671.
- U.S. Environmental Protection Agency. 2016. What climate change means for Michigan. Report No. EPA 430-F-16-024.
- Vetsch, J.A., and G.W. Randall. 2004. Corn production as affected by nitrogen application timing and tillage. *Agron. J.* 96:502-509.
- Vyn, T.J., J.G. Faber, K.J. Janovicek, and E.G. Beauchamp. 2000. Cover crop effects on nitrogen availability to corn following wheat. *Agron. J.* 92:915-924.
- Vyn, T.J., K.J. Janovicek, M.H. Miller, and E.G. Beauchamp. 1999. Soil nitrate accumulation and corn response to preceding small-grain fertilization and cover crops. *Agron. J.* 91:17-24.
- Wagger, M.G. 1989. Time of desiccation effects on plant composition and subsequent nitrogen release from several winter annual cover crops. *Agron. J.* 81:236-241.
- Wagger, M.G., M.L. Cabrera, and N.N. Ranells. 1998. Nitrogen and carbon cycling in relation to cover crop residue quality. *Soil Water Conserv.* 53:214-218.
- Warncke, D., J. Dahl, and L. Jacobs. 2009. Nutrient recommendations for field crops in Michigan. Bull. E2904, Michigan State University Extension, East Lansing, MI.
- Warncke, D., and J.R. Brown. 2015. Potassium and other basic cations. In: M.V. Nathan and R. Gelderman, editors, Recommended chemical soil test procedures for the North Central Region. North Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 7.1-7.3.

- Weinert, T.L., W.L. Pan, M.R. Moneymaker, G.S. Santo, and R.G. Stevens. 2002. Nitrogen recycling by nonleguminous winter cover crops to reduce leaching in potato rotations. *Agron. J.* 94:365-372.
- Williams, S.M., and R.R. Weil. 2004. Crop cover root channels may alleviate soil compaction effects on soybean crop. *Soil Sci. Soc. Am. J.* 68:1403-1409.
- Yin, X., and M.A. McClure. 2013. Relationship of corn yield, biomass, and leaf nitrogen with normalized difference vegetation index and plant height. *Agron. J.* 105:1005-1016.

CHAPTER 4

INFLUENCE OF COVER CROPS AND NITROGEN MANAGEMENT ON THE SOIL MICROBIOME TO OPTIMIZE CORN PRODUCTION

Abstract

Nitrogen fertilizer management may affect soil bacterial community composition in a corn (*Zea mays* L.) rotation but could be altered when following cover crops. Field studies were conducted in 2014 – 2015 (year 1) and 2015 – 2016 (year 2) to evaluate the effects of Daikon radish [*Raphanus sativus* (L). var. The Buster] and Forage oat [*Avena sativa* (L.) var. Magnum] on soil respiration, soil bacterial community diversity, richness, and evenness in response to N management strategies. Strategies consisted of 179 kg N ha⁻¹ applied as urea pre-plant incorporated (PPI), poultry litter (PL) PPI (2.2 Mg ha⁻¹) plus sidedress (SD) N V11, starter N (45 kg N ha⁻¹) applied 5 cm beside and 5 cm below the furrow (5x5) followed by SD at V4, V11, or V4 plus V11, and a zero N control. Cover crop autumn biomass production, C and N uptake, soil base cations, and corn yield were also measured. Forage radish increased autumn biomass production 39 and 120% in years 1 and 2, respectively, compared with oats, but 46% less C uptake by oats in year two resulted in a C:N ratio of 17:1 and 23:1 for oat and radish CC, respectively. Relative to an oat or no CC control radishes reduced soil bacterial diversity during active CC growth, but no differences were observed at corn seeding and R1. Nitrogen strategies affected soil bacterial diversity, richness, or evenness at 4 of 6 observations, compared with only 1 observation with CCs. Grain yields were correlated to R1 soil bacterial diversity in both years regardless of N fertilizer application ($r=-0.21$ to -0.63 ; $P \leq 0.08$) and indicate soil bacterial communities affected corn grain production. The inverse relationship of yield and R1 diversity

suggested the relative abundance of soil bacterial community membership rather than diversity may provide a better indication of yield.

Introduction

The soil is a living ecosystem and functions to sustain healthy plants, animals, and the environment (Doran and Zeiss, 2000). As a living component of soils the microbial biomass provides agricultural services which includes recycling of organic materials and functions as a labile source of carbon (C), N, phosphorus (P), and sulfur (S) (Brookes, 2001). Soil bacteria comprise one of the largest components of the microbial biomass and have direct impact on plant fitness as pathogens or beneficial mutualists, and indirectly as decomposers, or through suppression of deleterious pathogens (Nihorimbere et al., 2011). While soil bacteria include pathogens, beneficials such as genera *Streptomyces*, *Bacillus*, and *Pseudomonas* produce antibiotics antagonistic to other plant pathogens (Rosenzweig, 2014). Plant growth promoting bacteria associated with rhizosphere soil (PGPR) colonize plant roots and stimulate plant growth through N fixation, phytohormone production (e.g., auxin), or reduce the incidence of disease (Naseem and Bano, 2014; Rosenzweig, 2014, Vurukonda et al., 2016). Factors affecting soil bacterial communities may have impact on crop production. Multiple groups of soil bacteria may perform similar functions and influence cropping system resilience and stress adaptation through functional redundancy (Sharma et al., 2010). Cropping system resilience to moisture stress may improve consistency of Michigan corn grain yields. Since 2010 Michigan corn grain yields have varied >10% despite little variation in hectares planted (USDA, 2017). Recent studies suggest moisture stress as a result of increased climate variability during cash crop seasons may explain corn yield variation (Mishra and Cherkauer, 2010). A 70-yr projection for Michigan agriculture

predicts an additional 5 - 15 days per growing season with air temperatures $> 35^{\circ}\text{C}$ combined with increased spring rainfall and more intense summer droughts (USEPA, 2016). Increased understanding of production-related factors that affect soil bacterial communities may provide opportunities to enhance crop resilience to climate change (e.g. moisture stress) and improve grain yield consistency (Song et al., 2015).

Cover crops (CC) have been used to increase plant diversity and influence soil microbial communities (Buyer et al., 2010; Vukicevich et al., 2017). Following Michigan winter wheat harvest, soils may remain fallow for up to a 9 mo. period and provide a niche for CC incorporation. Plants, soils, and soil microbes function simultaneously to influence rhizosphere processes that contribute to plant health and productivity (Chaparro et al., 2012). Plants modulate microbial community composition through root morphology, C-rich root exudates (i.e., rhizodeposition), and litter quality (Berg and Smalla, 2009; Chaparro et al., 2012; Cleveland et al., 2014). Soil microbial communities provide agroecosystem services (i.e., nutrient cycling, carbon transformations, soil structure maintenance, water availability, and regulation of pests and disease) which in-turn influence crop production (Kibblewhite et al., 2008). Tillage, lack of crop rotations, and crop residue removal reduce soil bacterial community diversity (i.e., richness and evenness) (Lupwayi et al., 1998; Ceja-Navarro et al., 2010; Figuerola et al., 2015; Sun et al., 2016). Studies suggest increased bacterial community diversity may influence crop yield and N uptake, enhance functional resiliency, and protect ecosystem services under abiotic disturbance (Eisenhauer et al., 2012; Awasthi et al., 2014; Tautges et al., 2016). Meta-analysis indicated that crop diversity (i.e., rotations) increased microbial richness (i.e. number of species present) (15%) and diversity (3.4%) scores but was unable to conclude whether microbial diversity sustained

ecosystem function (e.g., nutrient cycling) under soil disturbance. Utilizing CCs to influence soil bacterial communities for optimal crop production requires further investigation (Venter et al., 2016).

To improve the synchrony between N availability and corn N uptake, Michigan corn growers utilize multiple N strategies including, pre-plant incorporation, at-plant applications, and multi-pass systems with reduced N rates at planting followed by in-season sidedress (SD) application. Contrasting literature results concerning the impact of N fertilizer on soil bacterial diversity and community composition may implicate both N rate and source (Fierer et al., 2012; Geisseler and Scow, 2014; Ikeda et al., 2014; Li et al., 2016). For example, Ikeda et al. (2014) observed increased bacterial species diversity (i.e., α -Proteobacteria and β -Proteobacteria) from urea-formaldehyde N relative to ammonium sulfate, nitrate-N, or urea. Zhao et al. (2014) observed increased bacterial diversity at the phylum level when manure and mineral N fertilizers were utilized as a combined source. However Orr et al. (2012) indicated organic and mineral N sources had less of an impact on soil bacterial communities as compared to changes in soil temperature and cumulative rainfall received 14 d prior to sampling. Recent data from the Morrow plots (fertilizer and crop-rotation study initiated in 1876) determined that manure increased bacterial diversity and lead to greater relative abundance of bacteria beneficial to plant fitness (Soman et al., 2017). Reduced soil microbial diversity due to N fertilizer applications may be due to changes in soil chemistry and not simply a direct osmotic effect (Lupwayi et al., 2012; Zhalnina et al., 2015; Li et al., 2016). Ammonium-based N fertilizers are a source of soil acidity due to H⁺ release during NH₄⁺ nitrification with varying degrees of acidification depending upon N source (i.e., anhydrous ammonia, urea, etc) (Havlin et al., 2014). Cation uptake by crops

also contributes to soil acidity, which in turn could influence bacterial communities. Soil microbial response to N fertilizer may also depend upon organic C inputs which are used by soil bacteria as a food source (i.e., heterotrophs). Meta-analysis indicated that the addition of mineral N fertilizer to promote crop growth and yield increased crop residue returned to the soil and increased both organic C (13%) and the microbial biomass (15%) (Geisseler and Scow, 2014). In fallow soils N fertilization decreased soil bacterial diversity and reduced copiotrophic (i.e., Actinobacteria) bacteria important for C and N cycling. However, when CCs were used this did not occur, indicating CCs were able to protect soil bacterial diversity from deleterious effects due to N fertilization (Verzeaux et al., 2016). Further research is needed to investigate impacts of N fertilization strategies on both soil bacterial communities and crop yield (Nielsen et al., 2014; Zhao et al., 2014).

High throughput next-generation sequencing (NGS) has enabled researchers to study plant and soil microbiome interactions which are predominated by bacteria not cultivable in laboratory environments (Rosenzweig, 2014). Illumina-based sequencing is a cost effective NGS strategy that can produce up to 500 cycles of a paired-end DNA fragment read while the MiSeq platform can generate 8.5 Gbp using paired 250-nucleotide reads (Kozich et al., 2013). Prior to sequencing, conserved primers which target regions of 16S rRNA are combined with template soil DNA and then amplified (Rosenzweig, 2014). The 16S rRNA gene provides phylogenetic bacterial community soil diversity due to its presence in most bacteria, stability over time, and size (1,500 base-pairs) for informatics purposes (Janda and Abbott, 2007). Sequencing of the 16S rRNA gene has revealed a higher prevalence of *Agrobacterium* spp. in the corn rhizosphere with variations in bacterial richness, diversity, and relative abundance when compared to bulk soil

(Sanguin et al., 2006; Peiffer et al., 2013). Analysis of corn rhizosphere phylogenetic diversity indicated enriched Proteobacteria, Firmicutes, Bacteroidetes, and Cyanobacteria which function to transform labile and recalcitrant nutrient sources (Li et al., 2014). Corn roots exude 5 – 62% of total net belowground C and may alter rhizosphere bacterial community composition, but C quality may depend on growth stage (Amos and Walters, 2006). Prior to tasseling copiotrophic bacteria, which prefer simple amino acids (i.e., r-strategists), predominate until after tasseling when oligotrophic bacteria, which utilize complex carbohydrates (k-strategists) become more abundant. When compared to the bulk soil corn has the capacity to influence and select bacterial communities (Li et al., 2014). Illumina sequencing of corn rhizosphere and bulk soil samples at multiple growth stages using field-derived studies may provide opportunities to assess the impact of cover crops when combined with N management strategies on the microbial community. The objective of this study was to investigate the effects of a radish, oat, and no cover crop in combination with corn N management strategies on corn grain production, spatial and temporal variation in soil CO₂ respiration, and soil bacterial community composition.

Materials and Methods

Site characteristics and cultural practices

Two field experiments were conducted on different sites (2014 – 15 and 2015 – 16) at the Michigan State University South Campus Research Farm in Lansing, MI. Each location consisted of a 2-yr cropping rotation of winter wheat/cover crop (CC) followed by corn on a Capac loam soil (fine-loamy, mixed, active, mesic Aquic Glossudalf). Following July wheat harvest, fields were chisel plowed, disk-harrowed, and leveled (10-cm depth) with a soil finisher. Prior to cover crop planting soils were sampled (20-cm depth), air-dried, and ground to pass

through a 2-mm sieve. Soil properties included 5.8 – 6.0 pH (1:1 soil/water) (Peters et al., 2015), 24 – 55 mg kg⁻¹ P (Bray-P1) (Frank et al., 2015), 122 – 136 mg kg⁻¹ K (ammonium acetate method) (Warncke and Brown, 2015), and 26 – 34g kg⁻¹ organic matter (loss-on-ignition) (Combs and Nathan, 2015).

To control volunteer wheat the following season, CCs were terminated each autumn with glyphosate [N-(phosphonomethyl) glycine]. To control vegetation encroachment, the no CC treatment received an additional glyphosate application prior to CC termination. Each site year received lime (2.2 Mg ha⁻¹) applied 39 to 42 d prior to corn planting to adjust soil pH to 6.5. Broadcast P and K fertilizer were pre-plant incorporated 1-d prior to corn planting with triple super phosphate (0-45-0 N-P-K) and muriate of potash (0-0-62) based on soil tests. Weeds were controlled with acetochlor [2-chloro-N-ethoxymethyl-N-(2-ethyl-6-methylphenyl) acetamide] and glyphosate followed by a second application of glyphosate 17 – 24 d later. On-site environmental data were collected using the Michigan Automated Weather Network (<http://www.agweather.geo.msu.edu/mawn/>, Michigan State University, East Lansing, Michigan; verified 04 Oct. 2017).

Experimental design and treatment application

Eighteen treatments were arranged in a randomized complete block design with four replications. The whole plot factor was CC (no cover crop, ‘The Buster™’ daikon forage radish, or ‘Magnum™’ forage oat [Weaver Seed of Oregon, Crabtree, OR]) and measured 54.9 m by 12.2 m in length. Cover crops were drill-planted (radish and forage oats at 11.2 and 28.0 kg ha⁻¹,

respectively) 14 Aug. and 17 Aug. in 2014 and 2015, respectively, with a Gandy orbit-Air Seeder coupled with John Deere double disk openers in 19 cm rows.

The subplot factor consisted of six N strategies that measured 4.6 m by 12.2 m in length. Total N rate was equalized to the maximum return to N rate (MRTN) (179 kg N ha⁻¹) (Steinke, 2015). Nitrogen management strategies included PPI N (1-d prior to corn planting) as 100% urea, dried poultry manure (4-3-2-0.42S-0.57Mg-7Ca) (PM) (2.2 Mg ha⁻¹) applied PPI plus V11 sidedress (SD), three starter N (45 kg N ha⁻¹ urea ammonium nitrate [UAN; 28-0-0]) treatments applied 5 cm beside and 5 cm below the furrow (5x5) followed by SD at V4, V11, or V4 and V11, and a zero N control. Corn V4 N was UAN coulter injected 5 cm deep and 38 cm to the side of each row. At V11 SD a urease inhibitor (CO(NH₂)₂ + n-(n-butyl) thiophosphoric triamide) (Koch Agronomic Services, LLC, Wichita, KS) was mixed with UAN to prevent N volatilization and banded 10 – 15 cm to the side of each row. Corn was seeded in 0.76 m rows at 84,510 seeds ha⁻¹ with Dekalb DKC48-12 RIB (98 d relative maturity) (Monsanto Co., St. Louis, MO).

Data collection and processing

Corn: The center two rows of each plot were used to determine V3 corn density (plants m⁻¹) and harvested for grain yield, moisture, and test weight on 14 Oct. and 18 Oct. 2015 and 2016, respectively, using a Massey Ferguson 8XP research plot combine (Kincaid Equipment Manufacturing, Haven, KS). Final grain yields were corrected to 155 g kg⁻¹ moisture.

Cover crops: Whole radish plants and oat canopies were sampled for biomass production and total tissue N and organic C content within 5 d of autumn termination and again mid-March (oats

only) from three 0.25 m² quadrats. Cover crop tissues were dried at 60°C, ground using a 1-mm mesh screen, and sent for commercial analysis (A&L Great Lakes Laboratories, Inc., Fort Wayne, IN) of total C and N via dry combustion. Total C and N uptake were calculated as the product of tissue concentration and dry biomass.

Soil chemical properties: Soils were sampled for NO₃-N (cadmium reduction) (30 cm depth) (Huffman and Barbarick, 1981), and P, K, Mg, Ca, pH, and CEC (20 cm depth) (previously described). Soils were sampled from each replication at CC establishment, from whole plots during autumn and mid-March (inorganic N only) simultaneous to cover crop biomass sampling, during May corn planting, and from subplots not receiving N fertility June through October (inorganic N only).

Soil basal CO₂ respiration and labile amino-N (SLAN): Soils were sampled (15 cm depth) from each replication and analyzed for CO₂ respiration and SLAN (Solvita®, Mt Vernon, ME) at cover crop establishment, from whole plots at autumn biomass sampling and May corn planting, and during corn tasseling (R1) and harvest (R6) (38 cm from corn row). Observations in the autumn of 2014 and May of 2015 followed a protocol which required soils to be dried, ground, and rewet but was changed for soil samples thereafter to the ‘basal’ protocol which utilized unground soils at field moisture levels (https://solvita.com/wp-content/uploads/2014/04/Solvita-and-Soil-Structure_Rewetting-2015.pdf, Solvita® & Woods End Laboratories; verified 02 Nov. 2017). For the basal field test, 6 – 10 soil samples were collected and homogenized and a 110 g subsample was used for processing at field soil moisture

levels. Soils used for SLAN testing at R1 and R6 were sampled from the zero N strategy only and required soils to be dried.

Soil bacterial phylogenetic diversity: Soils were sampled (10 cm depth) for bacterial DNA analysis at cover crop establishment and termination, corn planting, R1, and R6. Soils at R1 and R6 were sampled from subplots and included bulk soil (i.e., 38 cm from corn row) and rhizosphere soil (i.e., between two corn plants within a row). Soils were air-dried and ground to pass through a 2-mm sieve.

Soil genomic DNA was extracted from 0.25 g with the MO-BIO Power Soil® kit (MO BIO Laboratories, Carlsbad, CA) and stored at -20°C. Polymerase chain reaction (PCR) was used to amplify the V4 hypervariable region (length 250 bp) of the 16S gene using genomic DNA as a template and high fidelity polymerase (Accuprime™ Pfx Supermix, ThermoFisher Scientific, Waltham, MA). Amplicons and libraries were constructed per previously described methods (Kozich et al., 2013). The MiSeq Illumina platform (San Diego, CA) was used to sequence amplicon libraries submitted to the Michigan State University Research and Technology Support Facility for next-generation sequencing (East Lansing, MI). Sequence data were processed through the Michigan State University High Performance Computing Center (East Lansing, MI) and analyzed using a previously described analysis pipeline (Kozich et al., 2013) and protocol (http://www.mothur.org/wiki/MiSeq_SOP; verified 5 Oct. 2017) with the mothur software package (ver. 1.33.2b; Schloss et al., 2009). Operational taxonomic units (OTU's) were based on a 97% sequence identity and classified to the SILVA database. Soil bacterial alpha diversity indicators (i.e., membership of taxa within a single sample) were assessed to describe diversity,

richness, and evenness. The inverse Simpson's diversity index (ISD) was used to describe bacterial diversity and represented the number of uniformly abundant OTUs needed to observe the same level of diversity found in the community (Schloss and Handelsman, 2006). Soil bacterial community richness and evenness were assessed using the Chao1 estimator and Shannon index, respectively.

Data analysis

Data were subjected to analysis of variance using the GLIMMIX procedure in SAS (ver. 9.4; SAS Institute, 2011). Whole plot data considered cover crop and year as fixed effects and assumed random block effects. Additionally, subplot data (soil respiration, stand density, and corn yield) included N management strategy as a fixed effect and block x cover as a random effect. Microbial DNA sequencing data were processed separately by year due to computational load (Smith et al., 2016) and included soil sampling time and sampling zone as additional fixed effects in the model. Normality and homoscedasticity assumptions were validated with the UNIVARIATE procedure and Levene's test, respectively, using residuals ($P \leq 0.05$). When ANOVA generated a sig. F value ($P \leq 0.10$) treatment means were separated using the LINES option of the LSmeans statement. The SLICE statement was used to investigate interacting effects ($P \leq 0.10$). The CORR procedure was used to generate Pearson correlation coefficients for investigating relationships between edaphic variables and grain yield each year ($P \leq 0.10$).

Results and Discussion

Environmental conditions

Each year cover crops (CC) received 27 – 35 mm cumulative rainfall within 1-wk after seeding in Aug. that helped to aid in establishment (data not shown). Following CC seeding cumulative rainfall in Aug. and Sept. was 5 mm below the 30-yr mean in 2014 and 42 mm above in 2015 (Table 1). Oct. and Nov. received a cumulative rainfall deficit 47 and 62 mm below the 30-yr in 2014 and 2015, respectively. Air temperatures in the non-cash crop growing season (Aug. – Nov.) were 1.0 – 3.7 °C below the 30-yr mean in 2014 and 0.7 – 2.3 °C above in 2015 (except Aug.).

At corn seeding cumulative May and June rainfall was above normal (128 mm) in 2015 but below normal (103 mm) in 2016 and resulted in contrasting wet and dry soils, respectively. August rainfall was 39 – 79 mm above normal in 2015 – 2016. Cumulative Sept. and Oct. rainfall was 19 and 26 mm below and above the 30-yr mean in 2015 and 2016, respectively. Mean air temperatures (Apr. – Oct.) were -0.5 – 2.1 °C of the 30-year mean in 2015 and -1.3 to 1.6 °C in 2016.

Soil min. and max. moisture and temperature were generally higher in year 2 relative to year 1 in the 14-d prior to each soil sampling time except for soil moistures at R1 and soil temperatures at CC planting. Soil moisture at R1 was less in year 2 (relative to year 1) and likely due to deficit May and June rainfall followed by less precipitation in the 14-d prior to R1 observations in 2016 (Table 2). Soil moisture and soil temperature can shift soil bacterial community composition (i.e., β -diversity) with different phyla taxon responding differently to moisture stress and alter

CO₂ efflux in soils under corn management (Ding et al., 2010; Bainard et al., 2016). Yearly variation in soil moisture and temperature at each observation likely influenced soil bacterial community composition each site year.

Studies were conducted on separate fields each year (183-m apart). Despite similar soil series (i.e., Capac loam soil) soil properties varied yearly and within each field. For example prior to CC planting in year 1 standard deviation between replicates for soil organic matter (OM) (g kg⁻¹), pH, K (mg kg⁻¹), and magnesium (Mg) (mg kg⁻¹) was 1.9, 0.15, 21, and 25, respectively. Year 2 standard deviation was 0.22, 29, and 31 for soil pH, K, and Mg, respectively. Soil bacterial diversity (i.e., inverse simpson index) and relative abundance of bacterial phyla can be influenced by soil properties such as OM, pH, K, and Mg and vary spatially within fields (Rosenzweig et al., 2017). Differences in soil edaphic properties (pH, soil moisture, soil temperature, available phosphate) have shifted soil bacterial community composition within a growing season regardless of crop species (Bainard et al., 2016). The current study was conducted for two years on similar sites but variation within fields or between years in soil edaphic properties may have also influenced soil bacterial diversity and relative abundance.

Cover crop biomass production and quality

At autumn CC harvest in 2014 radish biomass was 39% greater compared with oat cover crop (Table 3). Leaf tissue analyses indicated radishes increased total N uptake 63% while oats resulted in an increased C:N ratio. Similar biomass results were observed in 2015 where the radish CC increased production 120% relative to the oat CC. In year 2 (i.e., autumn 2015) oat CC biomass production was reduced 44% relative to a 12% reduction with radishes from year 1

due to crown rust (*Puccinia coronata*) in the oats. Due to less biomass production C and N uptake was reduced 46 and 32%, respectively, and resulted in a lower oat CC C:N ratio relative to the radish. Litter quality (i.e., CC) can influence soil bacterial community composition due to labile and recalcitrant C and N contributions and their relative proportions which provide sources of nutrients and energy (Paterson et al., 2008). Microbial metabolism can become N limited above a 20:1 C:N threshold and immobilize available soil N during CC litter decomposition (Mooshammer et al., 2014). The C:N ratios observed in the current study are near the 20:1 threshold observed in other studies and suggest that both CC's have the potential to affect N cycling. Additionally grass species (i.e., oats) contain more recalcitrant C (i.e., lignin) compared with radishes which has increased the relative abundance of Verrucomicrobia and Actinobacteria and may prolong decomposition and N mineralization (Sagova-Marechova et al., 2011; Jahanzad et al., 2016; Ye et al., 2017).

In March of both years radish residues were nearly decomposed and only oat CC residues were able to be sampled for biomass, and C and N uptake which precluded a formal analysis. In March of 2014 oat CC residues constituted 2429 kg ha⁻¹ biomass, and 40 and 1041 kg ha⁻¹ of N and C, respectively. In 2015, 3415 kg ha⁻¹ oat residues remained and contained 52 and 1319 kg ha⁻¹ of N and C, respectively. Spring C/N ratios of remaining biomass were similar each year (26:1) but higher than observed in the fall indicating a recalcitrant C source for soil bacteria.

Cover crop effects on base cation levels

At autumn CC biomass harvest in 2014, soil test phosphorus (P), potassium (K), magnesium (Mg), and calcium (Ca) levels were 51 – 64, 95 – 105, 155 – 206, and 878 – 1052 mg kg⁻¹,

respectively. After 81-d of CC growth oats had significantly increased autumn soil P levels 25% relative to radishes ($P=0.0912$) while no differences were observed between CC and the no cover control. In contrast, radishes increased autumn soil Ca levels 18 – 20% ($P=0.0308$) relative to other treatments. A previous Michigan study indicated that CC (i.e., radish) can increase soil test nutrient levels, but was after CC termination and incorporation into the soil (Wang et al., 2008). In the current study CC were actively growing and soils were subject to CC root exudation. Studies suggest graminaceous plants root exudates contain phytosiderophores – iron chelating compounds (e.g., mugineic acids) and may help explain increased P levels due to oat CCs (Sugiura and Nomoto, 1984). Mean soil pH at was 6.1 at cover crop sampling and phytosiderophores could have reduced the potential for Fe- and Al-P binding. Other studies have suggested plant root exudates solubilize $FePO_4$ and $AlPO_4$ to release P which may have also contributed to P levels (Otani et al., 1996). Changes in autumn soil Ca and P levels in 2014 illustrate the potential of CC to impact soil properties and indirectly influence soil bacteria communities (Tao et al., 2017). At corn planting in May soil test P, K, Mg, and Ca levels were 48 – 52, 93 – 104, 169 – 173, and 1128 – 1167 but were unaffected by CC's.

At autumn CC biomass harvest in 2015, soil test P, K, Mg, and Ca levels were 20 – 23, 95 – 105, 155 – 206, and 878 – 1052 mg kg⁻¹, respectively, and at corn planting in May were 27 – 30, 164 – 180, 179 – 210, and 990 – 1178 but were unaffected by cover crops.

Cover crop effects on corn grain yield

In 2015, grain yield was affected by N strategy but not by CC despite numerically higher yields in the no cover treatment (Table 4). The zero N strategy reduced corn yields 22 – 24% from

strategies receiving N fertilizer. A single urea PPI application reduced grain yield 3% from strategies receiving a single SD application at V11 likely as a result of increased N loss opportunities with early applied N due to above-normal May (29%) and June (116%) rainfall. In 2016 grain yields were affected by CC's and N strategy. Oats reduced corn grain yield 4% relative to other CC treatments while the zero N strategy reduced grain yield 21 – 23% from other N fertilizer strategies. Unlike 2015 the 2016 field season was dry in May and June as a result of below normal rainfall in May (38%) and June (80%) rainfall. Reduced yields due to oat cover crops suggests that under dry soil conditions residue decomposition rates and subsequent N availability may not be in synchrony with corn N uptake. June soil sampling indicated that radish and oat cover crops reduced soil NO₃-N levels 28 – 32% from the no cover (P=0.0533) in 2015 and 20 – 29% (P=0.0394) (oats only) in 2016 (data not shown). At June soil sampling each year corn was at growth stage V6 to V8 where N stress has reduced yield potential (Binder et al., 2000). Results suggest that cover crops may impact soil N availability in June thus growers should consider crop N uptake periods when increasing plant diversity with using non-leguminous cover crops too stimulate soil bacterial communities.

Cover crop effects on SLAN content and soil basal CO₂ respiration

SLAN: Soil labile amino-N was observed in whole plots under the influence of CC treatments. Pearson correlations indicated SLAN was not affected by soil moisture, temperature, or precipitation experienced 14-d prior to each observation (Table 5). In 2014, the radish CC increased SLAN 15% relative to the no cover control at autumn biomass harvest (Table 6). In 2015 oats increased SLAN 18 and 14% relative to the radish CC and no cover control, respectively. Root exudates provided by actively growing cover crops in the autumn may contain

amino acids and contribute to SLAN (Huang et al., 2014). Soil microbes can directly assimilate free amino acids and increased SLAN due to covers may offer opportunity to affect soil bacterial communities (Jan et al., 2009). Despite non-significant correlations between SLAN and soil moisture, precipitation, or temperature, soil minimum and maximum moisture were correlated with soil NO₃-N ($r=-0.62$ to -0.64 ; $P \leq 0.06$) and NH₄-N were correlated ($r=-0.78$ to -0.79 ; $P \leq 0.01$). Significant correlations with soil NO₃-N and NH₄-N but not SLAN suggest that N contained in SLAN is less affected by climatic factors and a more stable criterion when evaluating CC to impact microbial communities. While observations the following May in 2015 indicated no differences in SLAN, the presence of root and shoot biomass from cover crop residues are a source of proteins for enzymatic amino acid synthesis and microbial assimilation (Jan et al., 2009).

Respiration: Soil respiration was observed in whole plots prior to corn planting and in subplots of the zero-N strategy at R1 and R6. Data will be discussed pertaining to whole plots first (not under the influence of corn) (i.e., autumn and the following May prior to corn planting) for year 1 and year 2, and then subplots for year 1 and year 2. In contrast to Han et al. (2007) Pearson correlations indicated CO₂ was not affected by soil moisture, temperature, or precipitation experienced 14-d prior to observations at CC planting, autumn CC harvest, corn planting, corn tasseling, and corn harvest (Table 5). In 2014 CC did not significantly impact soil respiration at CC harvest time in the autumn or at corn planting the following May in 2015 (Table 6). At CC harvest in the fall of 2015, radish and oat covers increased soil respiration 46 – 61%. Increased autumn CO₂ respiration due to cover crops are an indication of increased microbial activity as

both heterotrophic microbial oxidation of organic C by rhizosphere and bulk soil microbes and CC (autotrophic) root respiration could have contributed to CO₂ efflux (Raich and Mora, 2005).

After corn planting and subplot establishment in 2015 oat and radish cover crops increased soil CO₂ respiration 12 – 15% relative to the no cover control and indicated that CC residues were continuing to decompose between R1 and R6 (Table 7). Nitrogen strategies had no impact on CO₂ respiration, but when values were averaged across CC and N strategies respiration at R1 increased 21% relative to R6. Previous studies have indicated microbial community structure is influenced by corn growth stage with greatest bacterial diversity observed at tasseling and the least at physiological maturity (i.e., R6) (Cavaglieri et al., 2009). Increased CO₂ respiration at R1 may stimulate microbial activity due to the influence of an actively growing corn crop. In 2016 a CC by N strategy interaction was observed (P=0.0374) indicating that CO₂ respiration due to N strategy was affected by the previous CC (Table 8). When splitting the interaction by CC, CO₂ respiration was increased 22% in the radish CC when the early-applied SD N rate (i.e., V4) was reduced and the late-applied SD N rate (i.e., V11) was increased. In the same site year radish cover crops resulted in a higher C:N ratio than oats (previously mentioned). Previous studies have indicated that increased CC C:N ratio can have a legacy effect on soil CO₂ respiration in response to secondary low C:N inputs (i.e., starter N plus SD N) and reduce soil CO₂ respiration (Marschner et al., 2015). Deficit rainfall in May and June (i.e., 2016) may have prolonged cover crop decomposition and suggest consideration of CC C:N ratio management (e.g., termination dates) when using CC to stimulate soil microbial activity.

DNA sequence analysis

In year 1 (2014 – 2015) and year 2 (2015 – 2016) $7.1 - 9.2 \times 10^6$ valid sequences were obtained after quality filtering and represented a mean of 29,047 and 21,928 sequences per sample in each year, respectively. Despite soil sampling efforts each year rarefaction analysis did not produce a plateauing curve which indicated increased sampling frequency may have increased the number of recovered unique sequences. In spite of rarefaction results sequences were subsampled each year (i.e., 9,500=year 1 and 6,000=year 2) to help rarify data. Sequences covered 91,376 and 49,203 OTUs using 97% identity as the cutoff and suggests previous estimates (2,000 – 5,000 OTUs) are conservative (Schloss and Handelsman, 2006). In year 1 and year 2, 17,020 and 14,930 OTUs were represented by more than 5 reads which was 112 – 142% greater than reported in a previous study which utilized the Roche 454 platform to sequence soils treated with straw and N fertilizer (Pitombo et al., 2016). Sequencing identified 25 and 27 bacterial phyla in years 1 and 2 (24 phyla appearing across both years) and 682 and 655 genera, respectively. The top five phyla shared in each year represented 82 – 87% of the total OTUs identified. However, each year 42 – 44% of phyla and 76 – 78% of genera were unclassifiable and represented the largest group in each taxonomic rank. Collectively, singletons and doubletons (i.e., sequences observed 1x and 2x per sample, respectively) accounted for 73 and 56% of all sequences identified in years 1 and 2, respectively, and indicate rarity was common in the agricultural soils sampled.

Alpha diversity indicators (Inverse Simpson [ISD], Shannon Evenness [Shannon] and Chao1 Richness Estimator [Chao]) before and after corn establishment

Soil bacterial diversity (ISD), richness (Chao), and evenness (Shannon) was observed in whole plots prior to corn planting and in subplots of the zero-N strategy at R1 and R6. Data will be

discussed pertaining to whole plots first (not under the influence of corn) for year 1 (2014 – 2015) and year 2 (2015 – 2016), and then subplots for year 1 and year 2.

Before corn establishment: Estimation of OTU diversity using the inverse Simpson index (ISD), which accounts for richness (i.e., the number of species per sample) and evenness (i.e., the relative abundance of species in a sample) of soil bacterial communities indicated an effect of CC in year 1 (i.e., autumn, 2014, and May, 2015) (Table 9). Despite the abundance of singletons (previously mentioned) which can increase richness of samples, the ISD helps control for rarity and gives more weight to abundant species where OTU diversity increases as ISD increases. The radish CC reduced diversity 50 – 52% relative to oats or the no cover control which were similar in the autumn. Members of Brassicaceae (i.e., radish) contain quantities of glucosinolates that break down to isothiocyanates which may have biofumigant effects on soil microbial taxa and help explain reduced soil bacterial diversity to an ensuing corn crop observed in other studies (Kirkegaard and Sarwar, 1998; Fernandez et al., 2016; Tao et al., 2017). Results contrast the common assertion that increased plant diversity increases soil biological diversity. Instead radishes may have favored growth of copiotrophic taxa and decreased bacterial richness and evenness in the short term (Fernandez et al., 2016). Despite no sig. differences a trend for reduced OTU richness and evenness was evident which supports the previous hypothesis. In May (2015) radishes reduced diversity 23% relative to the oats while diversity in the no cover control remained similar to CCs which suggested that overwintering radish residues were continuing to affect soil bacterial membership more than the oat CC but to a lesser extent as CC decay progressed.

In year 2 CC did not affect diversity in the autumn (2015) or the following May (2016). Pearson correlations indicated that diversity in the May of corn establishment negatively corresponded to soil NO₃-N levels in both years ($r=-0.71$; $P \leq 0.01$), similar to other studies (Zhao et al., 2014; Li et al., 2016). Differing C:N ratios of radish CC biomass each autumn suggest differences in decomposition rates that could affect substrate availability as residue was broken down by soil microbiota. A 23:1 C:N ratio in year 2 (relative to 17:1 in year 1) may have reduced soil bacterial substrate availability and placed less emphasis on copiotrophic behavior (Koch, 2001; Fernandez et al., 2016). In May radish and oat CCs increased richness 7 – 11% and reduced evenness 0.9 – 1.0% which considering both factors may be an indication of increased number of rare OTUs. Within the two – year timeframe of this study it was unclear if CCs have a consistent effect on soil bacterial diversity, richness, and evenness prior to corn planting and warrants further investigation (Fernandez et al., 2016).

After corn establishment: When soil sample time and zone were included in the statistical model ANOVA of alpha diversity indicators did not indicate significant interactions of CC and N strategy when analyzed within soil sample zone, timing, or both and suggested soil bacterial communities responded similarly to N management strategies despite previous CCs (Table 10). Main effects of CC and N strategy are discussed separately. Soil bacterial diversity, richness, and evenness were not affected by CC while N strategy effects were significant in 4 of 6 observations which suggests soil bacterial communities were affected by N fertilization. Soil sampling zone and timing were significant in 5 of 6 observations which suggest that spatial and temporal variation had more impact on alpha diversity indicators than applied treatments.

In 2015 the PL plus V11 sidedress (SD) strategy increased ISD 7.2 – 9.4% relative to other N fertilizer strategies. The PL plus V11 SD treatment contained a natural organic plus inorganic NPK in addition to C, sulfur (S), Mg, Ca, boron (B), copper (Cu), iron (Fe), manganese (Mn), and zinc (Zn) provided by PL. Other studies have also observed increased soil bacterial diversity due to organic plus inorganic fertilization which may provide C and N substrates for bacterial function (Zhao et al., 2014). All N sources have the capacity to reduce soil pH which can indirectly due to release of H⁺ during corn root NO₃-N uptake, but urea and UAN-N sources may contribute directly to soil acidity due to 2H⁺ as a byproduct during decomposition (Havlin et al., 2014). Soil pH can be a determinant of soil bacterial diversity (Zhalnina et al., 2015). Urea and UAN fertilizers utilized in the current study may have created zones of acidity and reduced soil pH and reduced ISD relative to a natural organic N + inorganic N source (i.e., PL and V11 SD) although no differences were observed relative to the control. Richness was not affected within CCs and illustrates the dominance of spatial and temporal variation on richness. Cover crop residues had less influence to soil bacterial communities than N strategies as measured by ISD, chao richness, and Shannon evenness when combined across R1 and R6 sample times but growers may supplement a natural organic N source in their fertilizer program to increase diversity.

In 2015 a CC x soil sample zone x sample timing interaction was observed for diversity (i.e., ISD) (P=0.0091) and richness (P=0.0431) (Table 10). Slicing the ISD interaction indicated no diversity differences in soil sampling zone x timing within the radish CC only and suggests radish residues were able to reduce the spatial and temporal variation of soil bacterial diversity relative to oat and no CC soils (data not shown). Radish and oat CCs increased diversity 11 –

15% relative to no cover in bulk soils at R6 and not R1 and suggests 1) an actively growing cash crop may select for bacterial communities and reduce the influence of CC residues, 2) CC residues affecting diversity at corn maturity indicate recalcitrant residues may still influence soil bacterial diversity 11 months after termination, and 3) reduced influence of a corn rhizosphere after reaching physiological maturity. Studies observing no relationship of plant species richness and bacterial diversity have suggested that rhizosphere bacterial community composition may depend on the composition of bulk soil communities (de Ridder-Duine et al., 2005) or due to the functional redundancy of soil bacterial communities to utilize common root exudates (Zhalnina et al., 2015). Slicing the CC x soil sample zone x sample timing interaction for richness indicated CC did not have an effect on soil bacterial community richness regardless of soil sample timing or zone. Data suggest that future studies investigating CC influence to bacterial diversity in an actively growing corn cash crop may consider actively growing CC during the corn cash crop season due to the potentially interacting effects of plant rhizospheres and surrounding bulk soil bacterial communities.

In 2016 soils were dry following N applications in May and June due to deficit monthly rainfall. Cover crops did not affect ISD, Chao, or Shannon but each were affected by N strategy and suggests N fertilization may have more influence on bacterial OTUs in dry soils than wet (i.e., 2015) (Table 10). Limited rainfall to assimilate N fertilizer into the soil matrix may have increased soil solution osmotic potential, become toxic due to V4 UAN SD, and reduced community diversity 6 – 8% from soils receiving urea or PL PPI or no zero N and evenness 1.5% from the zero N strategy (Li et al., 2016). Strategies including V4 SD were coulter injected which may have preserved fertilizer band prevalence to reduce diversity relative to surface

banded V11 SD strategies. Pearson correlations indicated soil moisture was positively associated with bacterial diversity and evenness and negatively with soil inorganic N levels (Table 5). Associations suggest cumulative rainfall near N application timing may influence the effect of N strategy on soil bacterial communities. Similar to 2015, the PL and V11 SD resulted in similar diversity, richness, and evenness as the zero N strategy and suggest that an inorganic + organic N source may promote bacterial community stability across space and time. In both years diversity and evenness were increased at R6 relative to R1. Bulk soil diversity was increased in 2016 relative to the rhizosphere. Previous studies have suggested that environmental heterogeneity (e.g., pH and soil moisture) and field location interact to impact corn rhizosphere soil bacterial diversity (Pieffer et al., 2013). Deficit rainfall in May and June may have limited corn growth as evidenced by reduced yields in 2016 and reduced the influence corn had on bacterial diversity in dry spring soils as bulk soil diversity was similar each year (ISD 342 – 343).

In 2016 a N x Z x T interaction was observed for ISD ($P < 0.0001$) and Shannon evenness ($P = 0.0159$) (Table 10). Inspection of the interaction for ISD values indicated significant differences at R1 between strategies receiving N fertilizer in the bulk soil, but not the rhizosphere. In contrast Shannon evenness at R1 was affected by N strategies in the rhizosphere but not in the bulk soil. Other studies have illustrated the selection pressure plant rhizospheres can have on soil bacterial communities (Kowalchuk et al., 2002). Despite the influence of N strategies on bulk soil diversity no differences in ISD observed due to strategies receiving N fertilizer within R1 corn rhizosphere soils indicate corn influenced ISD more than N placement x timing. In lieu of ISD values, differences in Shannon evenness values in R1 rhizosphere soils indicate specific communities were influenced by the corn rhizosphere in regard to N fertilizer

strategy and suggest that N fertilizer strategy may have influenced corn selection of rhizosphere soil bacterial membership.

Pearson correlations indicated that R1 diversity in the rhizosphere was neg. associated with corn grain yield each year where N fertilizer had been applied ($r=-0.21$ to -0.44 ; $P \leq 0.08$) or had not been applied ($r=-0.53$ to -0.63 ; $P \leq 0.08$). Negative correlations suggest that relative abundance may explain more variation in grain yield. Reduced diversity were sometimes associated with treatments receiving UAN SD at V4. Evenness was negatively correlated with yield in the bulk soil ($r=-0.25$ to -0.36 ; $P \leq 0.05$) which suggests OTU relative abundance was affected and further supports reduced diversity associated with increased grain yield. Studies often do not report associations of crop yield with soil bacterial diversity, or have observed no correlation of plant biomass yield with microbial diversity (Zhalnina et al., 2015) and suggested bacterial abundance as indicators of crop yield (Tautges et al., 2015). While data indicating reduced R1 corn rhizosphere diversity and increased crop yield were consistent in both years reduced grain yield in a dry year (i.e., 2016) suggest future studies investigate variation of this effect across landscapes and climates for improved understanding on cropping system and soil bacterial community resilience to stress.

Conclusions

Two years of data demonstrated that CCs preceding a corn cash crop were able to affect soil labile amino acid N, soil CO₂ respiration, soil P and Ca levels, and soil bacterial diversity (i.e., ISD), richness (i.e., chao), and evenness (i.e., Shannon) but varied temporally. After corn cash crop seeding decomposing CC residues continued to affect soil respiration when cumulative May

– June rainfall was excessive (128 mm above normal) while under deficit May – June cumulative rainfall N fertilizer placement x timing treatments preceded by a radish CC affected CO₂ respiration. Daikon radish increased autumn biomass production and N uptake relative an oat CC, but less residue remaining in March of the following year suggesting fewer opportunities to affect soil microbial diversity in the ensuing corn cash crop season. During the corn cash crop season CCs had less influence on soil bacterial diversity, richness, and evenness in relation to the effects observed due to N strategies while more instances (5 of 6 site years) of interacting factors of soil sampling time and soil sampling zone (i.e., rhizosphere or bulk soils) affected soil bacterial diversity, richness, and evenness and suggest spatial and temporal variation were a large determinant of bacterial community membership. Nitrogen fertilizer which included a natural organic N source (i.e., poultry litter) increased diversity in relation to strategies that included V4 SD. Full V4 SD reduced diversity 6 – 8% from soils receiving urea, PL, or zero N in dry soils. An actively growing corn cash crop reduced the impact of N strategies on R1 rhizosphere soil bacterial diversity in relation to differences observed in the bulk soil. Grain yield was inversely proportional to soil bacterial diversity at R1 ($r=-0.21$ to -0.63 ; $P \leq 0.08$) regardless of fertilizer N application and indicates corn tasseling may be a time in corn ontogeny to increase opportunities to affect grain yield but suggests relative abundance of bacterial community membership may explain more variation in grain yield than diversity. However reduced grain yields in dry soils due to an oat CC suggest growers may need to select CC species according to agronomic objectives. Two site years of data with variable May and June rainfall suggest the main driver of soil microbial diversity, richness, and evenness during a corn cash crop season is N fertilizer strategy and not previous CC, but it remains unclear due to spatial and temporal variation between years. Relationship of alpha diversity indicators to corn grain yields and

cropping system resilience may depend on multi-year observations of crop rotations where the rotation itself is replicated across site years. A more extensive soil sampling and sequencing effort may be necessary to characterize relationships occurring between bacterial OTUs and corn rhizospheres. Future studies may be enhanced with analysis of microbial functional groups and not taxonomic relatedness.

Acknowledgements

The authors would like to thank the USDA National Institute of Food and Agriculture, Michigan AgBioResearch, the Michigan State University's College of Agriculture and Natural Resources, and Corn Marketing and Promotion of Michigan for partial funding and support of this research.

APPENDIX

Table 4.01. Monthly precipitation and temperature departures from normal and 30-yr means for Lansing, MI, 2014 – 2016.

Month	precipitation [†]				air temperature [‡]			
	2014	2015	2016	30-yr mean	2014	2015	2016	30-yr mean
	difference (mm) from avg.			mm	difference (°C) from avg.			°C
Aug.	+2	+39	+79	84	-1.0	-1.0	+1.6	21.3
Sept.	-7	+3	+14	92	-1.5	+2.1	+0.5	16.9
Oct.	-13	-22	+12	70	-1.3	+0.7	-1.2	10.4
Nov.	-34	-40	-22	71	-3.7	+2.3	+0.1	4.6
Apr.	-51	-50	+2	73	-0.7	-0.5	-1.3	8.7
May	-2	+25	-32	85	-0.3	+1.6	0.0	14.7
June	+34	+103	-71	89	0.0	-0.8	+0.3	20.0
July	-22	-21	+13	83	-3.3	-1.2	+1.0	22.1

[†]Precipitation and air temperature data were collected from Michigan Automated Weather Network (<http://www.agweather.geo.msu.edu/mawn/>).

[‡]30-yr mean sources for precipitation and air temperatures: NOAA (<https://www.ncdc.noaa.gov/cdo-web/datatools/normals>; verified 17 Aug. 2017).

Table 4.02. Two-year soil moisture (30-cm), precipitation (cm), and soil temperature (5-cm) data for the 14 d period prior to each soil sampling event (cover crop planting [CC plant], autumn cover crop harvest [CC harvest], corn planting [corn plant], corn tasseling [corn R1], and corn harvest [corn R6]) in Lansing, MI, 2014 – 2016.

	CC plant [†]	CC harvest	Corn plant	Corn R1	Corn R6
<u>2014-15</u>	14-Aug	3-Nov	1-May	20-Jul	9-Oct
Max. soil moist. (cm ³ cm ⁻³)	0.110	0.211	0.197	0.273	0.174
Min. soil moist. (cm ³ cm ⁻³)	0.110	0.202	0.190	0.257	0.170
Precip. (mm)	17.0	17.8	7.1	46.2	14.0
Max. soil temp. (°C)	24.0	4.0	12.0	27.0	16.5
Min. soil temp. (°C)	18.4	2.5	6.3	16.5	13.2
<u>2015-16</u>	17-Aug	30-Oct	17-May	20-Jul	11-Oct
Max. soil moist. (cm ³ cm ⁻³)	0.200	0.232	0.250	0.085	0.237
Min. soil moist. (cm ³ cm ⁻³)	0.179	0.224	0.235	0.079	0.222
Precip. (mm)	65.3	32.5	37.3	42.2	80.8
Max. soil temp. (°C)	22.9	11.4	16.5	30.0	19.3
Min. soil temp. (°C)	18.9	8.1	10.5	21.4	14.6

[†]Cover crop planting and harvest relative to 2014 and 2015, respectively. Corn planting, R1, and R6 relative to 2015 and 2016, respectively.

Table 4.03. Cover crop biomass and quality at autumn cover crop harvest, Lansing, MI, 2014 - 2015.

Sample	Biomass		C uptake		N uptake		C/N	
	2014							
			kg ha ⁻¹				mg kg ⁻¹	
Radish†	10567	a‡	3866	a	230	a	17:1	b
Oats	7604	b	3149	a	141	b	23:1	a
<i>P</i> > <i>F</i>	0.0667		0.1651		0.0550		0.0724	
2015								
Radish	9351	a	3298	a	149	a	23:1	a
Oats	4255	b	1765	b	101	b	17:1	b
<i>P</i> > <i>F</i>	0.0030		0.0072		0.0065		0.0508	

†Radish biomass unable to be sampled in spring due to winter decomposition.

‡Means followed by the same letter in a column within each year are not sig. different at alpha=0.10.

Table 4.04. Main effects of cover crop and N strategy[†] on corn grain yield, Lansing, MI, 2015 – 2016.

Main effect	Grain Yield			
	2015	2016		
Cover crop		Mg ha ⁻¹		
No cover	15.8	a [‡]	13.6	a
Radish	15.4	a	13.5	a
Oat	15.5	a	13.0	b
N Strategy				
Zero N	12.4	c	10.8	b
Urea [PPI]	15.9	b	14.0	a
PL [PPI] + V11	16.4	a	13.8	a
5x5 starter + V4	16.2	ab	13.7	a
5x5 starter + V11	16.4	a	13.7	a
5x5 starter + V4/V11	16.3	ab	13.9	a
ANOVA				
Cover (C)	0.1309		0.0311	
N strategy (N)	<.0001		<.0001	
C x N	0.2262		0.3271	

[†]All strategies requiring N fertilizer received total maximum return to N rate of 179 kg N ha⁻¹ in all years.

[‡]Means within each column followed by same lower case letters are not significantly different at alpha=0.10.

Table 4.05. Correlation coefficients (r) ($n=10$) relating indicators of soil microbial alpha diversity[†] indicators (inverse Simpson diversity [ISD], Chao1 richness [Chao] and Shannon evenness [Shannon]) or soil parameters[‡] (soil labile amino acid [SLAN], CO₂ respiration, soil NO₃-N and NH₄-N) to weather factors observed during the 14 d period prior to cover crop planting, autumn cover crop harvest, corn planting, corn tasseling, and corn harvest in Lansing, MI, in 2015 - 2016.

	ISD	Chao	Shannon	SLAN	Respiration	NO ₃ ⁻	NH ₄ ⁺
Max. Soil Moist.	0.81 ^{**}	0.02	0.55 ^{0.10}	-0.26	-0.12	-0.62 ^{0.06}	-0.78 ^{**}
Min. Soil Moist.	0.82 ^{**}	0.06	0.54	-0.28	-0.17	-0.64 [*]	-0.79 ^{**}
Precipitation	0.18	-0.28	0.25	0.25	0.38	-0.19	0.03
Max. Air Temp.	-0.52	-0.11	-0.29	0.27	0.39	0.35	0.60 ^{0.07}
Min. Air Temp.	-0.48	-0.05	-0.30	0.32	0.52	0.40	0.63 [*]

^{*}, ^{**}, ^{***} Significant at the 0.05, 0.01, and 0.001 probability levels, respectively.

[†]Soil parameters were an average of means observed across 1) replications at cover planting 2) whole plot replications in the autumn and spring, and 3) subplot replications within the zero-N strategy at corn R1 and R6.

[‡]Soil alpha diversity indicators were an average of means observed across 1) replications at cover planting 2) whole plot replications in the autumn and spring, and 3) subplot replications combined across soil sampling zones at corn R1 and R6.

Table 4.06. Effect of cover crop and no N fertilizer on soil labile amino N (SLAN) and soil respiration (CO₂) means observed in the autumn and the ensuing May at corn planting, Lansing, MI, 2014 - 2016.

Treatment	2014 – 2015							
	Autumn				Spring			
	SLAN [†]		CO ₂ [‡]		SLAN		CO ₂	
	kg N ha ⁻¹	mg CO ₂ kg ⁻¹			kg N ha ⁻¹	mg CO ₂ kg ⁻¹		
No Cover	115	b [§]	15.7	a	108	a	16.2	a
Radish	132	a	19.8	a	129	a	19.9	a
Oats	125	ab	17.8	a	118	a	19.5	a
P > F	.0716		0.5054		0.1369		0.5190	
	2015 – 2016 [¶]							
No Cover	157	b	14.7	b	142	a	15.5	b
Radish	152	b	23.6	a	155	a	24.1	a
Oats	179	a	21.5	a	158	a	22.0	a
P > F	0.0973		0.0896		0.5037		0.0358	

[†]Mean SLAN was 142 and 150 kg N ha⁻¹ at cover crop planting in 2014 and 2015, respectively.

[‡]Basal CO₂ respiration was 35.6 and 35.9 mg kg⁻¹ at cover crop planting in 2014 and 2015, respectively.

[§]Means followed by the same letter in a column within each year not sig. different at alpha=0.10.

[¶]Field soil moisture content at sampling time was 1.8 and 1.7 x 10⁻⁷ mg kg⁻¹ in the autumn of 2015 and May 2016, respectively. Moistures not presented for autumn of 2014 and spring of 2015 as a dry method sampling protocol was used.

Table 4.07. Main effects of cover crop and N strategy[†] on basal soil respiration combined across observations at flowering (R1) and physiological maturity (R6), Lansing, MI, 2015.

Main effect	Respiration	
Cover crop	mg kg ⁻¹	
No cover	18.1	b [‡]
Radish	20.2	a
Oat	20.9	a
N Strategy		
Zero N	19.8	a
Urea [PPI]	19.8	a
PL [PPI] + V11	20.0	a
5x5 starter + V4	18.5	a
5x5 starter + V11	19.9	a
5x5 starter + V4/V11	20.6	a
Timing		
Corn tasseling (R1)	21.6	a
Corn maturity (R6)	17.9	b
	ANOVA	
Cover (C)	0.0408	
N strategy (N)	0.7192	
C x N	0.2723	
Timing (T)	0.0089	
C x T	0.3177	
N x T	0.7337	
C x N x T	0.8163	

[†]All strategies requiring N fertilizer received total maximum return to N rate of 179 kg N ha⁻¹ in all years.

[§]Means within each column followed by same lower case letters are not significantly different at alpha=0.10.

Table 4.08. Cover crop and N management interaction ($P=0.0374$) on soil basal CO₂ respiration combined across R1 and R6 sampling times, Lansing, MI[†] 2016.

N strategy	Cover					
	No Cover		Radish		Oat	
	mg CO ₂ kg ⁻¹					
Zero N	21.6	a [‡]	21.1	abc	22.2	a
Urea [PPI]	19.3	a	24.6	a	18.0	a
PL [PPI] + V11	20.8	a	24.7	a	22.4	a
5x5 starter + V4	22.5	a	18.5	c	23.9	a
5x5 starter + V11	17.6	a	22.6	ab	23.4	a
5x5 starter + V4/V11	18.9	a	19.7	bc	22.4	a
$P > F$	0.3287		0.0597		0.2330	

[†]All strategies requiring N fertilizer received total maximum return to N rate of 179 kg N ha⁻¹ in all years.

[‡]Means within each column followed by same lower case letters are not significantly different at alpha=0.10.

Table 4.09. Effect of cover crop on alpha diversity indicators[†] (inverse Simpson [ISD], Chao1 Richness [Chao] and Shannon evenness [Shannon]) sampled in the autumn and spring prior to corn planting, Lansing, MI, 2014 – 2016.

Treatment	2014-15						
	Fall			Spring			
	ISD	Chao	Shannon	ISD	Chao	Shannon	
No Cover	392 a [‡]	5581 a	0.848 a	326 ab	3786 a	0.856 a	
Radish	190 b	3874 a	0.827 a	290 b	3011 a	0.866 a	
Oat	380 a	5821 a	0.857 a	379 a	4313 a	0.851 a	
<i>P</i> > <i>F</i>	0.0470	0.3670	0.1531	0.0695	0.1806	0.3319	
Treatment	2015-16 [§]						
	No Cover	339 a	4032 a	0.859 a	350 a	3924 b	0.864 a
	Radish	343 a	3828 a	0.864 a	340 a	4208 a	0.854 b
	Oat	367 a	4339 a	0.860 a	371 a	4362 a	0.856 b
	<i>P</i> > <i>F</i>	0.4458	0.4287	0.8873	0.1417	0.0305	0.0985

[†]Diversity indices were assessed for operational taxonomic units (OTUs) based on a 97% sequence identity.

[‡]Means followed by the same letter in a column within each year are not sig. different at alpha ≤ 0.10.

[§]2015 ISD normalized using exponential transformation. Untransformed means are presented.

Table 4.10. Main effects of cover crop and N strategy[†] on alpha diversity indicators (inverse Simpson diversity [ISD], Chao1 richness [Chao] and Shannon evenness [Shannon]) across soil sampling zones (rhizosphere and bulk soil) and sample timings (corn tasseling and harvest), Lansing, MI, 2015- 2016.

Treatment	2015			2016								
	ISD	Chao	Shannon	ISD	Chao	Shannon						
Cover Crop												
No cover	340	a [‡]	4525	a	0.851	a	324	a	3715	a	0.858	a
Radish	354	a	4546	a	0.856	a	320	a	3789	a	0.855	a
Oat	355	a	4436	a	0.853	a	333	a	3908	a	0.855	a
N Strategy												
Zero N	356	ab	4714	a	0.854	a	329	ab	3930	ab	0.862	a
Urea [PPI]	347	b	4527	a	0.853	a	336	a	3680	bc	0.862	a
PL [PPI] + V11	372	a	4608	a	0.855	a	331	a	4141	a	0.856	abc
5x5 starter + V4	341	b	4286	a	0.853	a	309	c	3686	bc	0.849	c
5x5 starter + V11	341	b	4451	a	0.854	a	329	ab	3797	bc	0.856	ab
5x5 starter + V4/V11	340	b	4427	a	0.850	a	319	bc	3589	c	0.853	bc
Zone												
Bulk soil	343	b	4382	b	0.852	a	342	a	3888	a	0.859	a
Corn rhizosphere	357	a	4623	a	0.855	a	310	b	3719	a	0.853	b
Sample time												
Corn tassel (R1)	342	b	4510	a	0.850	b	291	b	3861	a	0.843	b
Corn maturity (R6)	357	a	4495	a	0.856	a	361	a	3747	a	0.870	a
ANOVA ($P > F$)												
Cover (C)	0.3460	0.7785	0.3653	0.6247	0.6834	0.4827						
N strategy (N)	0.0242	0.5059	0.5827	0.0124	0.0970	0.0302						
C x N	0.8190	0.5083	0.4326	0.7487	0.4082	0.7117						
Zone (Z)	0.0112	0.0341	0.1048	<.0001	0.1543	0.0242						
C x Z	0.1921	0.2677	0.6214	0.6658	0.4137	0.2639						
N x Z	0.7730	0.7478	0.6892	0.2744	0.1833	0.4358						
C x N x Z	0.7675	0.6087	0.4783	0.7570	0.4796	0.4114						
Timing (T)	0.0238	0.9043	0.0005	<.0001	0.3286	<.0001						
C x T	0.4758	0.4015	0.9450	0.8872	0.7064	0.7028						
N x T	0.1981	0.6802	0.1492	<.0001	0.9340	0.0041						
C x N x T	0.9578	0.4999	0.5223	0.4173	0.8466	0.8244						
Z x T	0.0907	<.0001	<.0001	<.0001	<.0001	0.4922						
C x Z x T	0.0091	0.0413	0.9131	0.1669	0.5067	0.9546						
N x Z x T	0.3786	0.9459	0.9282	<.0001	0.7223	0.0159						
C x N x Z x T	0.8802	0.5765	0.6152	0.7955	0.9288	0.7904						

[†]All strategies requiring N fertilizer received total maximum return to N rate of 179 kg N ha⁻¹ in all years.

[‡]Means within each column followed by same lower case letters are not significantly different at alpha=0.10.

LITERATURE CITED

LITERATURE CITED

- Amos, B., and D.T. Walters. 2006. Maize root biomass and net rhizodeposited carbon: An analysis of the literature. *Soil Sci. Soc. Am. J.* 70:1489-1503.
- Awasthi, A., M. Singh, S.K. Soni, R. Singh, and A. Kalra. 2014. Biodiversity acts as insurance of productivity of bacterial communities under abiotic perturbations. *ISME J.* 8:2445-2452.
- Bainard, L.D., C. Hamel, and Y. Gan. 2016. Edaphic properties override the influence of crops on the composition of the soil bacterial community in a semiarid agroecosystem. *App. Soil Ecol.* 105:160-168.
- Berg, G., and K. Smalla. 2009. Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *FEMS Microbiol. Ecol.* 68:1-13.
- Binder, D.L., D.H. Sander, and D.T. Walters. 2000. Maize response to time of nitrogen application as affected by level of nitrogen deficiency. *Agron. J.* 92:1228-1236.
- Brookes, P. 2001. The soil microbial biomass: Concept, measurement and applications in soil ecosystem research. *Microbes and Environ.* 16:131-140.
- Buyer, J.S., J.R. Teasdale, D.P. Roberts, I.A. Zasada, and J.E. Maul. 2010. Factors affecting soil microbial community structure in tomato cropping systems. *Soil Biol. Biochem.* 42:831-841.
- Cadotte, M.W., J. Cavender-Bares, D. Tilman, and T.H. Oakley. 2009. Using phylogenetic, functional, and trait diversity to understand patterns of plant community productivity. *PLoS ONE* 4:e5695.
- Cavaglieri, L., J. Orlando, and M. Etcheverry. 2009. Rhizosphere microbial community structure at different maize plant growth stages and root locations. *Microbiol. Res.* 164:391-399.
- Ceja-Navarro, J.A., F.N. Rivera-Orduña, L. Patiño-Zúñiga, A. Vila-Sanjurjo, J. Crossa, B. Govaerts, and L. Dendooven. 2010. Phylogenetic and multivariate analyses to determine the effects of different tillage and residue management practices on soil bacterial communities. *App. Environ. Micro.* 76:3685-3691.
- Chaparro, J.M., A.M. Sheflin, D.K. Manter, and J.M. Vivanco. 2012. Manipulating the soil microbiome to increase soil health and plant fertility. *Biol. Fertil. Soils* 48:489-499.
- Cleveland, C.C., S.C. Reed, A.B. Keller, D.R. Nemergut, S.P. O’Niell, R. Ostertag, and P.M. Vitousek. 2014. Litter quality versus soil microbial community controls over decomposition: a quantitative analysis. *Oecologia* 174:283-294.
- Combs, S.M., and M.V. Nathan. 2015. Soil organic matter. In: M.V. Nathan and R. Gelderman, editors, *Recommended chemical soil test procedures for the North Central Region*. North

- Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 12.1-12.6.
- de Ridder-Duine, A.S., G.A. Kowalchuk, P.J.A.K. Gunnewiek, W. Smant, J.A. van Veen, and W. de Boer. 2005. Rhizosphere bacterial community composition in natural stands of *Carex arenaria* (sand sedge) is determined by bulk soil community composition. *Soil Biol. Biochem.* 37:349-357.
- Ding, W., H. Yu, Z. Cai, F. Han, and Z. Xu. 2010. Responses of soil respiration to N fertilization in a loamy soil under maize cultivation. *Geoderma* 155:381-389.
- Doran, J.W., and M.R. Zeiss. 2000. Soil health and sustainability: Managing the biotic component of soil quality. *Appl. Soil. Ecol.* 15:3-11.
- Eisenhauer, N., S. Scheu, and A. Jousset. 2012. Bacterial diversity stabilizes community productivity. *PLoS ONE* 7:e34517.
- Fernandez, A.L., C.C. Sheaffer, D.L. Wyse, C. Staley, T.J. Gould, and M.J. Sadowsky. 2016. Structure of bacterial communities in soil following cover crop and organic fertilizer incorporation. *Appl. Microbiol. Biotechnol.* 100:9331-9341.
- Fierer, N., C.L. Lauber, K.S. Ramirez, J. Zaneveld, M.A. Bradford, and R. Knight. 2012. Comparative metagenomic, phylogenetic and physiological analyses of soil microbial communities across nitrogen gradients. *The ISME J.* 6:1007-1017.
- Figuerola, E.L.M., L.D. Guerrero, D. Türkowsky, L.G. Wall, and L. Erijman. 2015. Crop monoculture rather than agriculture reduces the spatial turnover of soil bacterial communities at a regional scale. *Environ. Micro.* 17:678-688.
- Frank, K., D. Beegle, and J. Denning. 2015. Phosphorus. In: M.V. Nathan and R. Gelderman, editors, Recommended chemical soil test procedures for the North Central Region. North Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 6.1-6.6.
- Geisseler, D., and K.M. Scow. 2014. Does long-term use of mineral fertilizers affect the soil microbial biomass? *Better Crops* 98:13-15.
- Han, G.X., G.S. Zhou, Z.Z. Xu, Y. Yang, J.L. Liu, and K.Q. Shi. 2007. Biotic and abiotic factors controlling the spatial and temporal variation of soil respiration in an agricultural ecosystem. *Soil Biol. and Biochem.* 39:418-425.
- Haney, R.L., W.F. Brinton, and E. Evans. 2008. Soil CO₂ respiration: Comparison of chemical titration, CO₂ IRGA analysis and the Solvita gel system. *Renew. Agric. Food Syst.* 23:171-176.

- Havlin, J.L., S.L. Tisdale, W.L. Nelson, and J.D. Beaton. 2014. *Soil Fertility and Fertilizers. An Introduction to Nutrient Management*. 8th ed. Pearson Education Inc., Upper Saddle River, NJ.
- Huang, X.F., J.M. Chaparro, K.F. Reardon, R. Zhang, Q. Shen, and J.M. Vivanco. 2014. Rhizosphere interactions: root exudates, microbes, and microbial communities. *Botany* 92:267-275.
- Huffman, S.A. and K.A. Barbarick. 1981. Soil nitrate analysis by cadmium reduction. *Communications in Soil Science and Plant Analysis* 12:79-89.
- Ikeda, S., K. Suzuki, M. Kawahara, M. Noshiro, and N. Takahashi. 2014. An assessment of urea-formaldehyde fertilizer on the diversity of bacterial communities in onion and sugar beet. *Microbes Environ.* 29:231-234.
- Jahanzad, E., A.V. Barker, M. Hashemi, T. Eaton, A. Sadeghpour, and S.A. Weis. 2016. Nitrogen release dynamics and decomposition of buried and surface cover crop residues. *Agron. J.* 108:1735-1741.
- Jan, M.T., P. Roberts, S.K. Tonheim, and D.L. Jones. 2009. Protein breakdown represents a major bottleneck in nitrogen cycling in grassland soils. *Soil Biol. Biochem.* 41:2272-2282.
- Janda, J.M., and S.L. Abbott. 2007. 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: Pluses, perils, and pitfalls. *J. Clinical Microbiol.* 45:2761-4.
- Kibblewhite, M.G., K. Ritz, and M.J. Swift. 2008. Soil health in agricultural systems. *Phil. Trans. R. Soc. B* 363:685-701.
- Kirkegaard, J.A., and M. Sarwar. 1998. Biomfumigation potential of brassicas I. Variation in glucosinolate profiles of diverse field-grown brassicas. *Plant and Soil* 201:71-89.
- Koch, A.L. 2001. Oligotrophs versus copiotrophs. *Bioessays* 23:657-661.
- Kowalchuk, G.A., D.S. Buma, W. de Boer, P.G.L. Klinkhamer, and J.A van Veen. 2002. Effects of above-ground plant species composition and diversity on the diversity of soil-borne microorganisms. *Antonie van Leeuwenhoek* 81:509-520.
- Kozich, J.J., S.L. Westcott, N.T. Baxter, S.K. Highlander, and P.D. Schloss. 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform. *App. Environ. Micro.* 79:5112-5120.
- Lehman, R.M., V. Acosta-Martinez, J.S. Buyer, C.A. Cambardella, H.P. Collins, T.F. Ducey, J.J. Halvorson, V.L. Jin, J.M.F. Johnson, R.J. Kremer, J.G. Lundgren, D.K. Manter, J.E. Maul, J.L. Smith, and D.E. Stott. 2015. Soil biology for resilient, healthy soil. *J. Soil Water Conserv.* 70:12A-18A.

- Li, Q., X. Song, H. Gu, and F. Gao. 2016. Nitrogen deposition and management practices increase soil microbial biomass carbon but decrease diversity in Moso bamboo plantations. *Scientific Reports* 6:1-11.
- Li, X., J. Rui, J. Xiong, J. Li, Z. He, J. Zhou, A.C. Yannarell, and R.I. Mackie. 2014. Functional potential of soil microbial communities in the maize rhizosphere. *PLoS ONE* 9:e112609.
- Li, X., J. Rui, Y. Mao, A. Yannarell, and R. Mackie. 2014. Dynamics of the bacterial community structure in the rhizosphere of a maize cultivar. *Soil Biol. Biochem.* 68:392-401.
- Lupwayi, N.Z., W.A. Rice, and G.W. Clayton. 1998. Soil microbial diversity and community structure under wheat as influenced by tillage and crop rotation. *Soil Biol. Biochem.* 30:1733-41.
- Lupwayi, N.Z., G.P. Lafond, N. Ziadi, and C.A. Grant. 2012. Soil microbial response to nitrogen fertilizer and tillage in barley and corn. *Soil Till. Res.* 118:139-146.
- Marschner, P., Z. Hatam, and T.R. Cavagnaro. 2015. Soil respiration, microbial biomass and nutrient availability after the second amendment are influenced by legacy effects of prior residue addition. *Soil Biol. Biochem.* 88:169-177.
- Mishra, V., and K.A. Cherkauer. 2010. Retrospective droughts in the crop growing season: Implications to corn and soybean yield in the Midwestern United States. *Agric. Forest Meteor.* 150:1030-1045.
- Mooshammer, M., W. Wanek, I. Hämmerle, L. Fuchslueger, F. Hafhansl, A. Knoltsch, J. Schnecker, M. Takriti, M. Watzka, B. Wild, K.M. Keiblinger, S. Zechmeister-Boltenstern, and A. Richter. 2014. Adjustment of microbial nitrogen use efficiency to carbon:nitrogen imbalances regulates soil nitrogen cycling. *Nat. Commun.* 5:3694.
- Naseem, H., and A. Bano. 2014. Role of plant growth-promoting rhizobacteria and their exopolysaccharide in drought tolerance of maize. *J. Plant Inter.* 9:689-701.
- Nielsen, S., T. Minchin, S. Kimber, L. van Zwieten, J. Gilbert, P. Munroe, S. Joseph, and T. Thomas. 2014. Comparative analysis of the microbial communities in agricultural soil amended with enhanced biochars or traditional fertilisers. *Agric. Ecosys. Environ.* 191:73-82.
- Nihorimbere, V., M. Ongena, M. Smargiassi, and P. Thonart. 2011. Beneficial effect of the rhizosphere microbial community for plant growth and health. *Biotechnol. Agron. Soc. Environ.* 15:327-337.
- Orr, C.H., C. Leifert, S.P. Cummings, and J.M. Cooper. 2012. Impacts of organic and conventional crop management on diversity and activity of free-living nitrogen fixing bacteria and total bacteria are subsidiary to temporal effects. *PLoS ONE* 7:e52891.

- Otani, T., N. Ae, and H. Tanaka. 1996. Phosphorus (P) uptake mechanisms of crops grown in soils with low P status. *Soil Sci. Plant Nutr.* 42:553-560.
- Paterson, E., G. Osler, L.A. Dawson, T. Gebbing, A. Sim, and B. Ord. 2008. Labile and recalcitrant plant fractions are utilized by distinct microbial communities in soil: Independent of the presence of roots and mycorrhizal fungi. *Soil Biol. Biochem.* 40:1103-1113.
- Peiffer, J.A., A. Spor, O. Koren, Z. Jin, S.G. Tringe, J.L. Dangl, E.S. Buckler, and R.E. Ley. 2013. Diversity and heritability of the maize rhizosphere microbiome under field conditions. *PNAS* 110:6548-6553.
- Peters, J.B., M.V. Nathan, and C.A.M. Laboski. 2015. pH and lime requirement. In: M.V. Nathan and R. Gelderman, editors, *Recommended chemical soil test procedures for the North Central Region*. North Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 4.1-4.7.
- Pitombo, L.M., J.B. Do Carmo, M. De Hollander, R. Rossetto, M.V. López, H. Cantarella, and E.E. Kuramae. 2016. Exploring soil microbial 16S rRNA sequence data to increase carbon yield and nitrogen efficiency of a bioenergy crop. *GCB Bioenergy* 8:867-879.
- Raich, J.W., and G. Mora. 2005. Estimating root plus rhizosphere contributions to soil respiration in annual croplands. *Soil Sci. Soc. Am. J.* 69:634-639.
- Rosenzweig, N., 2014. The importance and application of bacterial diversity in sustainable agricultural crop production ecosystems. In *Bacterial Diversity in Sustainable Agriculture* (pp. 341-367). Springer International Publishing.
- Rosenzweig, N., J.M. Bradeen, Z.J. Tu, S.J. McKay, and L.L. Kinkel. 2013. Rhizosphere bacterial communities associated with long-lived perennial prairie plants vary in diversity, composition, and structure. *Can. J. Microbiol.* 59:494-502.
- Rosenzweig, N, L. Steere, B. Gerondale, and W.W. Kirk. 2017. A geostatistical approach to visualize the diversity of soil inhabiting bacteria and edaphic qualities in potato (*Solanum tuberosum*) production systems. *Am. J. Potato Res.* 93:518-532.
- SAS Institute. 2011. *The SAS system for windows*. Version 9.4. SAS Inst., Cary, NC.
- Sanguin, H., B. Remenant, A. Dechesne, J. Thioulouse, T.M. Vogel, X. Nesme, Y. Moënne-Loccoz, and G.L. Grundmann. 2006. Potential of a 16S rRNA-based taxonomic microarray for analyzing the rhizosphere effects of maize on *Agrobacterium* spp. and bacterial communities. *App. Environ. Microbiol.* 72:4302-4312.
- Sagova-Marechova, M., M. Omelka, L. Cermak, Z. Kamenik, J. Olsovska, E. Hackl, J. Kopechy, and F. Hadacek. 2011. Microbial communities show parallels at sites with distinct litter and soil characteristics. *Appl. Environ. Microbiol.* 77:7560-7567.
- Schloss, P.D., and J. Handelsman. 2006. Toward a census of bacteria in soil. *PLoS ONE* 2:e92.

- Schloss, P.D., S.L. Westcott, T. Ryabin, J.R. Hall, M. Hartmann, E.B. Hollister, R.A. Lesniewski, B.B. Oakley, D.H. Parks, C.J. Robinson, J.W. Sahl, B. Stres, G.G. Thallinger, D.J. Van Horn, and C.F. Weber. 2009. Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Appl. Environ. Microbiol.* 2009. 75:7537-41.
- Schultze, S.R., P. Sabbatini, and L. Luo. 2016. Effects of a warming trend on cool climate viticulture in Michigan, USA. *SpringerPlus* 5:1119.
- Sharma, S.K., A. Ramesh, M.P. Sharma, O.P. Joshi, B. Govaerts, K.L. Steenwerth, and D.L. Karlen. 2010. Microbial community structure and diversity as indicators for evaluating soil quality. p. 317-358. *In* E. Lichtfouse (ed.) *Biodiversity, biofuels, agroforestry and conservation agriculture. Sustainable Agriculture Reviews. Vol. 5.* Springer Netherlands, 2011.
- Smith, C.R., P.L. Blair, C. Boyd, B. Cody, A. Hazel, A. Hedrick, H. Kathuria, P. Khurana, B. Kramer, K. Muterspaw, C. Peck, E. Sells, J. Skinner, C. Tegeler, and Z. Wolfe. 2016. Microbial community response to soil tillage and crop rotation in a corn/soybean agroecosystem. *Ecol. Evol.* 6:8075-84.
- Soman, C. D. Li, M.M. Wander, and A.D. Kent. 2017. Long-term fertilizer and crop-rotation treatments differentially affect soil bacterial community structure. *Plant Soil* 413:145-159.
- Song, Z., H. Gao, P. Zhu, C. Peng, A. Deng, C. Zheng, M.A. Mannaf, M.N. Islam, and W. Zhang. 2015. Organic amendments increase corn yield by enhancing soil resilience to climate change. *The Crop J.* 3:110-117.
- Steinke, K. 2015. 2015 MRTN suggested N rates for corn. Michigan State University Ext. Online. Available at <https://soil.msu.edu/wp-content/uploads/2014/06/2015-MRTN-N-Rates-for-Corn.pdf> (posted 27 Mar. 2015; verified 05 Oct. 2017).
- Sugiura, Y., and K. Nomoto. 1984. Phytosiderophores structures and properties of mugineic acids and their metal complexes. *Structure and Bonding* 58:107-135.
- Sun, B., S. Jia, S. Zhang, N.B. McLaughlin, A. Liang, X. Chen, S. Liu, and X. Zhang. 2016. No tillage combined with crop rotation improves soil microbial community composition and metabolic activity. *Environ. Sci. Pollut. Res.* 23:6472-6482.
- Tao, J. X. Liu, Y. Liang, J. Niu, Y. Xiao, Y. Gu, L. Ma, D. Meng, Y. Zhang, W. Huang, D. Peng, and H. Yin. 2017. Maize growth responses to soil microbes and soil properties after fertilization with different green manures. *Appl. Microbiol. Biotechnol.* 101:1289-1299.
- Tautges, N.E., T.S. Sullivan, C.L. Reardon, and I.C. Burke. 2016. Soil microbial diversity and activity linked to crop yield and quality in a dryland organic wheat production system. *App. Soil Ecol.* 108:258-268.

- United States Department of Agriculture-National Agriculture Statistic Service (USDA- NASS) (2017). Quick stats. Online. Accessed 04 Dec. 2017. <https://quickstats.nass.usda.gov/>. U.S. Environmental Protection Agency. 2016. What climate change means for Michigan. Report No. EPA 430-F-16-024.
- Venter, Z.S., K. Jacobs, and H.J. Hawkins. 2016. The impact of crop rotation on soil microbial diversity: A meta-analysis. *Pedobiologia* 59:215-223.
- Verzeaux, J., A. Alahmad, H. Habbib, E. Nivelles, D. Roger, J. Lacoux, G. Decocq, B. Hirel, M. Catterou, F. Spicher, F. Dubois, J. Duclercq, and T. Tetu. 2016. Cover crops prevent the deleterious effect of nitrogen fertilization on bacterial diversity by maintaining the carbon content of ploughed soil. *Geoderma* 281:49-57.
- Vukicevich, E., T. Lowery, P. Bowen, and J.R. Úrbez-Torres. 2017. Cover crops to increase soil microbial diversity and mitigate decline in perennial agriculture. A review. *Agron. Sustain. Dev.* 36:1-14.
- Vurukonda, S.S.K.P., S. Vardharajula, M. Shrivastava, and A. Skz. 2016. Enhancement of drought stress tolerance in crops by plant growth promoting rhizobacteria. *Micro. Res.* 184:13-24.
- Wang, G., Ngouajio, M., and Warncke, D.D. 2008. Nutrient cycling, weed suppression, and onion yield following brassica and sorghum sudangrass cover crops. *HortTechnology* 18:68-74.
- Warncke, D., and J.R. Brown. 2015. Potassium and other basic cations. In: M.V. Nathan and R. Gelderman, editors, *Recommended chemical soil test procedures for the North Central Region*. North Central Region Res. Publ. 221 (rev.). SB 1001. Missouri Agric. Exp. Stn., Columbia. p. 7.1-7.3.
- Ye, J., P.G. Perez, R. Zhang, S. Nielsen, D. Huang, and T. Thomas. 2017. Effects of different C/N ratios on bacterial compositions and processes in an organically managed soil. [Online]. *Biol. Fertil. Soils* <https://doi.org/10.1007/s00374-017-1246-5>
- Zhalnina, K., R. Dias, P.D. de Quadros, A. Davis-Richardson, F.A.O. Camargo, I.M. Clark, S.P. McGrath, P.R. Hirsch, and E.W. Triplett. 2015. Soil pH determines microbial diversity and composition in the Park Grass Experiment. *Microb. Ecol.* 69:395-406.
- Zhao, J., T. Ni, Y. Li, W. Xiong, W. Ran, B. Shen, Q. Shen, and R. Zhang. 2014. Responses of bacterial communities in arable soils in a rice-wheat cropping system to different fertilizer regimes and sampling times. *PLoS ONE* 9:e85301.